




Transverse Single-Spin Asymmetries and Cross Section of Weak Bosons in p+p collisions at $\sqrt{s} = 510$ GeV

Oleg Eyser

Brookhaven National Laboratory
for the STAR Collaboration



Particles and Nuclei International Conference
September 5-10, 2021

Supported by




QCD, Universality, and the Proton Structure

- Transverse spin asymmetries are an ideal tool to study the multidimensional structure of the nucleon.
 - Spin-orbit correlations
Sivers effect: correlation between proton spin and transverse momentum of partons
 - Non-universality exhibits the process dependence
Attractive color force in SIDIS turns into repulsive force in p+p



O. Eysler / PANIC 2021



Gamberg, Kang, Prokudin
Phys. Rev. Lett. 110, 232301 (2013)
with HERMES data

Prospects for RHIC Run 2017

- Increased theoretical interest
- Significant uncertainty in sea-quark Sivers-TMD function
- Very different Q^2 range in SIDIS, Drell-Yan, and W-boson production
- TMD evolution not the same as DGLAP

from the STAR beam use request

www.bnl.gov/npp/pac.php






Figure 4-1: (left) Prediction for Sivers asymmetry A_N for DY lepton pair production at $\sqrt{s}=500$ GeV, for the invariant mass $4 < Q < 8$ GeV and transverse momenta $0 < q_T < 1$ GeV [19] **before any TMD evolution is applied.** (middle and right) A_N as a function of W^\pm boson rapidity at $\sqrt{s}=500$ GeV, **both before and after TMD evolution is applied.**

- Transverse asymmetries need the full reconstruction of the W-boson kinematics
 - Predicted asymmetries as function of rapidity and transverse momentum
 - Measurement of azimuthal modulation
 - Proof of principle measurement: PRL 116 (2016) 132301



RHIC as a Polarized Proton Collider

$$\vec{p} + \vec{p} / \vec{p} + A \quad \sqrt{s_{NN}} = 200 - 510 \text{ GeV}$$



RHIC as a Polarized Proton Collider


$$\vec{p} + \vec{p} / \vec{p} + A \quad \sqrt{s_{NN}} = 200 - 510 \text{ GeV}$$



Lepton Candidate

- Well established method (helicity asymmetry and cross section ratio measurements)
 - Includes Z^0 and τ decays
 - Data driven QCD normalized at low p_T -range
 - Missing EMC estimated from cuts with / without EMC
- Lepton candidate $p_T > 25$ GeV

Recent STAR W-boson results:
 Phys. Rev. D 103 (2021) 012001
 Phys. Rev. D 99 (2019) 051102



- 2017 data
 - Luminosity leveled (ZDC rate 330 kHz)
 - Barrel EMC high tower trigger
 - StiCA track reconstruction
- Candidate track with matched EMC hit:
 - TPC hits (≥ 15) and fit fraction ($> 51\%$)
 - $EMC E_{2 \times 2} / E_{4 \times 4} > 0.96$
 - $E_{2 \times 2} / E_{R=0.7} > 0.88$
 - $E_{T,away} < 10$ GeV
 - $p_{T,bal} > 0.8 p_T$


W-Boson Reconstruction

$$p + p \rightarrow W^\pm \rightarrow e^\pm + \nu$$

- W-boson decay
 - $p_{T,W}$ is lost
 - Almost no azimuthal angle correlation
- Measure recoil from the collision (tracks and EMC)

$$p_{T,W} = p_{T,e} + p_{T,\nu} = p_{T,recoil}$$

$$p_{T,recoil} = \sum(p_{T,TPC} + E_{T,EMC})$$



W-Boson Reconstruction



- W-boson decay
 - $p_{T,W}$ is lost
 - Almost no azimuthal angle correlation
- Measure recoil from the collision (tracks and EMC)

$$p_{T,W} = p_{T,e} + p_{T,\nu} = p_{T,recoil}$$

$$p_{T,recoil} = \sum(p_{T,TPC} + E_{T,EMC})$$

- Limited barrel acceptance
 - Comparison with embedded simulation
 - Recoil p_T correction
 - $p_{z,\nu}$ is more problematic


$$M_W^2 = (E_e + E_\nu)^2 - (\vec{p}_e + \vec{p}_\nu)^2$$

$$p_{\nu,z} = \frac{A}{p_{e,T}^2} \left[p_{e,z} \pm p_e \cdot \sqrt{1 - \frac{p_{e,T}^2 \cdot p_{\nu,T}^2}{A^2}} \right]$$


$$A = M_W^2 + \vec{p}_{e,T} \cdot \vec{p}_{\nu,T}$$

$$R = 1 - \frac{p_{e,T}^2 \cdot p_{\nu,T}^2}{A^2}$$

$$p_{T,W} = f(p_{T,meas}) = A + \frac{b}{(p_{T,meas})^c}$$



W^-



Azimuthal Angle Smearing

- Transverse spin asymmetries are measured through azimuthal modulations:

$$d\sigma(\phi) = \sigma_0[1 + PA_N \cos(\phi)]$$

$$A_N = \frac{d\sigma(\phi) - d\sigma(\phi + \pi)}{d\sigma(\phi) + d\sigma(\phi + \pi)}$$


$$A_N = \frac{1}{P} \frac{N_\phi - N_{\phi+\pi}}{N_\phi + N_{\phi+\pi}}$$

- Toy Monte Carlo study → determine asymmetry dilution

- 100k MC samples based on input distribution from embedding (per η -bin)

$$D = A_{N,meas}/A_{N,input}$$



$p_{T,recoil}$



$$\Delta\phi = \phi_W - \phi_{recoil}$$

$$\sigma_{A_N} \propto \frac{1}{\sqrt{ND}}$$



O. Eyser / PANIC 2021



$$\text{Asymmetry correction: } A_N = A_{N,meas}/D$$

Transversal Helicity Function g_{1T}

- Transversal helicity can also be measured in W-production
- χ^2 of fit is improved
- Uncertainties in A_S are similar to A_N
- A_S consistent with 0
- Cross talk in A_N is very small
 - W^- : $\Delta A_N / \sigma_{A_N} < 20\%$
 - Included in $\sigma_{syst}(A_N)$



$$\phi_V - \phi_{S_A} = \phi - \pi/2$$

$$A_N: \sin(\phi_V - \phi_{S_A}) = -\cos \phi$$



$$A_S: \cos(\phi_V - \phi_{S_A}) = \sin \phi$$

$$\frac{d\sigma^W}{dy d^2\vec{q}_T} = \sigma_0^W \left\{ F_{UU} + S_{AL}F_{LU} + S_{BL}F_{UL} + S_{AL}S_{BL}F_{LL} \right. \\ + |\vec{S}_{AT}| \left[\sin(\phi_V - \phi_{S_A}) F_{TU}^{\sin(\phi_V - \phi_{S_A})} + \cos(\phi_V - \phi_{S_A}) F_{TU}^{\cos(\phi_V - \phi_{S_A})} \right] \\ + |\vec{S}_{BT}| \left[\sin(\phi_V - \phi_{S_B}) F_{UT}^{\sin(\phi_V - \phi_{S_B})} + \cos(\phi_V - \phi_{S_B}) F_{UT}^{\cos(\phi_V - \phi_{S_B})} \right] \\ + |\vec{S}_{AT}| S_{BL} \left[\sin(\phi_V - \phi_{S_A}) F_{TL}^{\sin(\phi_V - \phi_{S_A})} + \cos(\phi_V - \phi_{S_A}) F_{TL}^{\cos(\phi_V - \phi_{S_A})} \right] \\ + S_{AL} |\vec{S}_{BT}| \left[\sin(\phi_V - \phi_{S_B}) F_{LT}^{\sin(\phi_V - \phi_{S_B})} + \cos(\phi_V - \phi_{S_B}) F_{LT}^{\cos(\phi_V - \phi_{S_B})} \right] \\ + |\vec{S}_{AT}| |\vec{S}_{BT}| \left[\cos(2\phi_V - \phi_{S_A} - \phi_{S_B}) F_{TT}^{\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})} + \cos(\phi_{S_A} - \phi_{S_B}) F_{TT}^1 \right. \\ \left. + \sin(2\phi_V - \phi_{S_A} - \phi_{S_B}) F_{TT}^{\sin(2\phi_V - \phi_{S_A} - \phi_{S_B})} + \sin(\phi_{S_A} - \phi_{S_B}) F_{TT}^2 \right] \left. \right\}.$$

$$F_{TU}^{\sin(\phi_V - \phi_{S_A})} = \mathcal{C}^W \left[(v_q^2 + a_q^2) \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} f_{1T}^1 \bar{f}_1 \right], \\ F_{TU}^{\cos(\phi_V - \phi_{S_A})} = -\mathcal{C}^W \left[2v_q a_q \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} g_{1T} \bar{f}_1 \right],$$

Z. Kang, CFNS Workshop on
RHIC physics for EIC, May 2021

Results: $A_N(W^\pm)$





- Comparison with new theory prediction, based on first global fit of world data
 - Updated for STAR kinematics from PRL 126 (2021) 112002
- New STAR data will have biggest impact on high-x region of quark Sivers function

New Results for Z^0

$$p + p \rightarrow Z^0 \rightarrow e^+ + e^-$$


- Experimentally very clean
 - Two high- p_T electrons (e^+ , e^-) from same vertex
- Leading systematic uncertainty from energy resolution
- Comparison with PRL 126 (2021) 112002
(more details in arxiv:2103.03270)




Unpolarized TMDs

$$p + p \rightarrow Z^0 \rightarrow e^+ + e^-$$

- Differential cross section of high interest for TMD-PDF fits
 - Pavia group, *JHEP* 07 (2020) 117



- 2017 data doubles the previous statistics
- Unfolded p_T spectrum
- Systematics from energy resolution and electron selection






Global luminosity uncertainty 8.5%
not included in the plot

Summary

➤ New results of transverse single-spin asymmetries of W^\pm/Z^0 -bosons

- Much improved precision over PRL 116 (2016) 132301
- Corrected for smeared recoil reconstruction
- Expect big impact in Sivers function at high- x in next global TMD fit



➤ Differential cross section of Z^0 -bosons

- Now 700 pb^{-1} of integrated luminosity



$$-\vec{p}_{rec} = \vec{p}_W = \vec{p}_e + \vec{p}_\nu$$

W^+

W^-

W-boson









MC true

recoil

W-boson

MC true

recoil



$\Delta = e - W$

$\Delta = e - rec$




O. Eyer / PANIC 2021

$\Delta = e - W$

$\Delta = e - rec$

Lepton Candidate Selection

- 2017 data
 - Luminosity leveled (ZDC rate 330 kHz)
 - Barrel EMC high tower trigger
 - StiCA track reconstruction
- Candidate track with matched EMC hit:
 - TPC hits (≥ 15) and fit fraction ($> 51\%$)
 - EMC $E_{2 \times 2} / E_{4 \times 4} > 0.96$
 - $E_{2 \times 2} / E_{R=0.7} > 0.88$
 - $E_{T,away} < 10 \text{ GeV}$
 - $p_{T,bal} > 0.8 p_T$



Electron Pseudorapidity

Compare Jae Nam's analysis:
W cross section ratio in run 17





$\eta_e < -0.5$

$-0.5 < \eta_e < 0$





$0 < \eta_e < 0.5$

$0.5 < \eta_e$

W^+



W^-



Azimuthal Angle Reconstruction


- Transverse spin asymmetries are measured through azimuthal modulations:

$$d\sigma(\phi) = \sigma_0 [1 + PA_N \cos(\phi)]$$

$$A_N = \frac{d\sigma(\phi) - d\sigma(\phi + \pi)}{d\sigma(\phi) + d\sigma(\phi + \pi)}$$

$$A_N = \frac{1}{P} \frac{N_\phi - N_{\phi+\pi}}{N_\phi + N_{\phi+\pi}}$$

- Toy Monte Carlo study → dilution factor $D = A_{N,meas}/A_{N,input}$




$$r_{BG} = \frac{BG}{S + BG}$$


$$S = \int f(x) dx = \sqrt{2\pi} \cdot p_2$$

$$BG = 2\pi p_3$$

O. Eyser / PANIC 2021



W-Bosons & Binning



Global Fit

- Original release: 2002.08384, Cammarota et al.
- Phys Rev. D102, 054002 (2020)
2009.10710, Echevarria et al.
- PRL 126, 112002 (2021)
2012.05135, Bury et al.
- 2103.03270, Bury et al.

