



JOINT MARCH MEETING AND APRIL MEETING  
Global Physics Summit

March 16–21, 2025, Anaheim, CA and virtual



# Probing Baryon Transport Dynamics and Strangeness Production with Hyperon-kaon Correlations

Siyuan Ping

Fudan University

# Outline

1. Baryon Number Transport

2. Hyperon-Kaon Correlation

3. p(d) + Au Collision



4. AMPT Result

5. Summary and Outlook

# Baryon Number Transport

- Baryons number transport from  $|y_{Beam}|$  to  $y \sim 0$ 
  - Significant  $N_{Baryon} > N_{anti-Baryon}$  at mid-rapidity
- Fragmentation scheme
  - Valence quark picture
  - Gluon-junction model


STAR Collaboration, arXiv:2408.15441



I. G. Bearden et al. (BRAHMS), Phys. Rev.L, 2004, 93(10):102301


# Hyperon-Kaon Correlation

- Baryons number transport from Proton/Neutron to Hyperon
  - Ratio Anti-Hyperon/Hyperon < 1
  - Strangeness conservation: Strange – Anti-Strange quarks are pair-produced




## Kaon-Hyperon Correlation:

*particles containing at least one valence strange quark:  
More primordial production than from weak decays  
compare to  $\pi$  and  $p$*



# p(d) + Au Collision

- Asymmetric collisions
  - The rapidity shift depends on the target A thickness
  - Baryon number is carried by the gluon junction during the interaction
  - Valence quark jets emerge with leading mesons
  - Hyperons in initial p/d hemi-sphere more likely carry baryons number from initial p/d
  - Clearer K-Hyperon correlation for hyperon which  $y > 0$



$$C_{K\Lambda}(\Delta y) = \frac{\bar{S}S}{N_{\Lambda}} \frac{dN_{K^+\Lambda}^{pairs}}{d\Delta y} - \frac{\bar{S}\bar{S}}{N_{\bar{\Lambda}}} \frac{dN_{K^+\bar{\Lambda}}^{pairs}}{d\Delta y}$$


$\Delta y = \theta(y_{\Lambda})(y_K - y_{\Lambda}) + \theta(-y_{\Lambda})(y_{\Lambda} - y_K)$   
 $\Delta y > 0$ : Kaon emit faster;  $\Delta y < 0$ :  $\Lambda$  emit faster

Yielding information of average Kaon number per Lambda  
 Combinatorial background subtraction

# AMPT Result

- Simulation for reference:
  - A Multi-Phase Transport Model (AMPT)
  - Version: 2.25t7cu (StringMelting)
  - Transport models treat chemical and thermal freeze-out dynamically
- Data Generated:
  - p+Au@39/62GeV
  - d+Au@200GeV
- Guiding the experimental investigation
  - STAR acceptance adopted
    - $|y| < 1$
    - $|\eta| < 1$

*Structure of AMPT model with string melting*



Z.W. Lin, C.M. Ko, B.A. Li, B. Zhang, S. Pal, Phys.Rev.C72:064901,2005

# AMPT Result

- p+Au@39GeV


$$\Delta y = \theta(y_{Hyperon})(y_K - y_{Hyperon}) + \theta(-y_{Hyperon})(y_{Hyperon} - y_K)$$

$\Delta y > 0$ : Kaon emit faster;  $\Delta y < 0$ : Hyperon emit faster


$$C_{K\Lambda}(\Delta y) = \frac{1}{N_\Lambda} \frac{dN_{K^+\Lambda}^{pairs}}{d\Delta y} - \frac{1}{N_{\bar{\Lambda}}} \frac{dN_{K^+\bar{\Lambda}}^{pairs}}{d\Delta y}$$

$$C_{K\Sigma}(\Delta y) = \frac{1}{N_\Sigma} \frac{dN_{K^+\Sigma}^{pairs}}{d\Delta y} - \frac{1}{N_{\bar{\Sigma}}} \frac{dN_{K^+\bar{\Sigma}}^{pairs}}{d\Delta y}$$


$$C_{K\Omega}(\Delta y) = \frac{1}{N_\Omega} \frac{dN_{K^+\Omega}^{pairs}}{d\Delta y} - \frac{1}{N_{\bar{\Omega}}} \frac{dN_{K^+\bar{\Omega}}^{pairs}}{d\Delta y}$$



$$\int C_{K\Lambda}(\Delta y) d\Delta y = 0.051 \pm 0.001$$



$$\int C_{K\Sigma}(\Delta y) d\Delta y = 0.104 \pm 0.002$$



$$\int C_{K\Omega}(\Delta y) d\Delta y = 0.152 \pm 0.007$$

# AMPT Result

- p+Au@62GeV


$$\Delta y = \theta(y_{\text{Hyperon}})(y_K - y_{\text{Hyperon}}) + \theta(-y_{\text{Hyperon}})(y_{\text{Hyperon}} - y_K)$$

$\Delta y > 0$ : Kaon emit faster;  $\Delta y < 0$ : Hyperon emit faster


$$C_{K\Lambda}(\Delta y) = \frac{1}{N_\Lambda} \frac{dN_{K^+\Lambda}^{\text{pairs}}}{d\Delta y} - \frac{1}{N_{\bar{\Lambda}}} \frac{dN_{K^+\bar{\Lambda}}^{\text{pairs}}}{d\Delta y}$$

$$C_{K\Sigma}(\Delta y) = \frac{1}{N_\Sigma} \frac{dN_{K^+\Sigma}^{\text{pairs}}}{d\Delta y} - \frac{1}{N_{\bar{\Sigma}}} \frac{dN_{K^+\bar{\Sigma}}^{\text{pairs}}}{d\Delta y}$$


$$C_{K\Omega}(\Delta y) = \frac{1}{N_\Omega} \frac{dN_{K^+\Omega}^{\text{pairs}}}{d\Delta y} - \frac{1}{N_{\bar{\Omega}}} \frac{dN_{K^+\bar{\Omega}}^{\text{pairs}}}{d\Delta y}$$



$$\int C_{K\Lambda}(\Delta y) d\Delta y = 0.042 \pm 0.001$$



$$\int C_{K\Sigma}(\Delta y) d\Delta y = 0.093 \pm 0.002$$



$$\int C_{K\Omega}(\Delta y) d\Delta y = 0.138 \pm 0.006$$



# AMPT Result

- d+Au@200GeV


$$\Delta y = \theta(y_{\text{Hyperon}})(y_K - y_{\text{Hyperon}}) + \theta(-y_{\text{Hyperon}})(y_{\text{Hyperon}} - y_K)$$

$\Delta y > 0$ : Kaon emit faster;  $\Delta y < 0$ : Hyperon emit faster


$$C_{K\Lambda}(\Delta y) = \frac{1}{N_\Lambda} \frac{dN_{K^+\Lambda}^{\text{pairs}}}{d\Delta y} - \frac{1}{N_{\bar{\Lambda}}} \frac{dN_{K^+\bar{\Lambda}}^{\text{pairs}}}{d\Delta y}$$

$$C_{K\Sigma}(\Delta y) = \frac{1}{N_\Sigma} \frac{dN_{K^+\Sigma}^{\text{pairs}}}{d\Delta y} - \frac{1}{N_{\bar{\Sigma}}} \frac{dN_{K^+\bar{\Sigma}}^{\text{pairs}}}{d\Delta y}$$


$$C_{K\Omega}(\Delta y) = \frac{1}{N_\Omega} \frac{dN_{K^+\Omega}^{\text{pairs}}}{d\Delta y} - \frac{1}{N_{\bar{\Omega}}} \frac{dN_{K^+\bar{\Omega}}^{\text{pairs}}}{d\Delta y}$$



$$\int C_{K\Lambda}(\Delta y) d\Delta y = 0.037 \pm 0.001$$




$$\int C_{K\Sigma}(\Delta y) d\Delta y = 0.099 \pm 0.002$$




$$\int C_{K\Omega}(\Delta y) d\Delta y = 0.166 \pm 0.007$$

# AMPT Result


- d+Au@200GeV Full Acceptance, estimate the detector acceptance effects in preparation for STAR data analysis



$$\int C_{K\Lambda}(\Delta y) d\Delta y = 0.174 \pm 0.000$$



$$\int C_{K\Xi}(\Delta y) d\Delta y = 0.417 \pm 0.001$$



$$\int C_{K\Omega}(\Delta y) d\Delta y = 0.619 \pm 0.005$$

STAR Acceptance:  $\int C_{K\Lambda}(\Delta y) d\Delta y = 0.037 \pm 0.001$

$\int C_{K\Xi}(\Delta y) d\Delta y = 0.099 \pm 0.002$

$\int C_{K\Omega}(\Delta y) d\Delta y = 0.166 \pm 0.007$

# Summary and Outlook


- p(d)+A collision:
  - A new way to study possible baryon junction interactions
- Hyperon-Kaon correlation:
  - Sensitive probe of baryon number transport dynamics
- AMPT simulation:
  - StringMelting version
  - Hyperons with positive rapidity correlate faster emitted Kaons
    - A hint of baryon junction?
  - $\int C_{K-Hyperon}(\Delta y) d\Delta y : K\Omega > K\Xi > K\Lambda$ 
    - A hint of baryon number transport to hyperons
- Analysis of d+Au STAR dataset is ongoing (Stay tuned!)

Thank you!


# Back Up

# AMPT Result


- p+Au@39GeV Full Acceptance



$$\int C_{K\Lambda}(\Delta y) d\Delta y = 0.172 \pm 0.000$$



$$\int C_{K\Xi}(\Delta y) d\Delta y = 0.373 \pm 0.001$$



$$\int C_{K\Omega}(\Delta y) d\Delta y = 0.536 \pm 0.003$$




STAR Acceptance:  $\int C_{K\Lambda}(\Delta y) d\Delta y = 0.051 \pm 0.001$

$$\int C_{K\Xi}(\Delta y) d\Delta y = 0.104 \pm 0.002$$

$$\int C_{K\Omega}(\Delta y) d\Delta y = 0.152 \pm 0.007$$

# AMPT Result

- p+Au@62GeV Full Acceptance



STAR Acceptance:  $\int C_{K\Lambda}(\Delta y) d\Delta y = 0.042 \pm 0.001$ 
                 
  $\int C_{K\Xi}(\Delta y) d\Delta y = 0.093 \pm 0.002$ 
                 
  $\int C_{K\Omega}(\Delta y) d\Delta y = 0.138 \pm 0.006$

