

1 Longitudinal De-correlation of Anisotropic Flow 2 at RHIC-STAR

3 Gaoguo Yan (for the STAR Collaboration)
4 Shandong University

5 Studies of longitudinal de-correlation of anisotropic flow can provide unique con-
6 straints on the three-dimensional structure of the initial stages and dynamical
7 evolution of the quark-gluon-plasma in heavy-ion collisions. Experimentally,
8 the factorization ratio, $r_n(\eta)(n = 2, 3)$, is used to quantify the amount of the
9 longitudinal flow de-correlation with pseudorapidity [1-3]. With data collected
10 by the STAR experiment at RHIC for 200 GeV Ru+Ru and Zr+Zr collisions as
11 well as 19.6, 27, and 54.4 GeV Au+Au collisions, we will present the centrality
12 and collision energy dependence of $r_n(\eta)(n = 2, 3)$. These results can provide
13 new insights into the three-dimensional modeling of the relativistic heavy-ion
14 collisions.

15 [1] The CMS Collaboration, Phys. Rev. C 92, 034911 (2015).

16 [2] The ATLAS Collaboration, Eur. Phys. J. C 78, 142 (2018).

17 [3] The ATLAS Collaboration, Phys. Rev. Lett. 126, 122301 (2021).