

STAR Results on Transversivity and TMD-Related Observables

Bassam Aboona, on behalf of the STAR Collaboration
(he/him/his)

bem4r@physics.tamu.edu

June 3rd -7th, 2024

Supported in part by:




U.S. DEPARTMENT OF
ENERGY

Outline


- Introduction
- Sivers Effect
- Transversity and the Collins Effect
- Selected Future Work
- Summary

RHIC: Relativistic Heavy Ion Collider




- The only machine in the world capable of colliding high-energy beams of polarized protons
- The beams travel in opposite directions around RHIC's 3.86 km two-lane racetrack
- Offers a wide range of center-of-mass energies (up to 510 GeV)

STAR: Solenoidal Tracker At RHIC



STAR, Phys. Rev. D **106**, 072010 (2022)






- Roman Pots (RP) at 15 meters from the interaction point
- FMS used to be where the current ECal and HCal are and covered $2.6 < \eta < 4.2$

Similar x coverage,
but at higher Q^2
when compared to
SIDIS

Transverse Single-Spin Asymmetries (TSSA's) - A_N


- Since the 1970's, surprisingly, large TSSA's have been observed at forward rapidities in $p^\uparrow + p$ collisions
- Perturbative Quantum Chromodynamics (pQCD) predicts very small values for A_N
- Twist-3 and **transverse momentum dependent (TMD)** theoretical frameworks have been developed to describe this observed large TSSA

$$A_N = \frac{N_L - N_R}{N_L + N_R}$$



Plot reference: Elke Aschenauer et al., arXiv:1602.03922.

Sivers Effect: A Mechanism for A_N



- \mathbf{k}_T : parton transverse momentum
- \mathbf{S}_p : proton spin
- \mathbf{P} : proton momentum


- $f_{1T}^\perp(x, k_T, Q^2)$: Describes the relationship between the transverse momentum distribution of unpolarized partons and the transverse spin polarization of the proton [1].
- Characterizes a scalar triple-vector correlation for an unpolarized parton and its polarized parent proton.
- Correlation between partonic orbital motion and proton's spin

[1] D. Sivers, Phys. Rev. D **41**, 83 (1990).


Probing The Sivers Effect Using Dijet Production

- A transversely polarized proton going in the longitudinal direction can have partons with a spin-dependent transverse momentum k_T
- The k_T provides a kick to the dijet and makes it fold in the direction of the transverse momentum

Side view:




Beam's eye view:




Probing The Sivers Effect Using Dijet Production

- A transversely polarized proton going in the longitudinal direction can have partons with a spin-dependent transverse momentum k_T
- The k_T provides a kick to the dijet and makes it fold in the direction of the transverse momentum

Side view:




Beam's eye view:



Mean k_T Flavor Dependence

STAR, arXiv:2305.10359




For the first time, there is evidence of non-zero Sivers effect using dijets.

- Jet charge tagging combined with unfolding used to determine the quark flavor.
- Tagged $\langle k_T \rangle$ represents a mixture of partons
- Obtaining parton fractions from simulation allows for measuring the individual parton $\langle k_T \rangle$

Results:

- $d\text{-quark } \langle k_T \rangle \approx -2 \text{ } u\text{-quark } \langle k_T \rangle$
- The $\langle k_T \rangle$ for gluon and sea quarks combined is consistent with zero

W^\pm Boson Reconstruction and A_N



Use TPC tracks and EMC hits to measure W^\pm recoil from collision:

$$\vec{p}_{T,W} = \vec{p}_{T,e} + \vec{p}_{T,\nu} = -\vec{p}_{T,recoil}$$



$$\vec{p}_{T,recoil} = \sum (\vec{p}_{T,TPC} + \vec{E}_{T,EMC})$$

- Left and right asymmetry of the W^\pm production with respect to the spin of the polarized proton
- Sensitive to the Sivers TMD function – $Q^2 = M_W^2 \sim 6500 \text{ GeV}^2$
- A_N is measured via azimuthal angle

$$A_N \cdot \cos(\phi) = \frac{1}{\langle P \rangle} \cdot \frac{\sqrt{N_\uparrow(\phi)N_\downarrow(\phi + \pi)} - \sqrt{N_\uparrow(\phi + \pi)N_\downarrow(\phi)}}{\sqrt{N_\uparrow(\phi)N_\downarrow(\phi + \pi)} + \sqrt{N_\uparrow(\phi + \pi)N_\downarrow(\phi)}}$$

$\langle P \rangle$: Mean beam polarization
 $N_\uparrow(N_\downarrow)$: Yield in spin up (down) state
 ϕ : Azimuthal angle


A_N Preliminary Results of W^\pm




- Results are generally consistent with the model predictions
- STAR results will have biggest impact on high- x region of the quark Sivers function

Z^0/γ^* Cross Section

STAR, *Phys. Lett. B* **854** (2024) 138715




- Z^0 events are reconstructed via:



- Serves as a test of the universality of unpolarized TMDs
- Provides insights into the x and Q^2 evolution of unpolarized TMDs
 - RHIC energies provide access to higher x compared to the Tevatron and LHC

$Z^0/\gamma^* A_N$


STAR, *Phys. Lett. B* **854** (2024) 138715



- $p_T^{Z^0}$ is limited to $< 10 \text{ GeV}/c$ to stay within the kinematic region where the polarized TMD approach is applicable
- This result will allow for the extraction of the Sivers TMD PDF, and especially for valence quarks in the region $x \geq 0.1$
- Unable to provide a conclusive statement regarding the sign-change hypothesis of the Sivers function

$$\text{Sivers}_{\text{DIS}} = -\text{Sivers}_{\text{DY}} \text{ or } \text{Sivers}_{W^\pm, Z^0}$$

Collins Effect: A Mechanism for A_N




- The relationship between the leading-twist (twist-2) collinear transversity, $h_1^a(x_1, Q^2)$, and the TMD Collins fragmentation function, $H_{1\pi/c}^\perp(z_\pi, j_T, Q^2)$

$$\frac{d\sigma^\uparrow(\phi_S, \phi_H) - d\sigma^\downarrow(\phi_S, \phi_H)}{d\sigma^\uparrow(\phi_S, \phi_H) + d\sigma^\downarrow(\phi_S, \phi_H)}$$

$$\propto A_{UT}^{\sin(\phi_S)} \sin(\phi_S) + A_{UT}^{\sin(\phi_S - \phi_H)} \sin(\phi_S - \phi_H)$$

$$\sim h_1^a(x_1, Q^2) \otimes H_{1\pi/c}^\perp(z, j_T, Q^2) + \dots$$


z_π : longitudinal momentum fraction of the pion




J. C. Collins, Nucl. Phys. B **396**, 161 (1993).
 Z.-B. Kang *et al.*, JHEP **11**, 068 (2017).
 Z.-B. Kang *et al.*, Phys. Lett. B **774**, 635 (2017).
 U. D'Aesio *et al.*, Phys. Rev. D **83**, 034021 (2011).

π^\pm Collins Asymmetry at $\sqrt{s} = 200$ GeV

- Integrated over a wide range of z and j_T to provide sensitivity to the collinear transversity, $h_1^a(x, Q^2)$
- The hadron j_T and z binning allows sensitivity to the Collins FF, $H_{1\pi/c}^\perp(z_\pi, j_T, Q^2)$
- In general, model calculations underestimate experimental data




STAR, Phys. Rev. D **106**, 072010 (2022).





π^\pm Collins Asymmetry at $\sqrt{s} = 200$ and 510 GeV

- Results from the two beam energies match each other very well
- Little, if any, energy dependence when comparing the 200 GeV results to the 510 GeV results
 - Q^2 values differ by a factor of 6 between 200 GeV and 510 GeV results
- Sets constraints on evolution effects



π^\pm Collins Asymmetry at $\sqrt{s} = 200$ and 510 GeV






z : longitudinal momentum fraction of the pion

j_T : transverse momentum of the pion with respect to the jet axis

- z and j_T binning allows sensitivity to the Collins FF, $H_{1\pi/c}^\perp(z_\pi, j_T, Q^2)$
- Good agreement between the 200 and 510 GeV results
- Little to no energy dependence

Di-pion Asymmetries and Cross-Section Measurements



$$A_{UT} \propto \frac{h_1^a(x) \otimes H_1^{\chi \pi^+ \pi^-}(z, M_h^2)}{f_1^a \otimes D_1^{\pi^+ \pi^-}}$$

- New measurements of A_{UT} at 200 and 510 GeV
 - First measurement of unpolarized $\pi^+\pi^-$ cross section at 200 GeV
- (a) + (b) \rightarrow model independent extraction of $h_1^q(x)$

Learn more during Bernd Surrow's talk this afternoon!

Λ and $\bar{\Lambda}$ Hyperon Transverse Spin Transfer - D_{TT}


$$D_{TT}^{\Lambda} = \frac{d\sigma(p^{\uparrow}p \rightarrow \Lambda^{\uparrow}X) - d\sigma(p^{\uparrow}p \rightarrow \Lambda^{\downarrow}X)}{d\sigma(p^{\uparrow}p \rightarrow \Lambda^{\uparrow}X) + d\sigma(p^{\uparrow}p \rightarrow \Lambda^{\downarrow}X)} = \frac{d\delta\sigma^{\Lambda}}{d\sigma^{\Lambda}}$$

$$d\delta\sigma^{\Lambda} = \sum \int dx_a dx_b dz \delta f_a(x_a) f_b(x_b) \delta\sigma(ab \rightarrow cd) \delta D^{\Lambda}(z)$$

transversity pQCD calculable polarized FF


- $\Lambda(\bar{\Lambda}) D_{TT}$ is sensitive to the (anti-)strange quark transversity in the proton
- Λ and $\bar{\Lambda}$ results are consistent with each other within uncertainties
- D_{TT} is consistent with zero

STAR, Phys. Rev. D **109**, 012004 (2024).




Theory: Q. H. Xu *et al.*, Phys. Rev. D, **73**(7), 077503 (2006).


Previous STAR Results



Theory: J. Cammarota *et al.*, Phys. Rev. D **102**, 054002 (2020).



Theory: Z.-B. Kang *et al.*, Phys. Lett. B **774**, 635 (2017).



Theory: L. Gamberg *et al.*, Phys. Rev. Lett. **110**, 232301 (2013).

Observations and Measurements: (STAR) J. Adam *et al.*, Phys. Rev. D **103**, 092009 (2021).

- Small A_N is observed for non-isolated π^0 in EM-jets (non-isolated = other photons are allowed in the jet)
- Small Collins asymmetry for π^0 in EM-jet
- Small jet A_N for inclusive EM-jets
- For $x_F \lesssim 0.3$: DIS-based model for the Sivers effect describes the non-isolated π^0 results well, but not the isolated results
- Large A_N is observed for isolated π^0 in EM-jets (isolated = no other photons in the jet)

Conclusion


Collins effect can't account for the large TSSA

Sivers effect can't account for large TSSA

Diffractive process?

Relevant Event Classes

Inclusive EM-Jet Event:




Relevant Event Classes

Rapidity Gap (RG) Event:

Vetoing hadrons in the BBC η range suppresses a large fraction of the non-diffractive events – RG events are highly enriched in diffractive processes

East BBC: $-5 < \eta < -2$




Relevant Event Classes


Single Diffractive Event:

Intact proton has an $\xi < 0.15$

East roman pots: $\eta < -6$



East BBC: $-5 < \eta < -2$



FMS: $2.5 < \eta < 4$


EM Jet

West

East

Forward Rapidity: A_N for Inclusive EM-Jets

- EM-jets are reconstructed using only photons
- Photon candidates are obtained from the Forward Meson Spectrometer (FMS) on the west side of STAR



- Three different photon multiplicity scenarios are considered
 - Multiplicity dependence is observed
- EM-jets with only 1 or 2 photons have the largest A_N
 - Could this point to a contribution to the observed A_N from diffractive processes?

Diffraction Process and A_N

$p + p \rightarrow \text{EM-jet} + X$

Inclusive EM-jet


$p + p \rightarrow \text{EM-jet} + X$

RG events (at least 50% of RG events are single diffractive)

$p + p \rightarrow p + \text{EM-jet} + X$


Single diffractive process

- A_N consistent within uncertainties for all three processes
- If A_N has significant contributions from diffractive processes, then A_N from diffractive events is expected to have a large magnitude
- Current results do not provide evidence in favor of a diffractive process having a large contribution to A_N




Mid-Rapidity: Inclusive Jet Asymmetry at 200 and 510 GeV

- At low p_T , the inclusive jet asymmetry is sensitive to the twist-3 correlators associated with the gluon Sivers function
- 510 GeV results extend the measurement to lower values of x
- Results are consistent with zero within uncertainties




Mid-Rapidity: Pion Tagged Jet Asymmetry at 200 and 510 GeV

STAR, Phys. Rev. D **106**, 072010 (2022)



u (for π^+) and d (for π^-) quark functions are enhanced by performing the pion tagging, providing sensitivity to the twist-3 correlators associated with the quark Sivers function




KPRY: Z.-B. Kang, A. Prokudin, F. Ringer, and F. Yuan, Phys. Lett. B **774**, 635 (2017), arXiv:1707.00913

Results are consistent with zero within uncertainties


Outlook

STAR Forward Upgrade:

- Installed and commissioned before 2022
- $2.5 < \eta < 4$
- Charged particle tracking using Si detectors and small-strip Thin Gap Chambers (sTGC)
- Electromagnetic and hadronic calorimetry
- Capable of measuring:
 - h^\pm, e^\pm (with good e/h discrimination)
 - Photons, π^0
 - Jets, hadrons in jets
 - Lambda's
 - Drell-Yan and J/ψ di-electrons
 - Mid-forward and forward-forward correlations
- Quarks up to $x \sim 0.5$ and gluons down to $x \sim 0.001$




Outlook



- The mid + forward rapidity capabilities of STAR complement the future EIC kinematic coverage
- The forward upgrade will bridge the kinematic region between mid-rapidity STAR and SIDIS
 - great for future Collins measurements

Outlook




- A_N for full jet reconstruction, combined with charge-sign tagging of a hadron fragment with $z > 0.5$
- Up to 10σ separation between plus-tagged and minus-tagged jet A_N

Summary

- EM-jet A_N results at forward rapidity for single diffractive processes show no large contribution for the observed large TSSA in the forward direction
- Spin-dependent $\langle k_T \rangle$ from dijet production and A_N from W^\pm/Z^0 studies at STAR provide probes for the Sivers effect
- The Z^0 cross section gives insights into the evolution of the unpolarized TMDs
- The Collins effect is studied at two energy levels and show little to no energy dependence
- Di-pion asymmetries and cross-section results from STAR can provide the initial steps to model-independent transversity extractions
- $\Lambda(\bar{\Lambda}) D_{TT}$ is sensitive to the (anti-)strange quark transversity in the proton
- The Forward Upgrade extends the kinematic range of the measurements at STAR, which are essential for universality studies at the future EIC

Backup

Probing The Sivers Effect Using Dijet Production



- φ_b : dijet bisector angle
- $\zeta > \pi$ if $\cos(\varphi_b) > 0$
- $\zeta < \pi$ if $\cos(\varphi_b) < 0$

- The signed opening angle, ζ , is sensitive to the spin-dependent partonic k_T involved in characterizing the Sivers effect.
- A Conversion from the spin-dependent ζ asymmetries ($\Delta\zeta$) to Sivers $\langle k_T \rangle$ can be achieved

$$\Delta\zeta = \frac{\langle \zeta \rangle^+ - \langle \zeta \rangle^-}{P}$$


$\langle \zeta \rangle^\pm$: the centroid of the distribution for spin-up/spin-down proton beams

P : magnitude of beam polarization

Tagged $\Delta\zeta$ and $\langle k_T \rangle$ From Tagged Dijet Production


STAR, arXiv:2305.10359

- $\sim 3.1\sigma$ separation between + and - tagging
- Asymmetry shifts from positive to negative when going from + to - tagging \rightarrow strong evidence that Sivers $\langle k_T \rangle$ in u and d are opposite




$\Delta\zeta - \langle k_T \rangle$
Correlation:

correct jet p_T
to parton level




map $\Delta\zeta - \langle k_T \rangle$
correlation

$$\eta^{\text{total}} = \eta_1 + \eta_2 \propto \ln(x_1/x_2)$$




- +: enhancing the fraction of u
- 0^+ : less enhancement in the u fraction
- 0^- : less enhancement in the d fraction
- : enhancing the fraction of d

Collins Asymmetry vs. z from 510 GeV



Collins Asymmetry from 510 GeV vs. Theory



Theory curves:

- **KPRY:** Z.-B. Kang, A. Prokudin, F. Ringer, and F. Yuan, Phys. Lett. B **774**, 635 (2017), arXiv:1707.00913
- **DMP+2013:** U. D'Alesio, F. Murgia, and C. Pisano, Phys. Lett. B **773**, 300 (2017), arXiv:1707.00914

- The 2011 and 2017 experimental results for A_{UT} agree with each other
- Overall, the theoretical models underestimate the experimental results

Detailed Future Work

- **EM-Jet A_N :**
 - Data from Run 2022 and 2024 using the Forward Upgrade will improve precision of measurement
- **Dijet Sivers:**
 - Combining existing results with data from 2017 and 2022 at 510 and 508 GeV, respectively, to explore the x -dependence of the measurement
- **W^\pm and Z^0 A_N :**
 - STAR recorded 400 pb⁻¹ during Run 2022 utilizing the Forward Upgrade detectors
 - iTPC extends the η coverage
- **Collins Asymmetries:**
 - Use polarized $p + Au$ data from 2015 to measure the Collins asymmetry
 - Use 2022 and 2024 data with the Forward Upgrade for Collins measurements in the forward direction
- **Di-pion Asymmetries:**
 - Use data from Run 2022 and 2024 to perform a precision measurement of IFF asymmetries of pion and kaons