

Determining π^0 A_{LL} from STAR 2012 Endcap Calorimeter Data

Claire Kovarik

For an Undergraduate CEU Poster Presentation

Valparaiso University

(for the STAR Collaboration)

The Solenoidal Tracker at RHIC (STAR) located at Brookhaven National Laboratory in New York uses longitudinally polarized proton-proton collisions to determine the gluon contribution to the spin of the proton. One analysis of the 2012 data set, at a proton-proton center of mass energy of 510 GeV, studies the production of neutral pions (π^0) which immediately ($\sim 10^{-16}$ s) decay into two photons. The neutral pion asymmetry, A_{LL} , can be determined through data collected by the Endcap Electromagnetic Calorimeter (EEMC). The EEMC, positioned in an intermediate pseudorapidity range of $1 < \eta < 2$, is able to measure the energy and position of each photon's electromagnetic shower. By using this information as well as the angle between the photons, the two-photon invariant mass can be reconstructed. The invariant mass spectra are fitted using a skewed Gaussian plus a background function to determine the total number of π^0 s present. The π^0 asymmetry is calculated from the number of π^0 s in collisions with different polarization directions of the colliding proton beams. The status of the measurement of A_{LL} of the π^0 s as a function of the transverse momentum, p_T , of the π^0 will be presented.