



ϕ -meson Production in Au+Au Collisions at $\sqrt{s_{NN}} = 3.0$ GeV from STAR


Guannan XIE (for the STAR Collaboration)
Lawrence Berkeley National Laboratory
Oct.29 – Nov. 1, 2020
DNP 2020, Virtual Meeting

Supported in part by



U.S. DEPARTMENT OF
ENERGY


Office of
Science





Motivation

- ϕ meson ($s\bar{s}$) has a small hadronic cross section.
- The strange hadron yield and ratios may be sensitive to the strangeness production mechanism.
- Grand canonical ensemble (GCE) and canonical ensemble (CE) calculations are quite different at low energy.



HADES:

Phys. Lett. B 778, 2018.403-407,

Phys. Rev. C. 80.025209. (2009)

E917: Phys. Rev. C. 69.054901 (2004)

P. Braun-Munzinger:


Nucl. Phys. A 772, 167 (2006)

K. Redlich:

Phys. Lett. B 603, 146 (2004)



FXT Setup @ STAR




Conventions: beam-going direction is the positive direction



Analysis Procedure


- 2018 FXT 3.85GeV ($\sqrt{s_{NN}} = 3.0$ GeV) $\sim 250M$, $y_{mid} \sim 1.05$.
- $\phi \rightarrow K^+ + K^-$ (hadronic channel).
- TPC (de/dx) and TOF (β) for Kaon particle identification.
- ϕ meson signal was fitted with breit-wigner function.
- Wide p_T and centrality coverage for ϕ meson.






Efficiency Correction

- Tracking efficiency and acceptance effects are estimated with GEANT simulations embedded into real events.
- A data-driven method is used to extract tof matching and pid related efficiency.





Efficiency Corrected m_T Spectra




ϕ -meson and K^- invariant yields in 0-10% most central collisions as a function of transverse kinetic energy ($m_T - m_0$) for various rapidity regions. Yields are fit with an exponential.



Systematic Estimation

STAR: Phys. Rev. C. 034909 (2009)



dN/dp_T extrapolation with various functions to capture the unmeasured p_T range: (systematic source)


- Levy
- Blast-wave
- m_T exponential
- $p_T^{3/2}$ exponential
- p_T^2 Gaussian

Other systematic sources:

Tracking quality cuts and PID.




Yield : Rapidity Distribution



Rapidity distribution of K^- and ϕ -meson and the Gaussian extrapolation in y_{cm} for various centrality regions. Solid symbols are measured data, open ones are reflection.



ϕ/K^- Ratio



HADES:

Phys. Lett. B 778, 2018.403-407, *Phys. Rev. C*. 80.025209. (2009)

E917: *Phys. Rev. C*. 69.054901 (2004)

P. Braun-Munzinger:

Nucl. Phys. A 772, 167 (2006)

K. Redlich:

Phys. Lett. B 603, 146 (2004)

r_c : correlation length, radius of the volume inside which the production of particles with open strangeness is canonically conserved.

ϕ/K^- ratio as a function of center of mass energy $\sqrt{s_{NN}}$ compared with statistical calculations with different r_c parameters. $\sim 5\sigma$ deviated from zero for 0-10% central and 10-40% central collisions.



Summary and Outlook

- First measurements of ϕ -meson production in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV with energy just above the NN threshold*. * A.I. Titov: *Eur. Phys. J. A* 7 (2000) 543-557
- ϕ/K^- ratio deviates from zero with $\sim 5\sigma$ for 0–10% and 10-40% central collisions.
- Data favors the CE with strangeness correlation length ($r_c \sim 3.2$ fm), while GCE show a clear discrepancy at low energy.

Outlook

- Precise measurements of ϕ/K^- on the centrality dependence from the STAR BES-II, to constrain the model calculations.

