



MICHIGAN STATE  
UNIVERSITY

# Local Parity Violation or Local Charge Conservation/Flow?

A Reaction-Plane-Dependent Balance Function Study





Hui Wang  
for the STAR Collaboration

# Motivation

- In heavy ion collisions, most of the detected charge is created during the evolution of the system.
- Balance functions are sensitive to charge formation mechanisms and relative diffusion
- A **three point correlator** has been proposed to measure the possible Chiral Magnet Effect


Phys. Rev. Lett.103,251601 (2009)



$$\gamma_{\alpha\beta} = \left\langle \cos \left( \phi_\alpha + \phi_\beta - 2\Psi_{EP} \right) \right\rangle$$

# Balance Function


$$B(\phi, \Delta\phi) = \frac{1}{2} \left\{ \frac{N_{+-}(\phi, \Delta\phi) - N_{++}(\phi, \Delta\phi)}{N_+(\phi)} + \frac{N_{-+}(\phi, \Delta\phi) - N_{--}(\phi, \Delta\phi)}{N_-(\phi)} \right\}$$



# Data Set




## STAR Detector






- 200 GeV Au+Au events from STAR Run 4
- 14 million events analyzed
- All charged particles
  - $|\eta| < 1.0$
  - Full azimuthal acceptance
  - $0.2 < p_t < 2.0$  GeV/c
  - Electrons are suppressed
  - 2<sup>nd</sup> order event plane from TPC

# Balance Function



- 40-50% centrality
- $45^\circ$  to event plane balance function is biased toward negative  $\Delta\phi$  region
- The out-of-plane balance function is wider than the in-plane balance function

Compare to blast wave model calculations<sup>1</sup>



<sup>1</sup> S. Schlichting and S. Pratt  
Phys. Rev. C 83, 014913 (2011)

# Blast Wave Model




∞ STAR

parameterization



(STAR, PRC, 72, 14904 (2005))

∞ Local charge  
conservation

S. Schlichting and S. Pratt  
Phys. Rev. C 83, 014913 (2011)



# Balance Function v.s. Centrality




Balance function narrows  
in central collision due  
to collective flow

# Three Point Correlator



Phys. Rev. Lett. 103, 251601 (2009)



- $\gamma_P$  is the difference between unlike- and like-sign correlations
- Blast wave model reproduces observed difference between unlike- and like-sign azimuthal correlations

$$\gamma_{\alpha\beta} = \langle \cos(\phi_\alpha + \phi_\beta - 2\Psi_{RP}) \rangle$$


$$\gamma_p = \frac{1}{2} (2\gamma_{+-} - \gamma_{++} - \gamma_{--}) = \frac{2}{M^2} \int d\phi d\Delta\phi \frac{dM}{d\phi} B(\phi, \Delta\phi) [\cos 2\phi \cos \Delta\phi - \sin 2\phi \sin \Delta\phi]$$



# Three Point Correlator



Phys. Rev. Lett. 103, 251601 (2009)



- $\gamma_P$  is the difference between unlike- and like-sign correlations
- Blast wave model reproduces observed difference between unlike- and like-sign azimuthal correlations

$$\gamma_{\alpha\beta} = \langle \cos(\phi_\alpha + \phi_\beta - 2\Psi_{RP}) \rangle$$

$$\gamma_p = \frac{1}{2} (2\gamma_{+-} - \gamma_{++} - \gamma_{--}) = \frac{2}{M^2} \int d\phi d\Delta\phi \frac{dM}{d\phi} B(\phi, \Delta\phi) [\cos 2\phi \cos \Delta\phi - \sin 2\phi \sin \Delta\phi]$$

# Summary




- ∞ The reaction-plane-dependent balance function analysis gives the same difference between the like-sign and unlike-sign charge dependent azimuthal correlations as the three point correlator results published by STAR
- ∞ This thermal blast wave model **reproduces most of the difference** between like- and unlike-sign charge-dependent azimuthal correlation incorporating **local charge conservation** and **flow**




# Back Up

# Beam Energy Dependence



# Weighted Average





- Compare data (points) with blast wave model calculations (solid lines)
- Data are not corrected for event plane resolution (differences between data and model)
- $c_b$  is related to the balance function width, while  $s_b$  quantifies the asymmetry of balance function
- Data show a stronger collective behavior in plane, while the asymmetry is most significant  $45^\circ$  to the reaction plane

$$c_b(\phi) \equiv \frac{1}{z_b(\phi)} \int d\Delta\phi B(\phi, \Delta\phi) \cos(\Delta\phi)$$

$$s_b(\phi) \equiv \frac{1}{z_b(\phi)} \int d\Delta\phi B(\phi, \Delta\phi) \sin(\Delta\phi)$$

$$z_b(\phi) \equiv \int d\Delta\phi B(\phi, \Delta\phi)$$

# Calculate $v_{2c}$ and $v_{2s}$



- Red arrows are the first particle in balance function calculation, while green ones are its opposite sign charge pairs
- $v_{2c} < c_b >$  is positive if more charged pairs are found in plane
- $v_{2c}$  is positive if charges are more correlated in plane
- $v_{2s}$  is negative if charges are more correlated on the in plane side

$$v_{2c} \equiv \langle c_b(\phi) \cos(2\phi) \rangle - v_2 \langle c_b(\phi) \rangle$$

$$v_{2s} \equiv \langle s_b(\phi) \sin(2\phi) \rangle$$

$$\langle f(\phi) \rangle \equiv \frac{1}{M} \int d\phi \frac{dM}{d\phi} z_b(\phi) f(\phi)$$

S. Schlichting and S. Pratt  
 Phys. Rev. C 83, 014913 (2011)