



Probing the Quark-Gluon Phase Transition with Long-Range Multiplicity Correlations in Heavy Ion Collisions from the STAR Experiment

Terence J Tarnowsky
for the STAR Collaboration
July 28, 2009



Outline


- Motivation
- Experimental Method
- Results
- Future Plans
- Summary



Study of Correlations and Fluctuations



- Expectation that correlations reflect features of multi-particle production.
 - e.g. Bose-Einstein/HBT (source size).
- Forward-backward correlations in (pseudo)rapidity characterize formation and decay of possible clusters.
 - Number of sources.
 - Size of sources.
- Changes in particle number fluctuations (π , K , p) expected near a phase transition.



F-B Multiplicity Correlations

- Predicted in context of **Dual Parton Model** [DPM] (and **Color Glass Condensate** [CGC]).
- Test of multiple elementary [partonic] scattering.
- Linear expression relating N_b and N_f (forward and backward multiplicity), found in hadron-hadron experiments (ex. UA5),

$$\langle N_b \rangle (N_f) = a + bN_f \quad \boxed{N = \# \text{ of hadrons}}$$

- “b” is correlation strength.


- Function of \sqrt{s} and A.

- Coefficient can be expressed as,

$$b = \frac{\langle N_f N_b \rangle - \langle N_f \rangle \langle N_b \rangle}{\langle N_f^2 \rangle - \langle N_f \rangle^2} = \frac{D_{bf}^2}{D_{ff}^2}$$

Short- and Long-Range Correlations

$$b = \frac{\langle n_f n_b \rangle - \langle n_f \rangle \langle n_b \rangle}{\langle n_f \rangle^2 - \langle n_f \rangle} = \frac{D_{bf}^2}{D_{ff}^2}$$



- $(D_{ff})^2$ characterizes short range correlations, related to the number of emitted particles per cluster.
- $(D_{bf})^2$ is related to the number of sources.

• Long-range is taken as a separation in η of forward-backward windows by > 1.0 .



Strings

Low Energy

$$\eta = -\ln \left[\tan \left(\frac{1}{2} \theta \right) \right]$$

Long Range
Backward n_b

Long Range
Forward n_f

Short + Long Range


High Energy

$-\eta_2$ $-\eta_1$ 0 η_1 η_2 η

Pseudorapidity interval

Pseudorapidity Gap

Probes longitudinal characteristics of the system.






Short and Long-Range FB Multiplicity Correlations

- Working definitions:
 - Short-range correlations (SRC):
 - SRC is defined as correlation for $|\Delta\eta| < 1.0$.
 - Long-range correlations (LRC):
 - LRC is defined as correlations for $|\Delta\eta| > 1.0$.


Energy Dependence: pp Min Bias

- Comparing correlation as function of $\sqrt{s_{NN}}$.
- 200 and 400 GeV in close agreement:
 - Larger SRC at 400 GeV
 - Plateau at same value of b at large $\Delta\eta$ as 200 GeV.
- 62.4 GeV goes smoothly to $b = 0$.
 - Not even a small LRC.



Centrality Dependence:

Au+Au at 200 GeV




- Central collisions demonstrate a **strong, LRC**.

- For mid-peripheral collisions:

- **LRC is absent.**


Centrality Dependence:

200 GeV Au+Au




Comparison to Parton String Model (PSM) and HIJING

- 2 Monte Carlo models: PSM (DPM) and HIJING.
- HIJING does not have long strings in η .
- PSM has these included.
- PSM shows qualitative agreement w/ data.
- HIJING agrees well w/ the short-range component.



Centrality Dependence: 40-50% Cu+Cu at 200 GeV

- As in mid-peripheral Au+Au:
- LRC absent.
- Approx same evolution w/ $\Delta\eta$.







Search for the QCD Critical Point

- Proposal for future running at RHIC to consist of an “energy scan” to search for predicted QCD critical point.
- Fluctuations and correlations (particle ratios, multiplicity, p_T , etc.) and behavior of flow (directed and elliptic) in vicinity of the critical point are expected to be primary signatures.
- F-B correlations and K/π , p/π fluctuations can be measured at **all energies**.

Energy Dependence: Central 0-10% Cu+Cu



Centrality Evolution: 22 GeV Cu+Cu




Energy Dependence: 40-50% Cu+Cu

• For mid-peripheral collisions:

• **LRC is absent at all energies.**

• All energies show approx. same evolution w/ η .





Particle Ratio Fluctuations

K/π and p/π

- NA49 uses the variable σ_{dyn}

$$\sigma_{\text{dyn}} = \text{sign}(\sigma_{\text{data}}^2 - \sigma_{\text{mixed}}^2) \sqrt{|\sigma_{\text{data}}^2 - \sigma_{\text{mixed}}^2|}$$

σ is relative width of K / π distribution

- Measure deviation from Poisson behavior using V_{dyn}


$$V_{\text{dyn},K\pi} = \frac{\langle N_K (N_K - 1) \rangle}{\langle N_K \rangle^2} + \frac{\langle N_\pi (N_\pi - 1) \rangle}{\langle N_\pi \rangle^2} - 2 \frac{\langle N_K N_\pi \rangle}{\langle N_K \rangle \langle N_\pi \rangle}$$

- It has been demonstrated that,


$$\sigma_{\text{dyn}}^2 = V_{\text{dyn}}$$

Excitation Function for $\sigma_{\text{dyn},K\pi}$

Compare STAR results for central Au+Au collisions with SPS results for central Pb+Pb collisions




Excitation Function for $\sigma_{\text{dyn},p\pi}$




Cu+Cu 22.4 GeV


$V_{dyn, K/\pi}$



Cu+Cu 22.4 GeV

$(dN/d\eta)v_{dyn,K/\pi}$







Summary I

- **Centrality Dependence:**
 - At 200 GeV, central A+A collisions exhibit a **strong, long-range correlation**.
 - For mid-peripheral collisions, correlation coefficient shows similar behavior at all energies and systems.
 - Dominated by SRC.
 - LRC is consistently absent.
- **System Size Dependence:**
 - Similar LRC at 200 GeV in Au+Au and Cu+Cu.
 - Correlation in pp at 200 GeV resembles that of mid-peripheral A+A.
- **Energy Dependence:**
 - For central collisions, correlation coefficient seems to depend more on energy than system size.
 - Differences in correlation in pp 200/400 vs. 62 GeV could indicate small LRC.
 - Jet contribution?

Summary II

- Preliminary look at $v_{\text{dyn},K/\pi}$ for Cu+Cu at 22.4 GeV.
 - More track quality cuts to be investigated.
- For $v_{\text{dyn},p/\pi}$ models show a transition effect.
 - Acceptance in HSD...?
 - Not acceptance in UrQMD. What is it?
- Study of fluctuations and correlations well established, will be an important part of QCD critical point search.