

Azimuthal anisotropic flow of identified hadrons in Au+Au collisions in BES-II energies

Prabhupada Dixit (for the STAR collaboration)

September 2022

1 Abstract

Anisotropic flow in heavy-ion collisions reveals crucial information about the created matter's properties. Elliptic (v_2) and triangular (v_3) flow coefficients, representing second and third order azimuthal anisotropic particle distributions in momentum space, offer sensitivity to the equation of state and transport properties like shear viscosity to entropy density ratio (η/s) of the medium.

Recently, STAR has completed the data taking of Beam Energy Scan phase-II (BES-II) with improved detector conditions and wider rapidity coverage. In this talk, using BES-II data, we will present the high precision measurements of v_2 and v_3 of identified hadrons, $\pi^+(\pi^-)$, $K^+(K^-)$, $p(\bar{p})$, K_S^0 , ϕ , $\Lambda(\bar{\Lambda})$, $\Xi^-(\bar{\Xi}^+)$, and $\Omega^-(\bar{\Omega}^+)$ in Au+Au collisions at $\sqrt{s_{NN}} = 14.6$ and 19.6 GeV. We will show the centrality, transverse momentum, rapidity dependence, and the number of constituent quark (NCQ) scaling of v_n . Finally, the physics implications of our measurements in the context of partonic collectivity will be discussed.