




Measurement of Total and Elastic Cross Sections in p+p collisions at $\sqrt{s} = 200$ GeV with the STAR Detector at RHIC



Włodek Guryn and Bogdan Pawlik

For the STAR Collaboration




1. Experimental setup at STAR
2. Data set
3. Analysis
4. Distributions of physics variables ($-t$, ϕ)
5. Simulations and efficiency, acceptance corrections
6. Results: $d\sigma/dt$, B-slope, σ_{tot} , σ_{el}

Experimental Setup




Only a dipole magnet DX between the Roman Pots (RPs) and the collision point. Bending only in (x,z) plane.



In this configuration, RP program at STAR was able to acquire large data samples without special running conditions – mostly for CEP, SDD and CP analyses (see talks by L. Fulek and R. Sikora at this conference).

Data Analysis



- Trigger was very inclusive: it required only a signal in at least one RP on each side.

$$\mathbf{RP_ET} = (\mathbf{E1U} \vee \mathbf{E2U} \vee \mathbf{E1D} \vee \mathbf{E2D}) \wedge (\mathbf{W1U} \vee \mathbf{W2U} \vee \mathbf{W1D} \vee \mathbf{W2D})$$

- Need to minimize background and maximize efficiency.
- To reduce background need angle reconstruction => two RPs on each side in up – down combination.

$$\begin{aligned} \mathbf{EU} &= (\mathbf{E1U} \wedge \mathbf{E2U}) ; \mathbf{ED} = (\mathbf{E1D} \wedge \mathbf{E2D}) \\ \mathbf{WU} &= (\mathbf{W1U} \wedge \mathbf{W2U}) ; \mathbf{WD} = (\mathbf{W1D} \wedge \mathbf{W2D}) \\ \mathbf{ET1} &= (\mathbf{EU} \wedge \mathbf{WD}) \\ \mathbf{ET2} &= (\mathbf{ED} \wedge \mathbf{WU}) \end{aligned}$$

- Use events with four track points – one track point per Roman Pot.
- Finally, choose fiducial region away from the apertures of DX magnet and beam pipe in front of the RPs.


Collinearity

$$\vec{p}_1 = -\vec{p}_2 \Rightarrow (\Theta_{x1}, \Theta_{y1}) = (-\Theta_{x2}, -\Theta_{y2}) \Rightarrow \Delta\Theta_x = \Delta\Theta_y = 0$$


Since the elastic events must satisfy collinearity condition collinearity within $2\sigma_\theta$.

Namely $|\theta_{\text{West}} - \theta_{\text{East}}| < 2\sigma_\theta$, where $\sigma_\theta = 255 \mu\text{rad}$, is required.

Events are well centered within 2σ and 3σ contours.



Geometrical Acceptance GEANT4 MC: I




Choice of geometrical acceptance (t, ϕ) plane

$$0.04 \leq |t| \leq 0.16 [(\text{GeV}/c)^2]$$

$$79.5 \leq |\phi| \leq 101.5 [\text{deg}]$$

$$2.00 \leq \theta \leq 4.00 [\text{mrad}]$$

Geometrical Acceptance and Event Yields




Choose region away from steep variation and edges of acceptance

Condition	# events
ET triggered	6.607M
ET accepted	3.974M
Collinear	2.696M
4 PT Collinear	1.100M
4 PT Collinear Geom.	0.667M

667K events used for the final analysis

GEANT4 MC: Background Study

1. Each distribution is normalized to 1, independently
2. Normalization MC to Data done by normalizing peaks
3. Background mostly due to the re-scattered protons in the the beam pipe and the DX magnet
4. Background is small – 0.3%, after $2\Delta\Theta$ cut and after geometrical acceptance cut



Results: Corrected $d\sigma/dt$ and Fits


$$\frac{d\sigma_{el}}{dt} = \frac{1 + \rho^2}{16\pi(\hbar c)^2} \cdot \sigma_{tot}^2 \cdot e^{-B|t|}$$

$$\sigma_{tot}^2 = \left(\frac{16\pi(\hbar c)^2}{1 + \rho^2} \right) \frac{d\sigma_{el}}{dt} \Big|_{t=0}$$

$$\sigma_{el} = \int \frac{d\sigma_{el}}{dt} dt$$

The value of $\rho = 0.128$ from COMPETE model was used*.

* Phys. Rev. Lett. 89 (2002) 201801



Results



Fit results

FILTER	$d\sigma_{el}/dt _{t=0}$ [mb/ GeV ²]	B [GeV ⁻²]	σ_{tot} [mb]	σ_{el} [mb]
4PT-COL	134.3 ±1.6	14.0 ±0.2	50.7 ±0.6	9.6 ±0.1
4PT-GEO	136.7 ±0.8	14.2 ±0.2	51.3 ±0.4	9.6 ±0.1

name	Quantity		Statistical uncertainty	Systematic uncertainty
	units	Value		
B	[(GeV/c) ⁻²]	14.2	±0.1	±0.3
σ_{el}	[mb]	9.6	±0.1	±0.7
σ_{tot}	[mb]	51.3	±0.4	+2.1 -1.9

The main source of systematic uncertainty are: luminosity measurement and beam tilt angle.

Comparison with the World Data



STAR point size reflects the total uncertainty

STAR results compare well with the world data and the COMPETE predictions: Phys. Rev. Lett. 89 (2002) 201801

Plots from the TOTEM Collaboration <https://arxiv.org/pdf/1712.06153v2.pdf> with STAR preliminary results added

Summary

1. The STAR experiment at RHIC measured elastic differential cross sections in the $|t|$ -range $[0.045, 0.125] \text{ (GeV/c)}^2$ in p+p collisions at $\sqrt{s} = 200 \text{ GeV}$.
2. The resulting values of B-slope, σ_{tot} , σ_{el} are:
 - Slope parameter $B = 14.2 \pm 0.1 \text{ (stat)} \pm 0.3 \text{ (syst)} \text{ (GeV/c)}^{-2}$
 - The total cross section $\sigma_{\text{tot}} = 51.3 \pm 0.4 \text{ (stat)} + 2.1 - 1.9 \text{ (syst)} \text{ (mb)}$
COMPETE Predictor, Phys. Rev. Lett. 89 (2002) 201801 $\sigma_{\text{tot}} = 51.76 \pm 0.12 \text{ (stat)} + 0.4 - 0.2 \text{ (syst)} \text{ mb}$
 - The elastic cross section $\sigma_{\text{el}} = 9.6 \pm 0.1 \text{ (stat)} \pm 0.7 \text{ (syst)} \text{ mb}$

At this point, the largest syst. uncertainties are: 1% due to the beam tilt angle and 7% due to the luminosity.

We expect the luminosity uncertainty to be about 3% after the careful calibration.