

STAR Heavy Flavor Overview

Dandan Shen (沈丹丹) for the STAR Collaboration

Shandong University (山东大学)


The 9th International Symposium on Heavy Flavor
Production in Hadron and Nuclear Collisions

Why heavy flavor?




- **Heavy quarks (c and b) are produced in early hard scatterings**
- **Test QCD Predictions**
 - Explore perturbative and non-perturbative QCD
- **Probe Quark-gluon plasma (QGP) Properties**
 - QGP dynamics and spin related effect

Open heavy flavor




D mesons, etc

Quarkonium



J/ψ , $\psi(2s)$, etc

The STAR detector




Presented here collision species:


- p+p: $\sqrt{s} = 500$ GeV
- Heavy-ion collisions:
 - Isobar (Ru+Ru, Zr+Zr): $\sqrt{s_{NN}} = 200$ GeV
 - Au+Au: BES-II($\sqrt{s_{NN}} = 14.6 - 27$ GeV), $\sqrt{s_{NN}} = 200$ GeV

Open Heavy Flavor

D⁰ suppression



✓ Energy loss in medium




Model: G. Qin, private communication
 Phys. Rev. C 99 (2019) 034908


$$R_{AA} = \frac{\sigma_{inel}^{NN} d^2 N_{AA}^{D^0} / dp_T dy}{\langle N_{coll} \rangle d^2 \sigma_{pp}^{D^0} / dp_T dy}$$

- Similar level of suppression in Isobar and Au+Au at same centrality
- Qualitatively reproduced by energy loss model calculations


D⁰-tagged jet yield and fragmentation



- ✓ Charm quark fragmentation function in medium




$$z_{\text{Jet}} = \frac{\vec{p}_{T,\text{Jet}} \cdot \vec{p}_{T,D^0}}{p_{T,\text{Jet}}^2}$$




- Hard-fragmented jets (large z) are more suppressed than soft-fragmented ones (small z)

$D^0-\bar{D}^0$ azimuthal correlations





✓ Energy loss and thermalization in medium \rightarrow weaker correlation at $\Delta\phi \approx \pi$ in HIC

Phys. Lett. B 647 (2007) 366–370



- No azimuthal correlation is seen for $D^0-\bar{D}^0$ pairs within large uncertainties
- Need better precision


D⁰ freeze-out property





STAR Phys. Rev. C 99 (2019) 034908

- D⁰ freezes out earlier than light hadrons
- No significant collision system dependence in isobar and Au+Au collisions

$D^0-\pi^\pm$ femtoscopic correlations






- ✓ Freeze-out dynamics and final state interaction



- No significant correlations, but also consistent with theoretical predictions with emission source size of 5 fm or larger within uncertainty

Quarkonium

J/ψ production mechanism



Model with different LDMEs describe the J/ψ p_T spectrum well, but differ in J/ψ fragmentation predictions.


Phys.Rev. Lett. 119 (2017) 032001

- No significant z dependence observed in data within uncertainties for $z < 1$
- Data show less isolated production than PYTHIA8


J/ψ + many charged jet activity

J/ψ + little charged jet activity

J/ψ R_{AA} vs. collision energy



✓ Hot and cold nuclear matter effect




- No significant energy dependence of J/ψ R_{AA} below 200 GeV
- The observed energy dependence is qualitatively described by the transport model



X. Zhao, R. Rapp, *Phys. Rev. C* 82 (2010) 064905 (private communication).

L. Kluberg, *Eur. Phys. J. C* 43 (2005) 145.

J/ψ R_{AA} and v_2 in isobar collisions




✓ J/ψ suppression and regeneration in HIC




- Indication of small regeneration effect in isobar collisions?
- Clear J/ψ suppression with little v_2

Charmonium sequential suppression



✓ QGP thermal properties at RHIC



➤ $\psi(2S)$ over J/ψ double ratio is smaller than that in p+A collisions

➤ First observation of charmonium sequential suppression in HIC at RHIC (3.5σ , 0-80%)

PHENIX, *Phys.Rev.D*, 85,092004 (2012)

NA51, *Phys.Lett.B* 438 (1998) 35-40


ISR, *Nucl.Phys.B* 142 (1978) 29

J/ψ polarization





✓ Angular distribution of the decayed leptons:

$$W(\cos\theta, \phi) \propto 1 + \lambda_\theta \cos^2\theta + \lambda_\phi \sin^2\theta \cos 2\phi + \lambda_{\theta\phi} \sin 2\theta \cos\phi$$



Faccioli et al, EPJC 69: 657-673 (2010)




STAR, Nature 614, 244-248 (2023)


✓ Definition of the axes

- **Helicity frame (HX):** J/ψ momentum direction
- **Collins-Soper frame (CS):** bisector of angle between beams
- **“Event plane”:** axis orthogonal to reaction plane $\lambda_\theta = \frac{1-3\rho_{00}}{1+\rho_{00}}$

J/ψ polarization vs. centrality




✓ Sequential melting of charmonium suppresses feed-down contribution ($\psi(2s), \chi_{cJ}$)




- No significant centrality dependence
- $\lambda_\theta, \lambda_\phi$ are consistent with zero
- Consistent λ_{inv} for the two frames, as expected

$$\lambda_{inv} = \frac{\lambda_\theta + 3\lambda_\phi}{1 - \lambda_\phi}$$

J/ψ global spin alignment at RHIC




- ✓ ϕ meson $\rho_{00} > 1/3$ at RHIC
 - ✓ J/ψ $\rho_{00} < 1/3$ at LHC forward rapidity
- How about J/ψ ρ_{00} at RHIC energy?




➤ ρ_{00} lower than $1/3$ with a significance of 3.5σ using 2nd order event plane

STAR, Nature 614, 244-248 (2023)
 Alice, Phys. Rev. L 131 4, 042303 (2023)

J/ψ global spin alignment: RHIC vs. LHC




Zr(Ru)+Zr(Ru) 200 GeV





➤ The ρ_{00} at RHIC energy is comparable to LHC results, with different collision energies, systems and rapidities

Photon-induced J/ψ decay anisotropy



✓ Photon-induced J/ψ decay anisotropy expected due to linearly polarized photon



$\Delta\phi[e^+ - e^-, \Psi_{EP}^{2nd}]$

Ψ_{EP}^{2nd} : second order TPC event plane

- Photon-induced $\cos(2\Delta\phi)$ modulation with a significance of $\sim 3.3 \sigma$
- Measured modulation consistent with EPA-VMD model prediction

X. Wu et al. Phys. Rev. Res. 4, L042048 (2022)

Summary and outlook



✓ Open heavy flavor

- **Energy loss:** similar level of D^0 suppression in Isobar and Au+Au 200 GeV; larger suppression of hard-fragmented D^0 jets in Au+Au
- **Freeze-out and final state interaction:** D^0 freezes out earlier than light hadrons; D^0 -hadron femtoscopy correlations suggest source size of 5 fm or larger



✓ Quarkonia

- **Production mechanism:** no significant z dependence of J/ψ in charged jet for $z < 1$ in p+p 500 GeV
- **Dissociation and regeneration:** charmonium sequential suppression and limited J/ψ v_2 in isobar collisions; no significant energy dependence of J/ψ R_{AA} in Au+Au collisions @14.6-200 GeV
- **Polarization and spin alignment:** J/ψ polarization is consistent with zero in HX and CS frames; J/ψ spin alignment and photon-induced J/ψ decay anisotropy are observed in isobar

Summary and outlook






- Run23-25: high statistics p+p(Au), Au+Au collisions
- STAR detector with enhanced capabilities
 - Higher DAQ rate; extended coverage (iTPC+ forward upgrade)



STAR BUR Run25 2024

Back up

J/ψ global spin alignment at RHIC



- ρ_{00} consistent with 1/3 with large uncertainty using 1st order event plane
- ρ_{00} lower than 1/3 with a significance of 3.5 σ using 2nd order event plane