

Measurements of non-photonic electrons in STAR

Olga Rusňáková for the STAR Collaboration

Faculty of Nuclear Sciences and Physical Engineering

Czech Technical University in Prague

Hot Quarks, Las Negras, Spain, 21-27 September, 2014




Outline

- Motivation.
- Non-photonic electrons (NPE) analysis method.
- NPE results:
 - p+p at 200 GeV.
 - Au+Au collisions, nuclear modification factor R_{AA} at 200 GeV.
 - Azimuthal anisotropy v_2 in Au+Au at 200 GeV.



Heavy ion collisions

- Heavy ion collisions:
 - hot and dense nuclear matter formation - **Quark-Gluon Plasma**
 - cold and hot nuclear matter effects
- p+p collisions:
 - baseline
- Medium effects quantified by **nuclear modification factor**:
 - R_{dA} – cold nuclear matter effects.
 - R_{AA} – hot nuclear matter effects.




$$R_{AA} = \frac{1}{\langle N_{coll} \rangle} * \frac{dN_{AA}/dy}{dN_{pp}/dy}$$



Heavy quarks

- Heavy quarks:
 - large masses
 - early production
 - p+p collisions - test of the validity of the pQCD



Non-photonic electron

- Heavy ion collisions:
 - energy loss
(nuclear modification factor R_{AA})
 - thermalization (elliptic flow v_2)

- Study of non-photonic electrons is a good way to measure production of bottom and charm hadrons via semi-leptonic decays.


$$b \rightarrow e^{\pm} + \text{anything} (10.86\%)$$

$$c \rightarrow e^{\pm} + \text{anything} (9.6\%)$$



STAR detector at RHIC

Solenoidal Tracker At RHIC : $-1 < \eta < 1, 0 < \phi < 2\pi$



Time Projection Chamber (TPC) – tracking, particle identification, momentum

Time of Flight detector (ToF) – particle identification at low p_T region.

BEMC – electron identification at high p_T region, triggering (High Tower triggers)



BSMD – electron identification at high p_T



STAR detector at RHIC

Solenoidal Tracker At RHIC : $-1 < \eta < 1, 0 < \phi < 2\pi$

Electron identification with TPC in low p region.








STAR detector at RHIC

Solenoidal Tracker At RHIC : $-1 < \eta < 1, 0 < \phi < 2\pi$

TPC and TOF together are great tool for distinguish electrons and hadrons in low p







STAR detector at RHIC

Solenoidal Tracker At RHIC : $-1 < \eta < 1, 0 < \phi < 2\pi$

TPC and BEMC together are used for electron identification at high p_T region.






NPE analysis method

$$NPE = N_{Inclusive} * purity_{Inclusive} - \frac{N_{Photonic}}{\epsilon_{Photonic}}$$




- **Inclusive electrons** – identification with TPC, TOF, BEMC.
- **Photonic electrons** – background
 - identified via small e^+e^- invariant mass
 - statistically reconstructed
 - main **background** comes from:
 - Dalitz decay: $\pi^0 \rightarrow \gamma + e^+ + e^-$ (BR: $\sim 1.2\%$)
 - Gamma conversions: $\gamma \rightarrow e^+ + e^-$
 - corrected for **reconstruction efficiency** via simulation.





Inclusive electron purity

$$NPE = N_{Inclusive} * \text{purity}_{Inclusive} \frac{N_{Photonic}}{\epsilon_{Photonic}}$$





- Multi-gaussian fit to the nsigele distribution.

$$nsigele = \ln \left(\frac{dE/dx_{exp}}{dE/dx_{Bethe/Bloch_e}} \right) / \sigma$$



Photonic background reconstruction efficiency





- Photonic electron reconstruction efficiency from simulation.

$$NPE = N_{Inclusive} * purity_{Inclusive} = \frac{N_{Photonic}}{\epsilon_{Photonic}}$$



NPE in p+p collisions at $\sqrt{s}=200\text{GeV}$

- p+p at 200GeV data (year 2009):
- Spectrum was extended to the low p_T region.




p+p 2008 - H. Agakishiev et al. [STAR Collaboration], Phys. Rev. D83, 052006,(2011)
 p+p PHENIX - A.Adare et al. [PHENIX Collaboration], , Phys. Rev. C84, 044905 (2011).
 FONLL - M. Cacciari, P. Nason and R. Vogt, Phys. Rev. Lett. 95, 122001 (2005)



Data to FONLL ratio

- Spectrum was extended to the low p_T region.
- Consistent with pQCD FONLL calculation (Fixed Order plus Next-to-Leading Logarithms).





p+p 2008 - H. Agakishiev et al. [STAR Collaboration], Phys. Rev. D83, 052006,(2011)
p+p PHENIX - A.Adare et al. [PHENIX Collaboration], , Phys. Rev. C84, 044905 (2011).
FONLL - M. Cacciari, P. Nason and R. Vogt, Phys. Rev. Lett. 95, 122001 (2005)



NPE in Au+Au collisions at $\sqrt{s}_{NN} = 200\text{GeV}$

- Au+Au at 200GeV (year 2010 data):
 - suppression at high p_T compared with FONLL calculations






Nuclear Modification Factor in Au+Au collisions at $\sqrt{s}_{NN} = 200\text{GeV}$

→ Strong suppression is observed at high p_T .

→ Strong suppression is similar as for D^0 mesons and light hadrons.




$$R_{AA} = \frac{1}{\langle N_{coll} \rangle} * \frac{dN_{AA}/dy}{dN_{pp}/dy}$$

D^0 - arXiv:1404.6185 [nucl-ex] – accepted by PRL





NPE elliptic flow in Au+Au collisions at $\sqrt{s}_{NN} = 200$ GeV

- Initial geometry asymmetry \rightarrow final momentum anisotropy.
- Results obtain using 2-particle and 4-particle correlations.
- Finite v_2 at low p_T together with strong suppression in high p_T indicates strong charm-medium interaction.
- Increase of v_2 at high p_T can arise from jet-like correlations or from path length dependence of heavy quark energy loss.





NPE R_{AA} and elliptic flow in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV



Glueon radiation scenario alone fails to explain large NPE suppression at high p_T .



Finite elliptic flow together with large suppression at high p_T at $\sqrt{s_{NN}} = 200$ GeV indicate that heavy quarks interact strongly with the surrounding partonic medium.

It's challenging for model calculations to describe the suppression and v_2 simultaneously.



Heavy flavor tracker (HFT)

Projected NPE results with HFT



Curves: H. van Hees *et al.* Eur. Phys. J. **C61**, 799(2009).

- Heavy Flavor Tracker (HFT) - Year 2014.
- HFT will allow measurement of $B \rightarrow e$ and $D \rightarrow e$ spectrum separately in Au+Au.



Conclusions

- Measurement of the NPE spectrum in p+p collisions was extended to the low p_T region.
- We observed **strong suppression** of NPE in **Au+Au collisions**. Similar suppression as for light hadrons.
- NPE v_2 at 200 GeV: finite v_2 at low p_T together with strong suppression at high p_T indicates a strong charm-medium interaction.
- The new HFT detector will allow measurement of $B \rightarrow e$ and $D \rightarrow e$ spectra separately.



Thank you!