

Kaon femtoscopy at the STAR experiment


Jindřich Lidrych for the STAR Collaboration

Faculty of Nuclear Sciences and Physical Engineering
Czech Technical University in Prague

Hot Quarks 2016

South Padre Island, Texas, USA

12th – 17th September 2016



Femtoscopy

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES



Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy



Model comparison

Conclusions



- Study space-time extents of the source at the thermal freeze-out
- Correlation function: $CF(p_1, p_2) = \int d^3 r S(r, k) |\psi_{1,2}(r, k)|^2$

$$r = x_1 - x_2 \quad q_{\text{inv}} = p_1 - p_2 = 2k^*$$



Femtoscopy with kaons – a cleaner probe

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions

In comparison with the most abundant pions, there are following advantages

- Less feed-down – smaller contamination with non-primary kaons from resonance decays
- Smaller cross section – information about a different stage of the collisions evolution
- Kaons contain strange quark

However, more difficult due to ~10 smaller statistics

STAR Experiment at RHIC

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES


Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions



Time Projection Chamber

- dE/dx PID
- Tracking
- $0 < \varphi < 2\pi, |\eta| < 1$

Time of Flight

- Time resolution $< 80\text{ps}$
- $1/\beta$ PID

Charged kaon femtoscopy for BES

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions

RHIC Beam Energy Scan

- One of the main physics program at the RHIC
- The goal of Beam Energy Scan:
 - Find the QCD critical point
 - 1st order phase transition signs
 - Turn-off sQGP signatures

\sqrt{s}_{NN} (GeV)	μ_B (MeV)	#Events	#Weeks	Year
200	20	350 M	11	2010
62.4	70	67 M	1.5	2010
39.0	115	130 M	2	2010
27.0	155	70 M	1	2011
19.6	205	36 M	1.5	2011
14.5	260	20 M	3	2014
11.5	315	12 M	2	2010
7.7	420	4 M	4	2010


1D femtoscopic analysis of charged kaons

- Motivation: Is there a difference between K^+ and K^- source?
- 6 energies: 7.7, 11.5, 14.5, 19.6, 27 and 39 GeV
- 2 centrality bins (0-30% and 30-80%)
- 2 k_T bins (0.2-0.4 GeV/c and 0.4-0.6 GeV/c)
- Fitting function: *Phys. Lett.*, B270:69–74, 1991

$$CF(q_{inv}) = \left[(1 - \lambda) + \lambda K(q_{inv}, R_{inv})(1 + e^{-R_{inv}^2 q_{inv}^2}) \right] \mathcal{N},$$

where R_{inv} – source radii, λ – correlation strength,

$K(q_{inv}, R_{inv})$ – Coulomb function and \mathcal{N} – normalization



Results from kaon femtoscopy for BES

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy


Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions

- No clear beam energy dependence visible
- Possible different behavior for K^+ and K^- for energy 10-20 GeV ?



Kaon femtoscopy at top RHIC energy

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison


Conclusions

Au+Au collisions at $\sqrt{s_{NN}} = 200 \text{ GeV}$

In the past, STAR has already performed the first measurements with kaons

Phys. Rev. C88 (2013) 34906

- Used data were recorded in 2004 and 2007
- Only TPC for PID
- Data favor models that break the m_T – scaling



Kaon femtoscopy at top RHIC energy

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison


Conclusions

Au+Au collisions at $\sqrt{s_{NN}} = 200 \text{ GeV}$

In the past, STAR has already performed the first measurements with kaons

Phys. Rev. C88 (2013) 34906

- Used data were recorded in 2004 and 2007
 - Only TPC for PID
 - Data favor models that break the m_T – scaling
 - Source imaging** was also performed
 - Technique to obtain kaon source function $S(r, k)$ directly
- > Kaon source can be well described by Gaussian shape**



Kaon femtoscopy at top RHIC energy

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions

Now, higher statistics which allow more precise measurements

- Data were recorded by the STAR in 2011
- One of the largest available statistics
- Time of Flight detector improves PID and extends identification to higher momenta

1D & 3D femtoscopic analysis of charged kaons

- 1D: 5 centrality bins: 0-5%, 5-10%, 10-30%, 30-50% and 50-75%
- 3D: 4 centrality bins: 0-10%, 10-30%, 30-50% and 50-75%
- 4 k_T bins: $(0.05-0.35)\text{GeV}/c$, $(0.35-0.65)\text{GeV}/c$, $(0.65-0.95)\text{GeV}/c$ and $(0.95-1.25)\text{GeV}/c$

Fitting – extraction of source size

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy




Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions

- Applied corrections for detector effects: kaon misidentification and momentum resolution
- **Fit example:** 1D correlation function & projection of 3D correlation function
 - data (points) vs the best fit (lines)
 - good agreement with data



Centrality 0-10%

K^+K^-

$0.35 < k_T < 0.65 \text{ GeV}/c$

STAR preliminary

Results – extracted source size

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES


Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions



$$k_T = \left(\frac{\vec{p}_1 + \vec{p}_2}{2} \right)_T$$

- **k_T and centrality dependence of HBT radii is observed**
 - Source radii increase with the centrality and decrease with pair transverse momentum
 - 1D & 3D: Uncertainty is dominated by systematic error, which is obtained by varying the fit range

Results – KK radii & Blast-wave model

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy


Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions

- Blast-wave parameterization can provide additional insight into the freeze-out configuration
- Simultaneous fit of kaon source radii and particle spectra (*Phys. Rev.*, C69:034909, 2004)



- Only statistical error; systematic errors are under study
- Comparison of PHENIX results with these results – consistent within errors
- Difference between pion and kaon parameters can indicate earlier decoupling of kaons

Femtoscopy with unlike-sign kaons

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions

Higher statistics also allow new possibilities:

Femtoscopy with narrow resonance


- Using strong final-state interaction via the resonance decay
 - Predicted to be sensitive to source spatial extent than measurement at low q_{inv}
 - Statistically advantageous
- Challenge - extension of femtoscopic formalism to higher q_{inv}

Lednický: Phys. Part. Nucl. 40 (2009) 307-352

Pratt et al.: PRC 68 (2003) 054901

K^+K^- correlations:

- Coulomb and strong final state interaction
- $\phi(1020)$ resonance
 - $k^* = 126 \text{ MeV}/c$, $\Gamma = 4.3 \text{ MeV}$
- First systematic study



Raw unlike-sign kaon correlation functions

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy


Results from 200 GeV

K^+K^- femtoscopy


Model comparison

Conclusions

Centrality dependence



k_T dependence



- CFs are sensitive to the source size
- In particular, **unlike-sign kaon CF is sensitive in the region of the resonance**
- In order to **compare experimental** unlike-sign kaon correlation functions to **theoretical predictions**, the purity corrections were done

Comparison of 1D unlike-sign to theoretical model

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV



K^+K^- femtoscopy

Model comparison

Conclusions

- Extracted radii from like-sign kaon femtoscopy are used for theoretical calculation of unlike-sign correlation function
 - Gauss + Lednický model of final-state interaction**
Lednický: Phys.Part.Nucl. 40 (2009) 307-352
 - Includes $\phi(1020)$ resonance due to the FSI
 - $$CF(p_1, p_2) = \int d^3r S(r, k) |\psi_{1,2}(r, k)|^2$$
 - Gaussian parameterization of source size – source size R_{inv} is extracted from the like-sign correlation function fit
 - The theoretical function is transformed to the experimental one via:
$$CF^{exp} = (CF^{theo} - 1)\lambda + 1$$
in order to compare to an experimental correlation function, which is corrected for impurities

Experimental data
for theoretical calculation



Comparison of 1D unlike-sign to Lednický model

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV





K^+K^- femtoscopy

Model comparison

Conclusions

- Lednický model reproduces overall structure of the observed correlation function

Centrality 0-5 %



Comparison of 1D unlike-sign to Lednický model

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV





K^+K^- femtoscopy

Model comparison

Conclusions

- Model under predicts the strength of the correlation functions in the region of resonance for smaller source – it can be interpreted as a breakdown of femtoscopic formalism in region of resonance

Centrality 30-50 %



Conclusions

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions

Charged kaon femtoscopy for RHIC Beam Energy Scan

- Extraction of source radii R_{inv} from 1D correlation function
- Possibly different emitting source radii for K^+ and K^- for energy below 20 GeV

Measurement of K^+K^+ & K^-K^- correlations in Au+Au collisions at 200 GeV

- Extraction of source radii R_{out}, R_{side} and R_{long} from 3D CF
- **Performed kaon source imaging**
 - Study kaon source function in Au+Au collisions at 200 GeV
 - Source can be well described by Gaussian shape
- Source radii used for Blast-wave model to extract freeze-out configuration
 - Results show difference between pion and kaon parameters

Measurement of K^+K^- correlations in Au+Au collisions at 200 GeV

- Strong centrality dependence in $\phi(1020)$ region
- k_T dependence in $\phi(1020)$ region
- Comparison of unlike-sign CF to Lednický model
 - Comparison indicates a breakdown of femtoscopic formalism in region of resonance in peripheral collisions



The End

Femtoscopy

Kaon femtoscopy

STAR Experiment

Kaon femtoscopy
for BES

Results from BES

Kaon femtoscopy
at top RHIC energy

Results from 200 GeV

K^+K^- femtoscopy

Model comparison

Conclusions

Thank you for your attention

