



Higher-Order Cumulants of Net-Proton Multiplicity Distributions in Zr+Zr and Ru+Ru Collisions at $\sqrt{s_{NN}} = 200$ GeV

Lawrence Berkeley National Laboratory

Ho-San Ko

for the STAR Collaboration

Supported by




Outline



1. Introduction & motivation
2. Analysis of $\sqrt{s_{NN}} = 200$ GeV isobaric collisions (Zr+Zr & Ru+Ru)
3. Net-proton cumulants & cumulant ratios
4. Summary & outlook

QCD phase diagram



STAR, Phys. Rev. Lett. 126, 092301 (2021)

1. QCD calculation and model

- Lattice QCD: Cross over at $\mu_B = 0$ [1] and $T = 156.5 \pm 1.5$ MeV [2~5]
- QCD based Model: A critical point followed by first-order phase transition at high μ_B [6]

2. Search for the possible signature of critical point by scanning T vs μ_B :

- by varying collision energy in heavy-ion collisions

[1] Y. Aoki, Nature 443, 675 (2006)

[2] Y. Aoki, JHEP 06, 088 (2009)

[3] HotQCD, Phys. Rev. D 85, 054503 (2012)

[4] S. Gupta, Science 332, 1525 (2011)

[5] HotQCD, Phys. Lett. B 795, 15 (2019)

[6] M. A. Halasz, Phys. Rev. D 58, 096007 (1998)

Fluctuation of conserved quantities

1. Cumulants of conserved quantities (B, Q, S) are related to correlation length of the system

$$\delta N = N - \langle N \rangle \quad C_1 = \langle N \rangle, C_2 = \langle (\delta N)^2 \rangle$$

$$C_3 = \langle (\delta N)^3 \rangle, C_4 = \langle (\delta N)^4 \rangle - 3 \langle (\delta N)^2 \rangle^2$$


$$C_2 = \sigma^2, S = C_3 / (C_2)^{3/2}, \kappa = C_4 / (C_2)^2$$

2. **The higher the cumulant order, the more sensitive to the correlation length**

$$C_2 \sim \xi^2, C_3 \sim \xi^{4.5}, C_4 \sim \xi^7 \quad [1\sim 3]$$

3. The cumulant ratios can be directly compared to theoretical calculations

$$\chi_q^{(n)} = \left(\frac{\partial^n p}{\partial \mu_q^n} \right)_T = \frac{1}{VT^3} \times C_q^n$$



[1] M. A. Stephanov, Phys. Rev. Lett. 102, 032301 (2009)


[2] M. Asakawa, Phys. Rev. Lett. 103, 262301 (2009)

[3] M. A. Stephanov, Phys. Rev. Lett. 107, 052301 (2011)

Net-proton number is used as a proxy to net-B number


Fourth-order cumulant for critical point search

Talk by Yu Zhang (Tue T03-I)



STAR, Phys. Rev. Lett. 126, 092301 (2021)

STAR, Phys.Rev.C 104, 024902 (2021)



M. A. Stephanov, Phys. Rev. Lett. 107, 052301 (2011)


B.-J. Schaefer, Phys. Rev. D 85, 034027 (2012)

J.-W. Chen, Phys. Rev. D 93, 034037 (2016)


J.-W. Chen, Phys. Rev. D 95, 014038 (2017)

- 4th order: predicts a non-monotonic energy dependence due to contribution from QCD critical point

Fifth- and Sixth-order cumulant




STAR, Phys. Rev. Lett. 127, 262301 (2021)



- Transition from QGP to hadronic matter is smooth crossover at $\mu_B \approx 0$.
 6th order: first principle lattice QCD calculation predicts $C_6/C_2 < 0$


Particle multiplicity dependent higher moment analysis



Poster by Ashish Pandav (S1 T07_1)




1. At $\sqrt{s_{NN}} = 200$ GeV: Zr+Zr and Ru+Ru ($A = 96$), and Au+Au ($A=197$) with p+p averaged
2. Large statistics: 2.0B Zr+Zr and 1.9B Ru+Ru events taken at STAR in 2018
3. **Inspect the systematic trend of multiplicity dependence of different collision systems (Zr, Ru, and Au) at the same collision energy**

Solenoid Tracker At RHIC (STAR)



- Time Projection Chamber (TPC):
vertexing & particle identification
- Time Of Flight (TOF) detector:
improves proton purity at $0.8 < p_T < 2.0 \text{ GeV}/c$

Proton identification





(Anti) Proton identification:

- $0.4 < p_T < 0.8$ GeV/c: deviation from the theoretical expectation **red line** $< 2\sigma$
- $0.8 \leq p_T < 2.0$ GeV/c: **red line** dev. $< 2\sigma$ & $0.6 < m^2 < 1.2$ GeV²/c⁴
- Purity: $> 99\%$

Acceptance: $|y| < 0.5$ & $0.4 < p_T < 2.0$ GeV/c

Net-proton distributions & multiplicity distributions



Number of charged particles in $|\eta| < 1$ excluding (anti-)protons

1. Detector efficiency correction [1~3]

- Binomial detector efficiency assumption
- TPC & TOF tracks

2. Centrality bin width correction [4]

- Corrects finite bin width effect

3. Statistical uncertainty calculated based on Delta theorem [5]

Measured distribution

Detector response

Actual distribution

$$p(n) = \sum_N \text{Binomial}(n; N, \epsilon) \times P(N)$$

“N” detector bins with efficiency ϵ

n_i : number of events in multiplicity bin i

$$C_n^{\text{corr}} = \frac{\sum_i n_i C_{n,i}}{\sum_i n_i} = \sum_i \omega_i C_{n,i}$$

[1] X. Luo, Phys. Rev. C 91, 034907 (2015)

[2] T. Nonaka, Phys. Rev. C 95, 064912 (2017)

[3] X. Luo, Phys. Rev. C 99, 044917 (2019)

[4] X. Luo, J. Phys. G: Nucl. Part. Phys. 40, 105104 (2013)




[5] X. Luo, J. Phys. G: Nucl. Part. Phys. 39, 025008 (2012)

Net-proton cumulants

UrQMD centrality determined in a similar way to the data: measure charged-pion & charged-kaon multiplicity



$\langle N_{part} \rangle$: Average number of participating nucleons



UrQMD: hadronic transport model.
 Calculated in the STAR acceptance

1. The cumulants of net-proton in Zr+Zr, Ru+Ru and Au+Au collisions follow the same $\langle N_{part} \rangle$ trend at $\sqrt{s_{NN}} = 200$ GeV
2. UrQMD overpredicts C₁ and C₃ while underpredicts C₂. In general, it shows a similar trend as the data

Au+Au C₁ ~ C₄:
 STAR, Phys. Rev. C 104, 024902 (2021)




Net-proton cumulant ratios

UrQMD centrality determined in a similar way to the data: measure charged-pion & charged-kaon multiplicity




$\langle N_{\text{part}} \rangle$: Average number of participating nucleons

UrQMD: hadronic transport model.
Calculated in the STAR acceptance



1. The cumulant ratios C_4/C_2 , C_5/C_1 , and C_6/C_2 of net-proton in Zr+Zr, Ru+Ru and Au+Au collisions show the same $\langle N_{\text{part}} \rangle$ trend at $\sqrt{s_{\text{NN}}} = 200 \text{ GeV}$
2. UrQMD overpredicts C_4/C_2 . In general, it shows a similar trend as the data

High-order net-proton cumulant ratio comparison



1. Zr+Zr and Ru+Ru collision results fit into the p+p \oplus Au+Au results at $\sqrt{s_{NN}} = 200 \text{ GeV}$
2. All cumulant ratios C_4/C_2 , C_5/C_1 , and C_6/C_2 decrease as the multiplicity increases
 \rightarrow Most central Au+Au collision results become consistent with Lattice QCD prediction for the formation of thermalized QCD matter and smooth crossover transition

Dziękujemy za uwagę!!

1. At $\sqrt{s_{NN}} = 200$ GeV: the highest precision data of Zr+Zr & Ru+Ru multiplicity-dependent cumulants and their higher order ratios fit into p+p and Au+Au results. All C_4/C_2 , C_5/C_1 , and C_6/C_2 show decreasing trend as multiplicity increases
2. Comparison with model and lattice QCD calculations
 - a. UrQMD over/underpredicts the results and in general, shows a similar trend as the data
 - b. **Higher order cumulant ratios at high multiplicity (top Au+Au collision centrality) consistent with the lattice QCD calculation result** (smooth crossover phase transition of thermalized medium)
3. Outlook: mixed cumulant ratios analysis to check a difference of B-field created in two isobar collisions
 - a. Expect $\sim 15\%$ difference in the B-field² between Zr+Zr & Ru+Ru collisions [1]
 - b. Mixed cumulant ratios offer an opportunity to experimentally assess the background B-field in the late stage of heavy-ion collisions [2]

[1] STAR, Phys. Rev. C 105, 014901 (2022)

[2] H.-T. Ding, Eur. Phys. J. A 57, 202 (2021)