

1 **Light nuclei production in Au+Au collisions at BES-II energies using the STAR**
2 **detector**

3 Sibaram Behera
4 (for the STAR Collaboration)

5 Indian Institute of Science Education and Research (IISER) Tirupati, India

6 In heavy-ion collisions, the production mechanism of light (anti-)nuclei can be explained by either
7 the thermal model or the coalescence model. By studying the yields and yield ratios of the light (anti-
8)nuclei, we can gain a better understanding of their production mechanisms and the properties of the
9 expanding system during freeze-out. Additionally, it has been proposed that the enhancement of light
10 nuclei compound ratios, e.g. $N_t N_p / N_d^2$ and $N_{^4He} N_p / N_{^3He} N_d$, with respect to the coalescence baseline
11 could serve as a probe of critical phenomena in the QCD phase diagram.

12 In the first phase of the Beam Energy Scan (BES-I) program at RHIC, an enhancement relative to the
13 coalescence baseline of the light nuclei yield ratio ($N_t N_p / N_d^2$) was observed in the most central Au+Au
14 collisions at $\sqrt{s_{NN}} = 19.6$ and 27 GeV, with a combined significance of 4.1σ . The precision of the
15 new measurements will be significantly improved by the large data sets ($\sim 10\times$ BES-I) obtained by the
16 STAR BES-II with upgraded detector capabilities.

17 In this talk, we will present the centrality and energy dependence of transverse momentum (p_T)
18 spectra of p , \bar{p} , d , \bar{d} , and 3He in Au+Au collisions at BES-II energies of $\sqrt{s_{NN}} = 7.7 - 27$ GeV. We will
19 also report on the centrality and energy dependence of integrated particle yields (dN/dy) and mean p_T
20 ($\langle p_T \rangle$) of light nuclei. Furthermore, we will discuss the centrality and p_T dependence of the coalescence
21 parameters ($B_2(d)$ and $B_3(^3He)$). The physics implications of these results will be discussed.