

**INPC 2025**

# **Femtoscopy of Strange Baryons in Heavy-ion Collisions at RHIC-STAR**

Boyang Fu (for the STAR Collaboration)

Central China Normal University

INPC 2025

In part supported by




U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science


1. Motivation
2. Femtoscopy
3. Lednicky-Lyuboshitz Approach
4. Analysis Details
5. Results
6. Summary

- Strange Dibaryons have never been found experimentally
  - Possible bound state: **H-dibaryon**  $\rightarrow \Lambda + \Lambda$  /  $\mathbf{p} + \Xi^-$ , **(Strange)Dibaryon**  $\rightarrow \mathbf{p} + \Omega^-$
- Hyperon-Nucleon (Y-N) and Hyperon-Hyperon (Y-Y) interactions provide important information to constrain the Equation-of-State and help to understand the inner structure of compact stars



Particle	Mass (MeV)	Quark composition	Decay mode
$f_0$	980	$q\bar{q}s\bar{s}$	$\pi\pi$
$a_0$	980	$q\bar{q}s\bar{s}$	$\pi\eta$
K(1460)	1460	$q\bar{q}q\bar{s}$	$K\pi\pi$
$\Lambda(1405)$	1405	$qqqs\bar{q}$	$\pi\Sigma$
$\Theta^+(1530)$	1530	$qqqq\bar{s}$	$KN$
H	2245	$uuddss$	$\Lambda\Lambda$
$N\Omega$	2573	$qqqsss$	$\Lambda\Xi$
$\Xi\Xi$	2627	$qqssss$	$\Lambda\Xi$
$\Omega\Omega$	3228	$ssssss$	$\Lambda K^- + \Lambda K^-$

Sungtae Cho, et al. (ExHIC), Phys. Rev. C 84, 064910 (2011)



- In high-energy experiment, Femtoscopy is inspired by Hanbury Brown and Twiss interferometry (Astronomy)
- Study the spatial and temporal extent of emission source
  - Quantum Statistics (Fermi-Dirac, Bose-Einstein)
  - Final-state Interactions (Coulomb, Strong interaction)
  - Collision Dynamics

• Two-particle correlation function:

<u>Statistical</u>	<u>Model</u>	<u>Experimental</u>
$C(k^*) = \frac{P(\vec{p}_a, \vec{p}_b)}{P(\vec{p}_a) \cdot P(\vec{p}_b)}$	$= \int d^3r^* S(r^*)  \Psi(r^*, k^*) ^2$	$= \mathcal{N} \frac{N_{same}(k^*)}{N_{mixed}(k^*)}$

>1: Attraction  
=1: No Correlation  
<1: Repulsion

$\vec{p}_a, \vec{p}_b$ : Single-particle momentum

$S(r^*)$ : Source function  
 $\Psi(r^*, k^*)$ : Pair wave function  
 PRF: Pair Rest Frame  
 $r^*$ : relative distance in PRF

$k^* = \frac{1}{2} |\vec{p}_a - \vec{p}_b|$ , relative momentum in PRF

$\mathcal{N}$ : Normalization factor  
 $N_{same}(k^*)$ : same event  
 $N_{mixed}(k^*)$ : mixed event

Nature 178 1046-1048(1956)  
ALICE Coll. Nature 588, 232–238 (2020)

$$CF(k^*) = \int d^3r^* S(r^*) |\psi(r^*, k^*)|^2$$

## ○ Assumptions:

- Smoothness approximation for source function
- Static and spherical Gaussian source
- Only consider s-wave
- Effective range expansion for  $\psi(r^*, k^*)$

## ○ Physics Parameters:

$R_G$ : Spherical Gaussian source

$f_0$ : Scattering length

$d_0$ : Effective range

## ○ Wave function:

$$\psi(r^*, k^*) = e^{-ik^*r^*} + f(k^*) \frac{e^{-ik^*r^*}}{r^*}$$


## ○ Scattering amplitude:

### ○ W/O Coulomb:

$$f_c(k^*) = \left[ \frac{1}{f_0} + \frac{1}{2} d_0 k^{*2} - ik^* \right]^{-1}$$


### ○ W/ Coulomb:

$$f_c(k^*) = \left[ \frac{1}{f_0} + \frac{1}{2} d_0 k^{*2} - \frac{2}{a_c} h(\eta) - ik^* A_c(\eta) \right]^{-1}$$



Relativistic Heavy Ion Collider (RHIC)  
Brookhaven National Laboratory, Upton

- p+p, Au+Au, Zr+Zr, Ru+Ru, d+Au...
- Beam Energy Scan Program I, II (Au+Au)

$$\sqrt{s_{NN}} = 3.0 - 200 \text{ GeV}$$


The Solenoidal Tracker At RHIC (STAR)

- Excellent particle identification
- Large, uniform acceptance at Mid-rapidity

○ Dataset:

Isobar collisions (Ru+Ru, Zr+Zr) @ 200 GeV

~ 3.9 billion minimum-bias events

Au+Au collisions @ 200 GeV (run11, run14, run16)

~ 2.5 billion minimum-bias events

Au+Au collisions @ 3 GeV (run21)







~ 2 billion minimum-bias events


○ Hyperon reconstruction via Helix-swimming method

$\Lambda \rightarrow p + \pi^-$ , BR = 63.9%

$\Xi^- \rightarrow \Lambda + \pi^-$ , BR = 99.9%

$\Omega^- \rightarrow \Lambda + K^-$ , BR = 67.8%






- CFs show enhancement at low  $k^*$
- Simultaneously fit with L-L function for 9 centralities (Au+Au and Isobar) to extract  $R_G, f_0$  and  $d_0$  by Bayesian method
- UrQMD + HAL QCD model is consistent with data

LL Fit	$f_0$ (fm)	$d_0$ (fm)	$\chi^2/\text{ndf}$
Ru+Ru			
Zr+Zr	$0.69^{+0.11}_{-0.10}$	$12.60^{+5.12}_{-7.00}$	<b>1.23</b>
Au+Au			

**p- $\Xi^-$  correlation**  
**STAR Preliminary**




- First experimental constraints of strong interaction parameters in p- $\Xi^-$  Paris in heavy-ion collisions

- Extracted spin averaged scattering length:

$$f_0 = 0.69^{+0.11}_{-0.10} \text{ fm (stat. + sys.)}$$


- **Attractive** strong interaction
- Shallow interaction compared to p-p interaction
- Consistent with HAL QCD prediction



- $\Lambda$ - $\Lambda$  correlation function shows suppression at small  $k^*$
- Compared with UrQMD + potential, it is found that the simulation with positive  $f_0$  is in better agreement with data  
=> Hints at an attractive interaction in  $\Lambda$ - $\Lambda$  pairs
- Need more precise data to confirm  
=> High statistics Isobar and Au+Au collisions

Potential	$f_0$ (fm)	$d_0$ (fm)	$\chi^2$ /ndf
NSC97a [1]	0.33	12.37	1.53
NF50 [2]	0.77	4.27	1.61
ND52 [3]	-23.96	2.59	2.24
ND50 [3]	10.63	2.04	4.02

[1] P. M. M. Maessen, et al, Phys. Rev. C 40 (1989) 2226  
 [2] M. M. Nagels, et al, Phys. Rev. D 20 (1979) 1633  
 [3] M. M. Nagels, et al, Phys. Rev. D 15 (1997) 2547




- Precise measurements of p- $\Omega^- \oplus \bar{p}-\bar{\Omega}^+$  correlation functions in Isobar collisions
  - CFs show enhancement at low  $k^*$  -> mainly due to Coulomb attractive interaction
  - CFs show depletion at  $k^* \sim 30-100 \text{ MeV/c}$  -> mainly due to the strong interaction

LL Fit	$\chi^2/\text{ndf}$
Spin ave.	0.65
Quintet	0.67

- Simultaneously fit with L-L function for 3 centralities to extract  $R_G, f_0$  and  $d_0$  by Bayesian method
- CFs obtained by HAL QCD theory with extracted  $R_G$  by L-L model is consistent with the data


Takumi Iritani, et al. (HAL QCD), Phys. Lett. B792 (2019)



- By taking CF ratio, Colomb effect can be largely canceled
- CF ratio shows enhancement at low  $k^*$  and depletion around  $k^* \sim 100$  MeV/c
  - Due to the presence of shallow bound state
- The potential  $V_{II}$ , with a p-value of 0.812, provides a better description of the data

Type	$f_0$ (fm)	$d_0$ (fm)	BE (MeV)	$\chi^2/ndf$	p-value	
$V_I$ [1]	1.12	1.16	--	1.66	0.014	No Bound
$V_{II}$ [2]	-3.38	1.31	2.15	0.76	0.812	Shallow Bound
$V_{III}$ [1]	-1.29	0.65	26.9	2.02	0.001	Deeply Bound

[1] Kenji Morita, et al., Phys. Rev. C 94, 031901 (2016)  
 [2] Kenji Morita, et al., Phys. Rev. C 101, 015201 (2020)



- First experimental constraints in heavy-ion collisions of strong interaction parameters in p- $\Omega^-$  pair

- Extracted **negative  $f_0$**  ( $|f_0| > 2d_0$ ) by Spin average method and Quintet method
  - **First experimental evidence of Strange Dibaryon**



- Calculate Binding Energy (BE) via Betha formula:

$$\text{Reduced mass: } m_{p\Omega} = \frac{m_p m_\Omega}{m_p + m_\Omega}$$

$$BE_{p\Omega} = \frac{1}{2m_{p\Omega}d_0^2} \left(1 - \sqrt{1 + \frac{2d_0}{f_0}}\right)^2$$

- Calculated BE are consistent with HAL QCD prediction



	Spin ave.	Quintet	HAL QCD
$f_0$ (fm)	$-4.9^{+0.5}_{-0.7}$	$-4.3^{+0.4}_{-0.7}$	-3.4
$d_0$ (fm)	$2.3^{+0.4}_{-0.5}$	$1.5^{+0.5}_{-0.7}$	1.3
BE (MeV)	$1.5^{+1.1}_{-0.6}$	$1.6^{+1.4}_{-0.5}$	2.3



- Extracted positive  $f_0$  in  $p-\Xi^-$  pair -> **Weakly attractive interaction**
- Extracted negative  $f_0$  in  $p-\Omega^-$  pair -> **Support the formation of bound state**

- Extracted source size in  $p-\Xi^-$ ,  $p-\Omega^-$  pairs show a linear distribution (Centrality dependence  $R_G^{central} > R_G^{peripheral}$ )

- Measured the correlation function of  $p - \Xi^- \oplus \bar{p} - \bar{\Xi}^+$ ,  $\Lambda - \Lambda$  and  $p - \Omega^- \oplus \bar{p} - \bar{\Omega}^+$  at STAR
- Extracted strong interaction parameters through L-L model
  - First experimental results of  $p - \Xi^-$  interaction  $\rightarrow$  Weakly attractive interaction
  - $\Lambda - \Lambda$  hints at weakly attractive interaction
  - **First experiment evidence of Strange-Dibaryon in  $p - \Omega^-$  channel**
- Extracted Binding Energy  $BE = 1.6_{-0.5}^{+1.4}$  MeV in  $p - \Omega^- \oplus \bar{p} - \bar{\Omega}^+$  pair



*Thanks for your attention!*