

Supported in part by




Global polarization of Ξ hyperons in Au+Au collisions in the STAR experiment

Outline:


- Introduction
- Global hyperon polarization
- Motivation
- The STAR experiment
- Hyperon polarization measurements
- Results
- Conclusions

Egor Alpatov (for the STAR Collaboration)
National Research Nuclear University MEPhI



Introduction



- The Quark-Gluon Plasma (QGP) formed in non-central nucleus-nucleus collisions is associated with large angular momentum, that leads to vorticity in the medium
- Spin-orbit coupling aligns spin directions of produced particles along the direction of vorticity
 - Z.-T. Liang and X.-N. Wang, PRL94, 102301 (2005)
 - S. A. Voloshin, arXiv:nucl-th/0410089
- Another possible source of particle polarization is magnetic field, created in non-central collisions in the initial stage
 - D. Kharzeev, L. McLerran, and H. Warringa, Nucl.Phys.A803, 227 (2008)
 - McLerran and Skokov, Nucl. Phys. A929, 184 (2014)



Vorticity



- In non-central Heavy-Ion Collisions the initial collective longitudinal flow velocity depends on the velocity gradient:



$$\omega_y = \frac{1}{2} (\nabla \times v)_y \approx -\frac{1}{2} \frac{dv_z}{dx}$$

- For small polarization:

Becattini, Karpenko, Lisa, Upsal, Voloshin PRC95.054902 (2017)

$$P_\Lambda \approx \frac{1}{2} \frac{\omega}{T} + \frac{\mu_\Lambda B}{T}$$

$$P_{\bar{\Lambda}} \approx \frac{1}{2} \frac{\omega}{T} - \frac{\mu_\Lambda B}{T}$$



How to measure global polarization?


- Hyperons are “self-analyzing” due to weak decay properties:
 - Daughter baryons are preferentially emitted along parent spin direction

- Daughter baryons of hyperons with polarization (\vec{P}) follows the distribution:

$$\frac{dN}{d\Omega^*} = \frac{1}{4\pi} (1 + \alpha_H |\vec{P}| \cdot \widehat{\vec{p}}_b^*) = \frac{1}{4\pi} (1 + \alpha_H P \cos \theta^*)$$


- α_H - decay parameter, unique for each hyperon species
- $\widehat{\vec{p}}_b^*$ is the daughter baryon momentum in the parent frame
- Projection to the transverse plane can be measured:

$$P_H = \frac{8}{\pi \alpha_H} \frac{\langle \sin(\psi_1 - \varphi_p^*) \rangle}{Res(\psi_1)}$$
 - ψ_1 is first-order event plane angle (proxy for reaction plane)
 - ψ_1 and its resolution $Res(\psi_1)$ can be calculated with spectator's signal.
- Ξ global polarization could also be measured via its daughter Λ polarization with transfer factor $C_{\Xi\Lambda} = 0.944$




Motivation



- Global polarization of Λ hyperons was measured for $\sqrt{s_{NN}} = 3\text{-}200$ GeV at STAR
- P_H decreases with increasing collision energy
- Recent BES-II Λ global polarization studies shows no significant difference between Λ and $\bar{\Lambda}$ global polarization
- Theoretical calculations can quantitatively explain the energy dependence of the Λ polarization, but many of them fail to explain differential measurements
- Nowadays there is a growing interest to measure the global polarization of other hyperons such as Ξ .
- Ξ and Ω hyperons global polarization was measured in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV
- Ξ polarization may provide new input for global polarization and vorticity studies



PRC 108, 014910 (2023)



The STAR experiment



Detectors with their η acceptance:


Hyperon reconstruction:

- Time Projection Chamber
 $|\eta| \in [-1, 1]$
- iTPC increases TPC acceptance to
 $[-1.5, 1.5]$
- Time-Of-Flight
 $|\eta| \in [-0.9, 0.9]$

Event plane angle measurement:

- Beam-Beam Counter
 $|\eta| \in [3.3, 5.0]$
- Event-Plane Detector
 $|\eta| \in [2.1, 5.1]$
- Bigger EPD acceptance and granularity improves event-plane resolution ~ 1.5 times compared to BBC in BES-I

Experimental technique




- Event plane Ψ_1 is determined by detectors at forward rapidity where directed flow is large

$$\Psi_1 = \tan^{-1} \left(\frac{\sum w_i \sin(\phi_i)}{\sum w_i \cos(\phi_i)} \right), \text{ where } w_i \text{ is detector's tile ADC}$$

A. M. Poskanzer, S. A. Voloshin, PRC58.1671(1998)



- $Res(\Psi_1, \text{Full } \eta) = \sqrt{2 \langle \cos(\Psi_{1, \text{Forward } \eta} - \Psi_{1, \text{Backward } \eta}) \rangle}$
- EPD was used to determine event-plane angle (BBC for systematics)

Zyzak, Maksym, Kisel, Ivan, Kulakov, Igor, & Vassiliev, Iouri (2013).
The KFPARTICLE package for the fast particle reconstruction
in ALICE and CBM



Hyperon reconstruction performed via decay topology with KFPARTICLE technique


- Λ daughters identified via TPC and TOF
- Ξ were reconstructed via $\Xi \rightarrow \Lambda + \pi$




$$P_H = \frac{8}{\pi \alpha_H} \frac{\langle \sin(\psi_1 - \phi_p^*) \rangle}{Res(\psi_1)}$$

Ξ global polarization: $\sqrt{s_{NN}}=19.6$ GeV


Centrality dependence




η dependence







p_T dependence



- Direct polarization measurements are consistent with measurements via daughter decays
- Polarization increases with centrality
- No obvious pseudorapidity or p_T dependence



Ξ global polarization: $\sqrt{s_{NN}}=27$ GeV




- Direct polarization measurements are consistent with measurements via daughter decays
- Polarization increases with centrality
- Weak pseudorapidity or p_T dependence if any

Ξ global polarization results



Global hyperon polarization $\sqrt{s_{NN}}$ dependence

- Observed positive Ξ global polarization
- Experimental results for Ξ global polarization consistent with each other with given large uncertainties at different collision energies
- Both Ξ global polarization measurement methods are consistent within uncertainties
- Ξ and inclusive Λ global polarization are consistent within statistical uncertainties
- Global polarization of Ξ hyperons consistent with AMPT predictions



Conclusions



- We presented $\Xi + \bar{\Xi}$ global polarization measurements in Au+Au collisions at $\sqrt{s_{NN}} = 19.6$ & 27 GeV, which helps to understand QCD spin dynamics and vorticity of QGP medium
 - This information can be used in theoretical development
- $\Xi + \bar{\Xi}$ global polarization is comparable with $\Lambda + \bar{\Lambda}$ global polarization within uncertainties, indicating a global nature of polarization

Thank you for your attention!