

Measurement of Intermittency for Charged Particles in Au + Au Collisions at $\sqrt{s_{NN}} = 7.7 - 200$ GeV from STAR

Jin Wu (for the STAR Collaboration)

Central China Normal University

1 One of the main goals of RHIC beam energy scan (BES) program is to search for the signatures
2 of QCD critical point in heavy-ion collisions. It is predicted that the local density fluctuations
3 near the critical point exhibit power-law scaling, which can be probed with an intermittency
4 analysis of the scaled factorial moments, $F_q(M)$, for charged particles. The power-law behavior
5 of q^{th} order scaled factorial moments can be expressed as: $F_q(M) \sim F_2(M)^{\beta_q}$, where M is the
6 number of equally sized cells in one dimension of momentum space, and β_q is the intermittency
7 exponent. The scaling exponent, ν , related to the critical component can be derived from the
8 equation: $\beta_q \sim (q - 1)^\nu$. The energy dependence of ν could be used to search for the signature
9 of the QCD critical point. Similar measurements have been carried out by NA49 and NA61
10 experiments in heavy-ion collisions with different system sizes.

11 In this talk, we will present the scaled factorial moments ($F_q(M)$, up to sixth order) of charged
12 particles in Au + Au collisions at $\sqrt{s_{NN}} = 7.7 - 200$ GeV measured by STAR experiment in the
13 first phase of RHIC BES. Then, we will show the energy and centrality dependence of the
14 extracted ν values. The physical implications of these results will be discussed.