

Baryon Stopping and Associated Production of Mesons in Au+Au Collisions at $\sqrt{s_{NN}} = 3.0$ GeV at STAR

XXIXth International Conference on Ultra-relativistic Nucleus-Nucleus Collisions
Kraków, Poland

Benjamin Kimelman for the STAR Collaboration



University of California - Davis



Supported in part by
U.S. DEPARTMENT OF
ENERGY

Office of Science


UC DAVIS
UNIVERSITY OF CALIFORNIA



- 1 Introduction
- 2 Charged Hadron Yields
- 3 Particle Production
- 4 Conclusions

Motivation

- Models indicate maximum baryon density $\sim \sqrt{s_{NN}} = 7.7$ GeV¹, suggesting changing dynamics of system
- π^\pm , K^\pm , and p dN/dy in central collisions have been measured at the AGS^{2,3,4}, SPS^{5,6}, and SIS⁷
- High statistics data sets collected by STAR extend these measurements to investigate centrality dependence
 - Observed Coulomb potential as a function of centrality is non-spherical
 - Baryon stopping measurements across centrality indicate constant rapidity loss



¹ J. Randrup and J. Cleymans, Phys. Rev. C **74**, 047901 (2006)

² J. L. Klay *et al.* (E895 Collaboration), Phys. Rev. C **68**, 054905 (2003)

³ J. L. Klay *et al.* (E895 Collaboration), Phys. Rev. Lett. **88**, 102301 (2002)


⁴ L. Ahle *et al.* (E866 and E917 Collaborations), Phys. Lett. **B490**, 53 (2000)

⁵ C. Alt *et al.* (NA49 Collaboration), Phys. Rev. C **77**, 024903 (2008)

⁶ S. Afanasiev *et al.* (NA49 Collaboration), Phys. Rev. C **66**, 054902 (2002)

⁷ A. Forster *et al.* (KaoS Collaboration), J. Phys. G **28**, 2011 (2002)

STAR Fixed-Target Acceptance




Outline


- 1 Introduction
- 2 Charged Hadron Yields
- 3 Particle Production
- 4 Conclusions




Particle Identification



- Slices of dE/dx and m^2 are made in rapidity, $m_T - m_0$, and centrality bins
- Each slice is fit with one or more Gaussians to determine yield of particle of interest, accounting for contamination from other particle species




Charged Pion Spectra



- Pion yields extracted from fitting dE/dx from TPC and $1/\beta$ from barrel ToF
- Pion spectrum described well by double thermal function¹, which describes thermal production at high $m_T - m_0$ and production from Δ resonance at low $m_T - m_0$
 - Dashed red line shows where leading production mechanism changes


¹ J. Klay et al. (E895 Collaboration), Phys. Rev. C 68, 054905 (2003)

Charged Kaon Spectra




- Kaon yields extracted from fitting dE/dx from TPC and $1/\beta$ from barrel ToF
- ToF identification required, which limits low $m_T - m_0$ reach
- Kaon spectrum fit with m_T exponential function since Bose-Einstein enhancement and radial flow effects mostly cancel

Proton Spectra



- Proton spectra extracted from fitting dE/dx from TPC and m^2 from barrel ToF
- Proton spectra near midrapidity described well using blast-wave function assuming cylindrical expansion of fireball

Radial Flow




- Shape indicates largest pressure gradient at midrapidity, decreasing toward target/beam rapidity
- Centrality dependence shows the size of overlapping region affecting pressure gradient
- Will investigate rapidity dependence to understand tension with Bjorken model

Surface velocity with linear profile ($n = 1$) from Blast-wave fit with constant temperature for each centrality and assuming Bjorken model

Centrality	0-5%	5-10%	10-20%	20-30%	30-40%	40-60%	60-80%
T_{kin} (MeV)	64.6	63.6	64.6	68.4	72.4	76.7	67.9



Charged Meson Rapidity Density Distributions



- Yields obtained from spectrum and integration of fits to extrapolate to $m_T - m_0 = 0$

- Yields consistent with results from E895¹ and E866/E917²


¹ J. Klay et al. (E895 Collaboration), Phys. Rev. C 68, 054905 (2003)

² L. Ahle et al. (E866 and E917 Collaborations), Phys. Lett. B490, 53 (2000)



Proton Rapidity Density Distributions

- Yields obtained from spectrum and integration of fits to extrapolate to $m_T - m_0 = 0$
- Centrality dependence shows participant proton peak shifting away from midrapidity for more peripheral collisions, indicating less baryon stopping





Outline

- 1 Introduction
- 2 Charged Hadron Yields
- 3 Particle Production
- 4 Conclusions



Coulomb Potential in Central Collisions




- Pions are susceptible to the Coulomb potential of stopped baryons due to their small mass^{1,2}
- Model includes Bose-Einstein nature of pions and an effective potential to account for momentum distribution of stopped protons

¹ D. Cebra *et al.* [arXiv:1408.1369 [nucl-ex]]

² J. Adamczewski-Musch *et al.* (HADES Collaboration), [arXiv:2202.12750 [nucl-ex]]


Coulomb Potential Correlation with Proton Yield

- Across centrality, the extracted Coulomb potential tracks monotonically with proton dN/dy
- Indication of less charge deposited in interaction region for peripheral collisions, matching expectations about geometry and baryon stopping
- Power law fit indicates that protons are not acting as a spherical source¹



¹J. Adamczewski-Musch *et al.* (HADES Collaboration), [arXiv:2202.12750 [nucl-ex]]

Kaon Production in Central Collisions



- Results at $\sqrt{s_{NN}} = 3.0$ GeV follows trend seen in SIS, AGS, SPS, and STAR BES-I data
- K^-/K^+ ratio at midrapidity is affected by baryon stopping
 - Production of K^+ in association with Λ baryon
 - Changes in μ_B and μ_S could suppresses K^-/K^+ ratio at low collision energy

¹A. Forster *et al.* (KaoS Collaboration), J. Phys. G **28**, 2011 (2002)

²L. Ahle *et al.* (E866 and E917 Collaborations), Phys. Lett. **B490**, 53 (2000)


³C. Alt *et al.* (NA49 Collaboration), Phys. Rev. C **77**, 024903 (2008)

⁴S. Afanasiev *et al.* (NA49 Collaboration), Phys. Rev. C **66**, 054902 (2002)



⁵L. Adamczyk *et al.* (STAR Collaboration), Phys. Rev. C **96**, 044904 (2017)

Baryon Stopping

- Proton rapidity density is fit with shape determined by counting the number of collisions each participating nucleon takes part in a Monte-Carlo Glauber model, smoothed using a Gaussian kernel
- The stopping, δy , is defined as the shift of the participant proton peak from beam rapidity



Baryon Stopping Trends



- Stopping at $\sqrt{s_{NN}} = 3.0 \text{ GeV}$ is consistent with measurements at similar energies
- Average loss of 0.19 ± 0.01 units of rapidity per nucleon-nucleon collision⁶

¹ W. Reisdorf *et al.* (FOPI Collaboration), Nucl. Phys. A **848**, 366 (2010)

² J. Klay *et al.* (E895 Collaboration), Phys. Rev. Lett. **88**, 102301 (2002)

³ B. Back *et al.* (E917 Collaboration), Phys. Rev. Lett. **86**, 1970 (2001)

⁴ L. Ahle *et al.* (E802 Collaboration), Phys. Rev. C **60**, 064901 (1999)

⁵ C. Blume. (NA49 Collaboration) J. Phys. G **34**, S951 (2007)

⁶ F. Videbæk and O. Hansen, Phys. Rev. C **52**, 2684 (1995)

Outline

- 1 Introduction
- 2 Charged Hadron Yields
- 3 Particle Production
- 4 Conclusions



- Rapidity density distributions have been presented at $\sqrt{s_{NN}} = 3.0$ GeV for π^\pm , K^\pm , and p
- The Coulomb potential from stopped protons has a noticeable effect on the charged pion spectra that correlates with the proton yields and indicates a non-spherical source
- K^-/K^+ ratio at midrapidity is affected by baryon stopping
- Baryon stopping can be modeled as an average rapidity loss of 0.19 ± 0.01 for each binary collision by observing trends in centrality
- Future studies of other STAR Fixed-Target data will identify energy trends in these observables, which will provide a more complete picture of the transition from a baryonic to a mesonic regime



29TH INTERNATIONAL
CONFERENCE ON ULTRARELATIVISTIC
NUCLEUS - NUCLEUS COLLISIONS

APRIL 4-10, 2022

KRAKÓW, POLAND

Thank you!





Backup

The STAR Fixed-Target




- Target located at $z = 200$ cm
- Target is 0.25 mm thick - 1% interaction probability
- Target is held 2 cm below center of beam axis
- Collider filled with 12 bunches

$\sqrt{s_{NN}} = 3.0$ GeV Vertex Position




Vertex centered well on target position and out-of-time pileup peaks (offset by 5 cm) easy to separate

Event and Track Selection Cuts



- Event selection
 - Select on minimum-bias events (using mixture of EPD, BBC, and VPD triggers)
 - $198 < V_z < 202$ cm
- Track selection
 - Track projects back to the primary vertex (at target location)
 - Distance of Closest Approach (DCA) ≤ 3

Coulomb-Corrected Double Bose-Einstein



- Coulomb correction applied to Bose-Einstein distribution to account for change in energy of pion
- Change in shape of spectra and fits at low $m_T - m_0$ indicate correction is necessary to describe spectra properly