


The 19th International Conference on Strangeness in Quark Matter

May 17-21, 2021, sponsored by Brookhaven National Laboratory, Upton, New York



Supported in part by



U.S. DEPARTMENT OF
ENERGY

Office of
Science

Measurement of global polarization of Λ hyperons in Au+Au $\sqrt{s_{NN}} = 7.2$ GeV fixed-target collisions at RHIC-STAR experiment

Kosuke Okubo for the STAR collaboration

University of Tsukuba


Strangeness in Quark Matter Conference 2021

20, May, 2021



筑波大学
University of Tsukuba





Introduction

◆ In non-central collisions...



▸ The created matter should exhibit strong vorticity.


-Z.-T.Liang and X.-N. Wang, PRL94, 102301

▸ The strong magnetic field would appear in the initial state.


-D. Kharzeev, L. McLerran, and H. Warring, Nucl.Phys.A803, 227 (2008)

-McLerran and Skokov, Nucl. Phys. A929, 184 (2014)





Global polarization



▸ Large angular momentum transfers to the spin degrees of freedom:

- Particle and anti-particle's spins are aligned with angular momentum.

▸ Spin alignment by magnetic field:


- Particles and anti-particles get aligned in the opposite direction due to the opposite signs of their magnetic moments.

✓ Both are considered to contribute to the global polarization.

How to measure the global polarization?

◆ Parity-violating decay of hyperon

- ▶ Daughter proton preferentially decays along the Λ 's spin (opposite for anti- Λ).



- ▶ Polarization can be measured via the distribution of the azimuthal angle of the daughter proton (in the hyperon rest frame).

◆ Projection onto the transverse plane

$$P_H = \frac{8}{\pi\alpha_H} \frac{\langle \sin(\Psi_1 - \phi_p^*) \rangle}{\text{Res}(\Psi_1)}$$


- STAR, PRC76, 024915(2007)


α_H : decay parameter ($\alpha_\Lambda = 0.732 \pm 0.014$)

P.A. Zyla et al. (PDG), Prog. Theor. Exp. Phys.2020, 083C01 (2020).


Ψ_1 : 1st-order event plane

ϕ_p^* : azimuthal angle of the daughter proton in the Λ 's rest frame







Motivation




- ▶ Λ global polarization has been measured from $\sqrt{s_{NN}} = 2.4 \text{ GeV}$ to 5.02 TeV.
- ✓ Polarization increases at low collision energy.
- ✓ No significant difference between Λ and anti- Λ .
- ✓ Preliminary Λ polarization measurement at $\sqrt{s_{NN}} = 2.4 \text{ GeV}$ from HADES is consistent with zero.
- ▶ New analysis of global polarization at $\sqrt{s_{NN}} = 7.2 \text{ GeV}$ with fixed-target experiment.
- ✓ 209M events at 7.2 GeV > 4M events at 7.7 GeV (BES I).
(Good minimum bias events)






The STAR detector




- **Time Projection Chamber (TPC)**
 - Main tracking detector, $|\eta| < 1.0$, full azimuth
- **Time-Of-Flight (TOF)**
 - Particle identification, $|\eta| < 0.9$, full azimuth
- **Event Plane Detector (EPD)**
 - Event plane reconstruction, $2.1 < |\eta| < 5.1$




STAR fixed-target program



- ✓ The gold target was installed inside the vacuum pipe at $z = 2.0$ m.
- ✓ Target is 0.25 mm thick and $\sim 1\%$ interaction probability.
- ✓ 209M good minimum bias events for Au+Au with fixed-target experiment at $\sqrt{s_{NN}} = 7.2$ GeV.
- ✓ Mid-rapidity for 7.2 GeV is -2.02.



Event plane correlation and resolution



✓ First-order event plane was reconstructed by the following formula.

$$\Psi_1 = \tan^{-1} \left(\frac{\sum w_i \sin(\phi_i)}{\sum w_i \cos(\phi_i)} \right)$$


✓ Event plane resolution was calculated by 3-subevent method.


$$\begin{aligned} \langle \cos([\Psi_1^A - \Psi_1^B]) \rangle &= \langle \cos([\Psi_1^A - \Psi_1^{\text{true}}]) \rangle \langle \cos([\Psi_1^{\text{true}} - \Psi_1^B]) \rangle \\ &= \sigma_n^A \sigma_n^B \end{aligned}$$

$$Res(\Psi_1^A) = \sqrt{\frac{\langle \cos([\Psi_1^A - \Psi_1^B]) \rangle \langle \cos([\Psi_1^A - \Psi_1^C]) \rangle}{\langle \cos([\Psi_1^B - \Psi_1^C]) \rangle}}$$

- A : EPD most inner 4 rings
- B : TPC ($-0.5 < \eta < 0$)
- C : TPC ($-1.5 < \eta < -1.0$)

A. M. Poskanzer and S. A. Voloshin, Phys. Rev. C 58, 1671 (1998).





Λ reconstruction

- Charged particles can be identified via specific ionization energy loss in the TPC and mass estimated from TOF.

Proton

$$\sqrt{|n\sigma|} < 3$$

$$\sqrt{0.5 < m^2 < 1.5 \text{ (GeV/c}^2\text{)}^2}$$

Pion

$$\sqrt{|n\sigma|} < 3$$

$$\sqrt{-0.06 < m^2 < 0.1 \text{ (GeV/c}^2\text{)}^2}$$

Topological cut

$$\sqrt{p\text{-DCA}} > 0.4 \text{ cm}$$

$$\sqrt{\pi\text{-DCA}} > 1.6 \text{ cm}$$




$$\sqrt{p\text{-}\pi \text{ DCA}} < 1.1 \text{ cm}$$


$$\sqrt{\Lambda\text{-DCA}} < 0.8 \text{ cm}$$

$$\sqrt{\text{Decay length}} > 5.0 \text{ cm}$$


(Centrality 30-40%)

These values of topological cut are slightly tuned depending on centrality.





Collision energy dependence of P_H




✓ First measurement in Au+Au collisions at $\sqrt{s_{NN}} = 7.2$ GeV.




• Observed positive Λ global polarization!

• The result follows the global trend of the energy dependence.


- Λ global polarization increases at lower collision energies.





Centrality dependence of P_H




✓ Polarization increases in more peripheral collisions.



Rapidity dependence of P_H




Z.T.Liang et al., Chin.Phys.C 45 (2021) 1, 014102




✓ Polarization is expected to depend on rapidity but the prediction is different among models.

✓ The result does not show significant rapidity dependence within our acceptance.

✓ Polarization in large rapidity region can be explored in the future with iTPC and forward upgrade (2023+2025).



p_T dependence of P_H




✓ One might expect ...

-decrease at low p_T due to the smearing effect caused by scattering at the later stage of the collisions.

-decrease at high p_T due to jet fragmentation.

J. Adams et al. (STAR), PRC98, 14910 (2018)

✓ **No significant p_T dependence.**



Summary

Summary

◆ We presented Λ global polarization in Au+Au collisions at $\sqrt{s_{NN}} = 7.2$ GeV with fixed-target configuration.

- Positive polarization is observed.
- Λ polarization increases at lower collision energies.
- Increasing trend towards peripheral collisions as expected.
- There are no significant rapidity and p_T dependences.

Outlook

- Measurement of anti- Λ polarization.

✓ We completed the data taking of BES II + FXT.

- $\sqrt{s_{NN}} = 7.2$ GeV : 267M(now) \rightarrow 267M + about 300M.


Collider mode data at BES II

Beam Energy (GeV/nucleon)	$\sqrt{s_{NN}}$ (GeV)	μ_B (MeV)	Number Events Requested (Recorded)	Date Collected
13.5	27	156	(560 M)	Run-18
9.8	19.6	206	400 M (582 M)	Run-19
7.3	14.6	262	300 M (324 M)	Run-19
5.75	11.5	316	230 M (235 M)	Run-20
4.59	9.2	373	160 M (162 M)	Run-20+20b
3.85	7.7	420	100 M (100 M)	Run-21

Fixed-target mode data at BES II



Beam Energy (GeV/nucleon)	$\sqrt{s_{NN}}$ (GeV)	μ_B (MeV)	Run Time	Number Events Requested (Recorded)	Date Collected
31.2	7.7 (FXT)	420	0.5+1.1 days	100 M (50 M+112 M)	Run-19+20
19.5	6.2 (FXT)	487	1.4 days	100 M (118 M)	Run-20
13.5	5.2 (FXT)	541	1.0 day	100 M (103 M)	Run-20
9.8	4.5 (FXT)	589	0.9 days	100 M (108 M)	Run-20
7.3	3.9 (FXT)	633	1.1 days	100 M (117 M)	Run-20
5.75	3.5 (FXT)	666	0.9 days	100 M (116 M)	Run-20
4.59	3.2 (FXT)	699	2.0 days	100 M (200 M)	Run-19
3.85	3.0 (FXT)	721	4.6 days	100 M (259 M)	Run-18

Back up



Pile up rejection

✓ Pile up events are removed using TOF start timing(T0).





T0 = the average of Time of Flight
- Time of Flight of each particle

Step1 : Count number of pion and proton with T0 from -0.3 to 0.3 [nsec].


Step2 : The event where the number of pion and proton is far from average are removed as pile up events in each multiplicity bin.

▶ About 90% pile up events would be removed.







Extract the polarization signal



✓ Observed polarization is more sharply peaked near Λ mass and it dips on the sides mass peak.

The width of the invariant mass depends on the daughter's azimuthal emission angle relative to the Λ .





Extract the polarization signal



✓ Calculate polarization each $\phi_\Lambda - \phi_p^*$ bin.

▸ Invariant mass method

• The data was fitted with the following formula.


$$\langle \sin(\Delta\phi) \rangle^{\text{obs}} = (1 - f^{\text{Bg}}(M_{\text{inv}})) \langle \sin(\Delta\phi) \rangle^{\text{Sg}} + f^{\text{Bg}}(M_{\text{inv}}) \langle \sin(\Delta\phi) \rangle^{\text{Bg}}$$

$$\begin{cases} \Delta\phi = \Psi_1 - \phi_p^* \\ f^{\text{Bg}}(M_{\text{inv}}) = f(M_{\text{inv}}^{\text{Bg}}) / f(M_{\text{inv}}^{\text{obs}}) \end{cases}$$



✓ Observed polarization is described as follows.

$$\frac{8}{\pi \alpha_H R^1_{EP}} \left\langle \sin(\Psi_1 - \phi_p^*) \right\rangle^{\text{Sg}} = P_\Lambda^{\text{true}} + c v_1 \sin(\phi_\Lambda - \phi_p^*)$$



Systematic uncertainty

✓ Different topological cut (~1.6%)

- Ten different cuts are applied.

• p-DCA ± 0.1 cm

• π -DCA ± 0.1 cm

• p- π DCA ± 0.1 cm

• Λ -DCA ± 0.1 cm

• Decay length ± 0.5 cm

✓ Method comparison for extracting polarization signal (~17.0%)

- Invariant mass method

- Event plane method

✓ Background assumption for polarization in the invariant mass method (~0.4%)

✓ Uncertainty from decay parameter α_H (~3.2%)

✓ Cumulant correction (~3.8%)

✓ Efficiency correction (~8.2%)