

# Investigating quantum interference in Drell-Söding process in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV at STAR

Xinbai Li (for the STAR Collaboration)  
University of Science and Technology of China

## Abstract

Relativistic heavy-ion collisions offer a unique environment for exploring quantum interference at an unprecedented femtometer scale through photon-nuclear interactions in photoproduction. In exclusive  $\pi^+\pi^-$  production, the resonance and continuum  $\pi^+\pi^-$  components arise from distinct production mechanisms in the  $\gamma\text{A}$  interaction. The continuum  $\pi^+\pi^-$  photoproduction is dominated by Drell-Söding process, in which a virtual  $\pi^+/\pi^-$  is diffraction-scattered on the nucleus. The  $\sim 5\%$  difference in the elastic scattering cross sections of  $\pi^- \text{A}$  and  $\pi^+ \text{A}$  around a  $\gamma\text{p}$  center-of-mass energy of approximately 12 GeV in photoproduction in 200 GeV Au+Au collisions at STAR may result in destructive interference. In contrast to  $\rho^0$  photoproduction, the Entanglement Enabled Spin Interference (EESI) in Drell-Söding process may differ due to the absence of the intermediate  $\rho^0$  and the specific dynamics of the virtual pion-nucleus interaction.

In this poster, we will present the first measurement of the diffractive  $p_T$  spectrum and the spin interference pattern through the amplitude of the second order final state angular cosine oscillation  $A_{2\Delta\phi}$  measurement for the Drell-Söding process in Au+Au collisions at  $\sqrt{s_{\text{NN}}} = 200$  GeV. The results indicate an obvious difference in  $p_T$  spectrum compared to  $\rho^0$  photoproduction. We also observe  $A_{2\Delta\phi}$  with no clear mass dependence for  $p_T < 0.1$  GeV/c and a notably stronger interference at the same  $M_{\pi^+\pi^-}$  in the Drell-Söding process compared to  $\rho^0$  photoproduction, which provides a unique opportunity to explore the effect of production mechanism on the EESI.