

1 **Nuclear modification factor of inclusive charged particles in Au+Au**
2 **collisions at $\sqrt{s_{NN}} = 7.7\text{--}27$ GeV with the STAR experiment and**
3 **comparing with URQMD and SMASH+vHLL model**

4 Aitbayev Alisher (for the STAR collaboration)

5 *Joint Institute for Nuclear Research (JINR)*

6 April 8, 2025

7 **Abstract**

8 The Quantum Chromodynamics (QCD) phase diagram, characterized by temperature
9 (T) and baryon chemical potential (μ_B), features a transition from hadronic matter to
10 a deconfined quark-gluon plasma (QGP). The Beam Energy Scan (BES) program at the
11 Relativistic Heavy Ion Collider (RHIC) explores this phase structure by systematically
12 varying the collision energy of Au+Au collisions, with a key focus on locating the QCD
13 critical point.

14 During the first phase (BES-I, 2010–2014), the STAR experiment measured the nuclear
15 modification factor (R_{CP}) for Au+Au collisions in energy range $\sqrt{s_{NN}} = 7.7\text{--}27$ GeV. In
16 2018, the STAR experiment initiated the second phase of the BES-II program, which has
17 a tenfold increase in statistics compared to the first phase. This will enable a deeper
18 understanding of the nuclear modification factor and improve its description. By 2021,
19 STAR collected 100 million Au+Au events at $\sqrt{s_{NN}} = 7.7$ GeV, two orders of magnitude
20 larger than the BES-I dataset at this energy.

21 In this talk, we present new measurements of charged-particle production and R_{CP}
22 from the high-statistics BES-II 7.7 GeV data, comparing them with BES-I results. We fur-
23 ther evaluate theoretical descriptions using URQMD and hydrodynamic (HYDRO) model
24 predictions, testing their consistency with experimental observations. By extending the
25 analysis to higher transverse momenta (p_T), we probe potential jet quenching effects and
26 assess implications for QGP formation and properties at lower collision energies.