

Nuclear modification factor of inclusive charged particles in Au+Au collisions at $\sqrt{s_{NN}} = 7.7\text{-}27$ GeV with the STAR experiment

Aitbayev Alisher (for the STAR collaboration)

Joint Institute for Nuclear Research (JINR)

April 8, 2025

Abstract

The Quantum Chromodynamics (QCD) phase diagram, characterized by temperature (T) and baryon chemical potential (μ_B), features a transition from hadronic matter to a deconfined quark-gluon plasma (QGP). The Beam Energy Scan (BES) program at the Relativistic Heavy Ion Collider (RHIC) explores this phase structure by systematically varying the collision energy of Au+Au collisions, with a key focus on locating the QCD critical point.

During the first phase (BES-I, 2010–2014), the STAR experiment measured the nuclear modification factor (R_{CP}) of inclusive charged particles in Au+Au collisions in energy range $\sqrt{s_{NN}} = 7.7\text{--}27$ GeV. In 2018, the STAR experiment initiated the second phase of the BES program (BES-II), which has a tenfold increase in statistics compared to the first phase. This will enable a This will enable better precision R_{CP} measurements. By 2021, STAR collected 100 million Au+Au events at $\sqrt{s_{NN}} = 7.7$ GeV, two orders of magnitude larger than the BES-I dataset at this energy.

In this talk, we present new measurements of charged-particle production and R_{CP} from the high-statistics BES-II data at $\sqrt{s_{NN}} = 7.7$ GeV, comparing them with BES-I results. We further evaluate theoretical descriptions using UrQMD and hydrodynamic (SMASH+vHLL) model predictions, testing their description of the experimental observations. By extending the analysis to higher transverse momenta (p_T), we probe potential jet quenching effects and assess implications for QGP formation and properties at lower collision energies.