




DNP2022

Fall Meeting of the Division of Nuclear Physics
of the American Physical Society

Oct. 27 – 30, 2022

Hyatt Regency Hotel, New Orleans, LA



Measurements of W^+ / W^- cross-section ratio in pp collisions at STAR

Jae D. Nam

Temple Univ.

For the STAR collaboration

Supported in part by:




U.S. DEPARTMENT OF
ENERGY


Office of Science



Relativistic Heavy Ion Collider



RHIC energies, species combinations and luminosities (Run-1 to 22)




- RHIC continues to serve as the world's first and only polarized pp collider.
- Features pp collisions at $\sqrt{s} = 500/508/510 \text{ GeV}$ and $\sqrt{s} = 200 \text{ GeV}$.
- $pA \sqrt{s_{NN}} = 200 \text{ GeV}$ and $AA \sqrt{s_{NN}} = 3 \sim 200 \text{ GeV}$ collisions

- At RHIC, protons can be polarized either:
 - Longitudinally (along the direction of the beam)
 - Proton spin composition
 - Transversely (perpendicular to the beam)
 - 3D image of the proton
 - Or can be unpolarized (if we choose not to look at the polarization)
 - **Parton distribution functions**
 - Non-linear gluon effects




Physics case: \bar{d}/\bar{u} asymmetry

NuSea, Phys.Rev.D 64 (2001) 052002



- While the valence quark (d, u) structure of the proton is well determined, the anti-quark counterpart (\bar{d}, \bar{u}) is much less constrained.
- Non-diminishing asymmetry between the anti-quarks in the proton sea \bar{d}, \bar{u} is a purely non-perturbative phenomenon.
- The anti-quark ratio \bar{d}/\bar{u} is typically measured in Drell-Yan type experiments with deuterons.

SeaQuest, Nature 590 (2021) 7847, 561-565



- Inconsistencies among these measurements have been found, especially in the proton momentum fraction range $x > 0.2$.
- W measurements at RHIC may provide insight around the same x region at higher Q^2 .






W production in pp collisions

- W^\pm cross sections at LO
 - $d\sigma^{W^+} \propto u(x_1)\bar{d}(x_2) + u(x_2)\bar{d}(x_1)$
 - $d\sigma^{W^-} \propto \bar{u}(x_1)d(x_2) + \bar{u}(x_2)d(x_1)$

$$\rightarrow R_W = \frac{\sigma^{W^+}}{\sigma^{W^-}} \sim \frac{u(x_1)\bar{d}(x_2) + u(x_2)\bar{d}(x_1)}{\bar{u}(x_1)d(x_2) + \bar{u}(x_2)d(x_1)}$$

- At LO, momentum scale set by the W mass, $Q^2 \sim M_W^2$.
- Leptonic decay via $W \rightarrow ev$
 - $\frac{d\sigma(W^\pm \rightarrow ev)}{dp_{T,e}^2} \propto \frac{(1 \pm \cos \theta)^2}{M_W \cos \theta}$
 - $p_{T,e} \sim \frac{M_W}{2} \sin \theta$
 - Jacobian peak at $p_{T,e} \sim M_W/2$
 - $y_e \sim y_W + \frac{\ln 1 + \cos \theta}{\ln 1 - \cos \theta}$
 - Charge discrimination as a function of y_e .


- Key features in experiment
 - High p_T electron.
 - Electron/hadron discrimination needed.
 - Large imbalance in p_T due to missing ν .



P. M. Nadolsky and C.P. Yuan, Nucl.Phys. B666 (2003) 31.
Jae D. Nam


Solenoid Tracker At RHIC (STAR)

- For measurements of W bosons, it is important to achieve near 4π detector acceptance.
- Time Projection Chamber (TPC)
 - Acceptance of $|\eta| < 1.3$.
 - Provides tracking & PID.
- Electro-Magnetic Calorimeter
 - Barrel (BEMC): $|\eta| < 1$.
 - Endcap (EEMC): $1 < \eta < 2$.
 - Assists in electron/hadron discrimination.
 - Assists in electron charge discrimination.
- Luminosity monitoring & Vertexing
 - Beam-Beam Counter (BBC)
 - Zero Degree Counter (ZDC)
 - Vertex Position Detector (VPD)



- The W bosons detected in the TPC + BEMC (barrel region) arise from a kinematic region of $0.1 < x < 0.3$.
- EEMC provides coverage in the intermediate rapidity region $1 < \eta < 2$, extending the kinematic reach to $0.06 < x < 0.4$.

RHIC pp Run Overview




Run	\sqrt{s} (GeV)	L (pb^{-1})
2009	500	10
2011	500	25
2012	510	75
2013	510	250
2017	510	350
2022	508	450 (estimate)


- $L \sim 700 + 450 pb^{-1}$ of pp collisions with $\sqrt{s} \geq 500 GeV$ has been collected at STAR.
 - Initial measurement based on Run 2009 with $L \sim 10 pb^{-1}$. (STAR, PRD 85 092010)
 - Follow up study with Run 2011-2013 with $L \sim 350 pb^{-1}$ has been published. (STAR, PRD 103,012001)
 - Preliminary study based on Run 2017 with $L \sim 350 pb^{-1}$.
 - New dataset with $L \sim 450 pb^{-1}$.



W tagging method





TPC track extrapolated to
BEMC tower grid




- W bosons that undergo the leptonic decay process, $W \rightarrow e\nu$, are tagged.
- Imbalance in p_T due to the missing neutrino. High $\vec{p}_{T,bal}$ ($= \vec{p}_{T,e} + \Sigma \vec{p}_{T,recoil}$) events are selected.
- Unlike hadrons, electrons deposit their energy in a highly concentrated region in the EMC. This isolated electron energy deposit is selected with $E_T^{2 \times 2} / E_T^{4 \times 4}$.
- Charge separation from TPC + EMC ($Q_e \times E_T / p_T$).
- Although not in this measurement, full W kinematics can be reconstructed.
 - Used for measurements of Sivers effect.

Signal/background description



- Electroweak ($N_{Z \rightarrow ee}$, $N_{W \rightarrow \tau \nu}$) background
 - Z decays with one unidentified electron
 - Leptonic decay of τ^W
 - Estimated with MC (Pythia)
- QCD background
 - Due to the limited acceptance and kinematic coverage, imbalance in p_T may appear in QCD events.
 - Two methods employed to estimate their contributions:
 - Second EEMC (N_{EEMC})
 - Accounts for missing backward coverage ($-2 < \eta < -1$)
 - Estimated by mirroring the effect of existing EEMC in the forward direction.
 - Data-driven QCD (N_{QCD})
 - Remaining background contribution that passes the selection process.
 - Distribution obtained by using events that do not pass the $p_{T,bal}$ cut.

Efficiencies



- In the W cross-section ratio measurement, the expression of the ratio reduces to:

$$\sigma_{W^+}/\sigma_{W^-} = \frac{N_{obs}^+}{\epsilon^+ \int L dt} / \frac{N_{obs}^-}{\epsilon^- \int L dt}$$


$$= \frac{\epsilon^-}{\epsilon^+} \cdot \frac{N_{sig}^+ - N_{bg}^+}{N_{sig}^- - N_{bg}^-}$$

where ϵ represents the product of the efficiencies of our selection process.

- Lower efficiency in Run 2012 and 2013 (compared to Run 2011) due to higher material budget and event pile-up caused by higher instantaneous luminosity.
- Higher tracking efficiency in 2013 than in 2012 due to improvements in tracking algorithm.
- MC study suggests that the efficiency ratio ϵ^-/ϵ^+ is consistent with unity and the deviation from unity is taken as a source of systematic uncertainty.




Results



- Measurement with STAR 2011-2013 data set has been published (PRD 103 (2021) 1, 012001).
- Additional data set taken in 2017 has been analyzed and is in preliminary release.
- These measurements are consistent with each other within their uncertainties.




Results (continued)



- Shown here is the result from the combined STAR 2011-2013 + 2017 data set.
 - Represents combined statistics of $L \approx 700 \text{ pb}^{-1}$.
 - Overall good agreement with the PDF distributions.
 - Latest PDF sets will be included in the final paper.


Results (projection)



- Projection for STAR 2022 data set
 - Combined statistics $\sim 1 \text{ fb}^{-1}$
 - Pushes the measurement to the systematic limit.
 - Concludes 500/510 GeV pp program at STAR

Absolute cross sections

STAR, PRD 103 (2021) 1, 012001




- Measurement of the total cross sections.

$$\sigma_{W/Z}^{fid} = \frac{N_{sig} - N_{bg}}{\epsilon \int L dt}$$


$$\sigma_{W/Z}^{tot} = \sigma_{W/Z}^{fid} / A_{W/Z}$$

- Acceptance correction on 2011 sample ($\sqrt{s} = 500 \text{ GeV}$) to match 2012 and 2013 samples ($\sqrt{s} = 510 \text{ GeV}$) by using FEWZ [PRD 86 (2012) 094034].



Z reconstruction

- The leptonically decaying $Z \rightarrow e^+e^-$ bosons are tagged by looking for electron-positron pairs.
- Additional selection process based on the reconstructed mass M_Z to reject $\gamma^* \rightarrow e^+e^-$ processes.




- Results with 2017 data set in progress.




PDF impacts

JAM, PRD 104 (2021) 7, 074031



S.Park, DIS2021



- Recent publication (STAR 2011+2012+2013) has been included in global fits.
- STAR data have a moderate impact on the sea quark distributions around $x \sim 0.2$.



Summary & Outlook

- W^+/W^- cross-section ratio has been measured with STAR pp $\sqrt{s} = 500, 510$ GeV datasets.
 - Probe \bar{d}/\bar{u} asymmetry in the proton sea, complementary to Drell-Yan measurements.
 - Results based on STAR 2011+2012+2013 ($L \approx 350$ pb^{-1}) have been published.
 - STAR 2017 (adds additional $L \approx 350$ pb^{-1}) dataset in preliminary state.
 - Combined results consistent with the current PDF distributions.
 - Global fit analyses confirm constraining power in the valence region.
- Outlook
 - STAR 2017 results nearing publication.
 - Measurement will reach systematic limit upon inclusion of new STAR 2022 dataset.



