

Measurements of Local Parton Density Fluctuations via Proton Clustering from STAR Beam Energy Scan

Dylan Neff
(For the STAR Collaboration)

UCLA

1 Quark-Gluon Plasma (QGP), a novel state describing the bulk properties of
2 QCD matter at high energies, can be experimentally probed with relativistic
3 nucleus-nucleus collisions. The nature of the transition between the QGP phase
4 and the final state hadron gas phase is yet to be established. The Beam Energy
5 Scan (BES) program at RHIC aims at searches for a possible critical point in
6 the QCD phase diagram.

7 Local density fluctuations are a characteristic signature of a first-order phase
8 transition. Baryons, formed via the coalescence of quarks at hadronization,
9 could be sensitive to these local parton density fluctuations at the phase tran-
10 sition boundary. In order to explore this, the multiplicity distribution of pro-
11 tons is studied in the subvolumes obtained by partitioning the azimuthal phase
12 space. Mixed events are constructed to wash out any event-by-event signal but
13 to preserve background effects, and thus serve as a baseline. Measurements of
14 transformed proton multiplicity distributions in azimuthal partitions of Au+Au
15 collisions at $\sqrt{s_{NN}} = 7 - 62$ GeV will be presented in this poster as a function
16 of beam energy and azimuthal partition size. These results are compared with
17 those obtained from the AMPT and MUSIC+FIST model calculations.