

Determining the longitudinal double-spin asymmetry (A_{LL}) for π^0 production from STAR
2013 Endcap Calorimeter Data
Emily Nelson, Madison Wallner
For an Undergraduate CEU Poster Presentation
Valparaiso University (for the STAR Collaboration)

The Solenoidal Tracker at RHIC (STAR) located at Brookhaven National Laboratory uses longitudinally polarized proton-proton collisions to study the gluon spin contribution to the known proton spin of $\frac{1}{2} \hbar$. The relative contributions of the quarks and gluons to the spin of the proton remain uncertain. Using data from the 2013 longitudinally polarized proton-proton collisions we study the asymmetry of proton spin-dependent production of neutral pions (π^0 s) from these collisions. π^0 s rapidly ($8.5 \cdot 10^{-17}$ s) decay into 2 photons that are detected by the Endcap Electromagnetic Calorimeter. By comparing the number of π^0 s produced when protons collide with different helicities, the asymmetry of π^0 production (A_{LL}), which can be related to the contribution of the gluon spin to the spin of the proton, can be measured. The two-photon invariant mass spectrum is reconstructed and then fit using a skewed Gaussian function to represent the π^0 signal and a Chebyshev function to characterize the background. Various checks must be made to assure the quality of the data being analyzed. The status of this analysis will be presented.