



DNP2020

Fall Meeting of the Division of Nuclear Physics
of the American Physical Society

Oct. 29 – Nov. 1, 2020 *Now Virtual Meeting!*

~~Hyatt Regency Hotel, New Orleans, LA~~




Longitudinal Double-Spin Asymmetry for Inclusive and Di-Jet Production in Polarized Proton Collisions at $\sqrt{s} = 200$ GeV

Nick Lukow

For the STAR collaboration

30 October 2020



STAR



TEMPLE
UNIVERSITY®

Supported in part by:



U.S. DEPARTMENT OF
ENERGY

Office of
Science

DOE NP contract: DE-SC0013405

Proton Spin Composition




Jaffe-Manohar Spin Sum Rule: $S = \frac{1}{2} = \frac{1}{2} \Delta\Sigma + \Delta G + L$

Nucl. Phys. B 337, 509 (1990)

The spin of the proton is composed of the **spin of the quarks and antiquarks**, the **spin of the gluons**, and the **orbital angular momentum** of the quarks and gluons

Quark Spin Contribution: $\Delta\Sigma = \int (\Delta u + \Delta d + \Delta s + \Delta\bar{u} + \Delta\bar{d} + \Delta\bar{s}) dx$



Gluon Spin Contribution: $\Delta G = \int \Delta g(x) dx$


Gluon Polarization & Longitudinal Double-Spin Asymmetry




- Polarized gluon distribution function can be constrained by measuring the longitudinal double-spin asymmetry of jets (A_{LL})



$$A_{LL} = \frac{\sigma^{++} - \sigma^{+-}}{\sigma^{++} + \sigma^{+-}} \propto \frac{\Delta f_a \Delta f_b}{f_a f_b} \hat{a}_{LL}$$



Low x –
Poorly
constrained



$$\text{DSSV: } \int_{0.1}^1 \Delta g(x) dx = 0.126 \pm 0.023$$

Phys. Rev. D 100, 114027 (2019)

$$\text{NNPDF: } \int_{0.05}^{0.2} \Delta g(x) dx = 0.17 \pm 0.06$$



Nucl. Phys. B 887, 276 (2014)

Subprocess Sensitivity



- STAR observables are dominantly sensitive to qg and gg scattering.
- Because of this sensitivity, A_{LL} allows access to $\Delta g(x)$

$$A_{LL} = \frac{\sigma^{++} - \sigma^{+-}}{\sigma^{++} + \sigma^{+-}} \propto \frac{\Delta f_a \Delta f_b}{f_a f_b} \hat{a}_{LL}$$




Longitudinal Double-Spin Asymmetry




$$A_{LL} = \frac{1}{P_1 P_2} \cdot \frac{N_{++} - \left(\frac{L_{++}}{L_{+-}}\right) N_{+-}}{N_{++} + \left(\frac{L_{++}}{L_{+-}}\right) N_{+-}}$$

- N_{++} - Yield of jets/dijets when helicities are the same sign
- N_{+-} - Yield of jets/dijets when helicities are opposite sign
- $P_{1,2}$ - Polarization of Beam 1,2
- $\left(\frac{L_{++}}{L_{+-}}\right)$ - Relative Luminosity



RHIC/STAR



Jet and Dijet Results from STAR



Central Rapidity ($|\eta| < 1.0$)

- Jets at 200 GeV in 2009
[Phys.Rev.Lett. 115 \(2015\) no.9, 092002](#)
- Dijets at 200 GeV in 2009
[Phys.Rev. D95 \(2017\) no.7, 071103](#)
- Jets at 510 GeV in 2012
[Phys.Rev. D100 \(2019\) no.5, 052005](#)
- Dijets at 510 GeV in 2012
[Phys.Rev. D100 \(2019\) no.5, 052005](#)
- Jets at 510 GeV in 2013
[\(STAR Preliminary\)](#)
- Dijets at 510 GeV in 2013
[\(STAR Preliminary\)](#)
- Jets at 200 GeV in 2015
[\(STAR Preliminary\)](#)
- Dijets at 200 GeV in 2015
[\(STAR Preliminary\)](#)


Intermediate Rapidity ($0.8 < \eta < 2.0$)

- Dijets at 200 GeV in 2009
[Phys.Rev. D98 \(2018\) no.3, 032011](#)
- Dijets at 510 GeV in 2012
[\(Preliminary\)](#)
- Dijets at 510 GeV in 2013
[\(Work in Progress\)](#)

Central Jets at 200 GeV in 2009



Phys.Rev.Lett. 115 (2015) no.9, 092002




- Jet p_T is corrected to the parton-level which doesn't include underlying event or beam remnants
- These results provided a significant increase in statistical precision compared to earlier measurements made in 2006
- These results show systematically larger A_{LL} than the DSSV 2008 global fit
- These results strongly suggest a positive gluon polarization value after inclusion in DSSV14 and NNPDF1.1 fits

Central Dijets at 200 GeV in 2009



Phys.Rev. D95 (2017) no.7, 071103




- Dijets probe a narrower region in x

- Plotted vs dijet invariant mass:


$$M = \sqrt{s x_1 x_2}$$

- Two topologies shown sample different x ranges:


- $\text{Sign}(\eta_1) = \text{Sign}(\eta_2)$



- $\text{Sign}(\eta_1) \neq \text{Sign}(\eta_2)$




Phys. Rev. D95 (2017) no.7, 071103




Intermediate Dijets at 200 GeV in 2009


Phys.Rev. D98 (2018) no.3, 032011




- By extending to intermediate rapidity, dijets can probe an even lower region in x
- Plotted vs dijet invariant mass
- Three topologies shown sample different x ranges:



Phys. Rev. D98 (2018) no.3, 032011




Central Jets at 200 GeV in 2015



- Most recent central jet analysis performed using data collected in 2015
- Includes underlying event subtraction technique first used in the analysis of 2012 data
- Results are consistent with previously published STAR results at $\sqrt{s} = 200$ GeV
- The figure-of-merit for the 2015 data set is twice as large as that for the 2009 data set

Central Dijets at 200 GeV in 2015



- New preliminary result from an analysis performed using data collected in 2015
- Dijets can provide further constraints by probing narrower regions in x and constraining the functional form of $\Delta g(x)$
- Includes underlying event subtraction technique first used in the analysis of 2012 data
- Results are consistent with previously published STAR results at $\sqrt{s} = 200$ GeV
- This result and the inclusive jet result using 2015 data are currently being prepared for publication

Summary



\sqrt{s} (GeV)	RHIC Run	Central Jets ($ \eta < 1.0$)	Central Dijets ($ \eta < 1.0$)	Intermediate Dijets ($0.8 < \eta < 2.0$)
200	2006	Published $x > 0.05$		
200	2009	Published $x > 0.05$	Published $x > 0.05$	Published $x > 0.01$
200	2015	Preliminary $x > 0.05$	Preliminary $x > 0.05$	
510	2012	Published $x > 0.02$	Published $x > 0.02$	Preliminary $x > 0.004$
510	2013	Preliminary $x > 0.02$	Preliminary $x > 0.02$	Work in Progress $x > 0.004$

- Nearly twice as much data was collected in 2015 as compared to 2009
- Results from data collected in 2015 will provide a significant reduction in the gluon polarization uncertainty for $x > 0.05$
- For details on the first A_{LL} analysis of intermediate-rapidity dijets at $\sqrt{s} = 510$ GeV, see Joe Kwasizur's talk later in this session

Backup

Dijet Correlation Measurements




$$x_1 = \frac{1}{\sqrt{s}} (p_{T3} e^{\eta_3} + p_{T4} e^{\eta_4})$$

$$x_2 = \frac{1}{\sqrt{s}} (p_{T3} e^{-\eta_3} + p_{T4} e^{-\eta_4})$$

$$M = \sqrt{x_1 x_2 s}$$

$$\eta_3 + \eta_4 = \ln \frac{x_1}{x_2}$$


$$|\cos \theta^*| = \tanh \left| \frac{\eta_3 - \eta_4}{2} \right|$$



Underlying Event Subtraction



- Off-axis cones are created at the same η as the reconstructed jet but $\pm \frac{\pi}{2}$ in ϕ
- Then the underlying event density is determined by taking the off axis cone p_T divided by the cone area. The densities of the two cones are then averaged together.
- The correction applied to the reconstructed jet is determined by multiplying the underlying event density by the area of the reconstructed jet. This correction is subtracted from the p_T of the reconstructed jet.



$$\sigma_{UE} = \frac{1}{2} \left(\frac{p_{T,cone(plus)}}{Area_{cone}} + \frac{p_{T,cone(minus)}}{Area_{cone}} \right)$$

$$p_{T,corrected} = p_T - \sigma_{UE} \cdot Area_{Jet}$$