



One-dimensional pion femtoscopy in d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV from STAR

Eugenia Khyzhniak


(for the STAR collaboration)

National Research Nuclear University MEPhI




NUCLEUS-2020

Outline

- Motivation
- Femtoscopy
- Correlation functions and their fits
- Systematic uncertainty
- k_T dependence of R_{inv} and λ
- System comparison



Motivation



Examination of the spatial and temporal scales of the particle-emitting source is one of the ways to study the process of particle production.


M. Podgoretzky 1989 *Particles & Nuclei* 20 630-68

In small systems (like p+p or d+Au) a collision area size is sensitive to fluctuations of initial conditions. Therefore, the detailed nature of particle production becomes important.

A. Bzdak et al. 2013 *Phys. Rev. C* 87, 064906

C. Plumberg 2020 arXiv:2008.01709

Femtoscscopy




Homogeneity region

Extracted radii measure the homogeneity lengths of the source


Akkelin SV, Sinyukov YM. Phys. Lett. B356:525 (1995)

- Femtoscopy allows one to measure:
 - Size of the emission source
 - Source shape & orientation
 - Lifetime & Emission duration
- System expansion dynamics are influenced by:
 - Transport properties
 - Phase transition/Critical point
 - Initial-state event shape



Analysis technique

Schematic view



1)

Construction of the correlation function:

$$C(Q_{inv}) = \frac{A(Q_{inv})}{B(Q_{inv})}$$

$$Q_{inv} = \sqrt{(\vec{p}_1 - \vec{p}_2)^2 - (E_1 - E_2)^2}$$

- $A(Q_{inv}) - Q_{inv}$ distribution with Bose-Einstein statistics (and final-state interactions – Coulomb and strong)
- $B(Q_{inv}) - Q_{inv}$ distribution without it

Fit of the correlation function:

2)


$$C(Q_{inv}) = N \left(1 - \lambda + \lambda K_{Coul}(Q_{inv})(1 + G(Q_{inv})) \right) D(Q_{inv})$$

$$G(Q_{inv}) = e^{-q_{inv}^2 R_{inv}^2}$$

- N - normalization factor
- λ - correlation strength parameter
- K_{Coul} - Coulomb correction factor
- $D(Q_{inv}) = 1$ (in this analysis) – Non-femtoscopic correlations



The STAR experiment



➤ Colliding system:



➤ d+Au@200 GeV

➤ Pion identification:

➤ Time Projection Chamber (TPC) - main tracking detector, $|\eta| < 1.0$, full azimuth





Example of the correlation functions and fits



Gaussian fit assumption:

$$G(Q_{inv}) = e^{-q_{inv}^2 R_{inv}^2}$$


$$\vec{k}_T = \frac{\vec{p}_{1T} + \vec{p}_{2T}}{2}$$



Correlation functions and their fits look reasonable



Lorentzian fit assumption:

$$G(Q_{inv}) = e^{-q_{inv} R_{inv}}$$





Statistical and systematic uncertainty

- For almost all cases statistical uncertainty smaller than marker size
- Sources of the systematic uncertainty:
 - Selection criteria of the events (position of the primary vertex): < 5%
 - Selection criteria of the tracks (momentum of the tracks, tracking efficiencies): < 6%
 - Selection criteria of the pairs (two track effects – merging, splitting): < 2%
 - Fit range: < 3%
 - Coulomb radius: < 3%
- Plan to investigate single track momentum resolution



k_T dependence of R_{inv} and λ



- Radius decreases with increasing k_T
- Radius increases with increasing particle multiplicity

- Correlation strength parameter decreases with particle multiplicities
 - Influences of the resonances increases?



Summary

- Femtoscopic parameters were obtained for d+Au colliding system
- The k_T dependence of the R_{inv} shows the dynamic of the system and allows to probe the different regions of the homogeneity in d+Au system
- Radius increases with increasing particle multiplicity
- Correlation strength parameter decreases with particle multiplicities

Thank you for your attention!



Back-up slide

Selection criteria

Event cuts	Track cuts	Pair cuts	Pion TPC cuts
$ Z_{TPC} \text{ (cm)} < 40$	$N_{Hits} > 15$	$-0.5 < \text{Splitting Level (quality)} < 0.6$	$ n\sigma_{pion} < 2$
$\sqrt{X_{TPC}^2 + Y_{TPC}^2} \text{ (cm)} < 2$	$N_{Hits}/N_{HitsFit} > 0.51$	$0.15 < k_T \text{ (GeV/c)} < 1.05$	$ n\sigma_{other} > 2$
$ Z_{TPC} - Z_{VPD} \text{ (cm)} < 5$	$DCA < 2 \text{ cm}$	Average Separation of two tracks within TPC volume (cm) > 10	
	$ \eta < 0.5$	$-1.1 < \text{Fraction of Merged Hits (\%)} < 0.1$	
	$0.15 < p \text{ (GeV/c)} < 0.8$		