



International Conference on New Frontiers in Physics (ICNFP), 2022

Azimuthal anisotropy of strange and multi-strange hadrons in isobar collisions at $\sqrt{s_{NN}} = 200$ GeV

Priyanshi Sinha

(for the STAR Collaboration)

Indian Institute of Science Education and Research (IISER) Tirupati, India



Supported in part by the




U.S. DEPARTMENT OF
ENERGY

Office of
Science


The STAR Collaboration

<https://drupal.star.bnl.gov/STAR/presentations>






Reaction plane: z-x plane



$$\frac{dN}{d\phi} \propto \frac{1}{2\pi} \left[1 + \sum_{n=1}^{\infty} 2v_n \cos(n(\phi - \Psi_R)) \right]$$

$$v_n = \langle \cos(n(\phi - \Psi_R)) \rangle$$

- Sensitive to early times in the evolution of the system
- Useful in understanding the nuclear structure








- Study of elliptic flow in isobar collisions may help in understanding the deformation of the colliding nuclei
- v_2 of strange and multi-strange hadrons, having lower hadronic cross-section, gives direct information on initial state anisotropies


Check if:

$$\frac{(v_2)_{\text{Ru+Ru}}}{(v_2)_{\text{Zr+Zr}}} \stackrel{?}{=} 1$$



- System size dependence of the azimuthal anisotropy



G. Giacalone, J. Jia, and V. Somà, Phys. Rev. C 104, L041903 (2021)



M. S. Abdallah et al. (STAR Collaboration), Phys. Rev. C 105, 14901 (2022)



Dataset: Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV (2018)

- Particle identification using TPC and TOF
- Systematic uncertainty sources: Event and track selections, Topological selection, Functional fitting for yield extraction



- Azimuthal angle of event plane is defined as :


$$\Psi_2 = \left[\tan^{-1} \left(\frac{\sum_i w_i \sin(2\phi_i)}{\sum_i w_i \cos(2\phi_i)} \right) \right] / 2$$

- Event plane has been calculated in two different pseudo-rapidity windows 'a' ($-1.0 < \eta < -0.05$) and 'b' ($0.05 < \eta < 1.0$)


- The event plane resolution using η -sub event approach :

$$R = \sqrt{\langle \cos 2(\Psi_2^a - \Psi_2^b) \rangle}$$


- Resolution correction is applied to obtain the final v_2



*Statistical error within marker size





- K_s^0 , ϕ , Λ , and Ξ have been reconstructed using invariant mass method; topological cuts using Helix method
- Background reconstruction using event-mixing method for ϕ -mesons and rotational method for K_s^0 , Λ , and Ξ



Event plane method:


- Particle raw-yields as a function of $\phi - \Psi_2$ are fitted with a Fourier function for different p_T ranges to extract v_2 coefficients



**Vertical bars indicate statistical error and shaded boxes denote systematic errors*

- v_2 shows a mass ordering at low p_T in isobar collisions
- Baryon-meson splitting at $p_T > 2$ GeV/c
- v_2 values have similar p_T dependence in minimum bias Ru+Ru and Zr+Zr collisions


Centrality dependence of $v_2(p_T)$



*Vertical bars indicate statistical error and shaded boxes denote systematic errors


- Strong centrality dependence is observed for v_2 of K_s^0 , Λ , and $\bar{\Lambda}$, in both Ru+Ru and Zr+Zr collisions
- $v_2(p_T)$ increases from central to peripheral collisions

Centrality dependence of $v_2(p_T)$





*Vertical bars indicate statistical error and shaded boxes denote systematic errors

- Strong centrality dependence is observed for v_2 of ϕ , Ξ^- , and Ξ^+ in both Ru+Ru and Zr+Zr collisions
- $v_2(p_T)$ increases from central to peripheral collisions



Transverse kinetic energy (KE_T) = $m_T - m_0$


- NCQ scaling holds good within 10% uncertainties for (multi-)strange in both Ru+Ru and Zr+Zr collisions
 - Indicative of partonic collectivity in the system



- p_T -integrated elliptic flow increases from central to peripheral collisions
- Ratios of v_2 between Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV for K_s^0 , Λ and $\bar{\Lambda}$ deviates from unity by $\sim 2\%$ at central and mid-central collisions
 → May indicate nuclear shape and structure difference between the two isobars

**No tracking efficiency correction since the effect would be largely cancelled*
**Error in the ratio includes statistical and systematic uncertainties*


System size dependence (strange)



- v_2 of K_s^0 , Λ , and $\bar{\Lambda}$ in isobar collisions is smaller than in $^{197}\text{Au}+^{197}\text{Au}$ and $^{238}\text{U}+^{238}\text{U}$ collisions at higher p_T
- v_2 in isobar collisions is larger as compared to $^{63}\text{Cu}+^{63}\text{Cu}$ collisions at higher p_T

B. I. Abelev et al. (STAR Collaboration), Phys. Rev. C 77, 054901 (2008)
 B. I. Abelev et al. (STAR Collaboration), Phys. Rev. C 81, 044902 (2010)
 M. S. Abdallah et al. (STAR Collaboration), Phys. Rev. C 103, 064907 (2021)

System size dependence (multi-strange)



$$\text{Fitting function : } f_{v_2}(n) = \frac{an}{1 + e^{-(p_T/n-b)/c}} - dn$$

n : number of quarks; a, b, c, d : free parameters

- v_2 of ϕ is similar in the measured p_T range for different collision systems within uncertainties
- v_2 of Ξ is lower than $^{238}\text{U}+^{238}\text{U}$ collisions at higher p_T

B. I. Abelev et al. (STAR Collaboration), Phys. Rev. C 77, 054901 (2008)

B. I. Abelev et al. (STAR Collaboration), Phys. Rev. C 81, 044902 (2010)

L. Adamczyk et al. (STAR Collaboration), Phys. Rev. Lett. 116, 062301 (2016)




- Elliptic flow of K_s^0 , Λ , $\bar{\Lambda}$, ϕ , and Ξ has been measured using event plane method for Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV
- Strong centrality dependence of v_2 for all particles has been observed
- NCQ scaling holds good within 10% uncertainties for all particles in all centralities for the isobar collisions
- Elliptic flow ratio for Ru+Ru over Zr+Zr shows a deviation of nearly 2% in central and mid-central collisions
 - Maybe related to nuclear shape and structure difference between the two isobars
- v_2 of strange hadrons in isobar collisions
 - At high p_T : Smaller compared to Au+Au and U+U collisions, and larger compared to Cu+Cu collisions
 - At low p_T : Similar for all collision systems studied



Thank you for your attention!

v_2 vs centrality



*Vertical bars indicate statistical error and shaded box denote systematic errors

*Error in the ratio includes statistical and systematic uncertainties

- p_T -integrated elliptic flow increases from central to peripheral collisions
- Ratio of v_2 between Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV for charged hadrons are comparable within the current uncertainties