

1 **Probing novel baryonic Spin Hall Effect via measurement of local** 2 **spin polarization of Λ hyperons in STAR Beam Energy Scan**

3
4 Qiang Hu

5 Institute of Modern Physics, Chinese Academy of Sciences
6 (for the STAR Collaboration)

7 The Spin Hall Effect (SHE) is instrumental in investigating quantum effects in many-body systems.
8 Recently, theoretical calculations indicate that the gradient of baryonic chemical potential
9 (analogous to the electric field) can induce a sizable spin Hall current. At the RHIC Beam Energy
10 Scan (BES) energies, the sign as well as the pattern of energy dependence of the difference in the
11 second harmonic coefficient between polarization of Λ and Λ -bar hyperons, could be significantly
12 different with and without the baryonic spin Hall current. In this talk, we will present the first

13 measurement of second harmonic coefficients of Λ hyperons' spin polarization ($P_x^{\Lambda-\Lambda\text{-bar}} \sin(2\Delta\varphi)$,

14 $P_y^{\Lambda-\Lambda\text{-bar}} \cos(2\Delta\varphi)$ and $P_z^{\Lambda-\Lambda\text{-bar}} \sin(2\Delta\varphi)$, where $\Delta\varphi = \varphi_\Lambda - \Psi_2$) as a function of transverse
15 momentum, rapidity, and collision centrality in RHIC BES-II Au+Au collisions at $\sqrt{s_{NN}} = 14.6, 19.6$
16 and 27 GeV. This can serve as the first probe of the baryonic SHE in heavy-ion collisions.