

1 Probing strong magnetic-field and baryonic Spin Hall effect via spin polarization of
2 Λ and $\bar{\Lambda}$ in Au+Au collisions from the RHIC Beam Energy
3 Scan
4

5 Qiang Hu

6 Institute of Modern Physics, Chinese Academy of Sciences
7 (for the STAR Collaboration)
8

9 A significant global spin polarization of Λ hyperons in the first phase of RHIC Beam
10 Energy Scan (BES) provides evidence of vorticity of the QGP created in heavy-ion collisions. The
11 data also hint at a larger polarization of $\bar{\Lambda}$ than that of Λ , which is
12 expected from a strong late-stage magnetic field sustained by the medium. A decisive
13 experimental test of this splitting would be highly significant, as it could reveal valuable
14 information about the electric conductivity of the QGP [1]. Similarly, the local polarization of
15 Λ and $\bar{\Lambda}$ hyperons are predicted to be different due to the polarization
16 induced by the gradient of baryonic chemical potential (analogous to the electric field) and called
17 baryonic spin Hall effect. This effect is expected to be observable through the energy dependence
18 of the angular modulation of the net polarization [2], $P_{\{x,y,z\}}^{\text{net}} = P_{\{x,y,z\}}^{\Lambda} -$
19 $P_{\{x,y,z\}}^{\bar{\Lambda}}$, as measured by $P_{\{x,z\}}^{\text{net}} \sin\left(2\phi_{\Lambda} -$
20 $2\Psi_2\right)$ and $P_y^{\text{net}} \cos\left(2\phi_{\Lambda} - 2\Psi_2\right)$, where Ψ_2 is the
21 second-order event-plane.

22 We report the first measurement of splitting in global and local spin polarization between
23 Λ and $\bar{\Lambda}$ hyperons in heavy-ion collisions using data collected by the
24 STAR experiment during the second phase of the RHIC Beam Energy Scan (BES II) with the
25 upgraded STAR detectors. We present results of Λ global polarization as a function of
26 centrality, transverse momentum, and rapidity in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 19.6$ and
27 27 GeV. We also present local polarization measurements in Au+Au collisions at
28 $\sqrt{s_{\text{NN}}} = 7.7 \sim 27$ GeV. Our measurements provide important insights into the late-
29 stage magnetic field sustained by the QGP, as well as spin Hall currents possibly created in a
30 highly dense baryonic environment.

31 [1]L. McLerran, V. Skokov, Nucl. Phys. A 922, 184 (2014).

32 [2]B. Fu et al, arXiv: 2201.12970 (2022).