

# The centrality and energy dependence of the elliptic flow of light nuclei and hadrons in STAR

Rihan Haque  
(for the STAR Collaboration)  
NISER, Bhubaneswar, India

## Outline

- Introduction & motivation
- STAR experiment at RHIC
- Results
- Summary


Quark Matter  
Darmstadt

May 19-24, 2014







# Azimuthal anisotropy



Interactions  
↓  
Pressure (P)

→

$$y > x \longrightarrow \frac{\partial P}{\partial x} > \frac{\partial P}{\partial y}$$



Azimuthal distribution of produced particles can be described as a Fourier series. The second order coefficient,



$$v_2 = \langle \cos(2(\phi - \psi_R)) \rangle = \left\langle \frac{p_x^2 - p_y^2}{p_x^2 + p_y^2} \right\rangle$$

– Sensitive to early times in the evolution of the system

An estimate of  $\psi_R$ , namely Event Plane ( $\psi_2$ ) is calculated using produced particles in mid-rapidity.

*A. M. Poskanzer and S. A. Voloshin, Phys. Rev. C 58, 1671 (1998)*





✓ Particle anti-particle  $v_2$  shows difference.

→ How does the difference depend on centrality and energy?

✓ hadron  $v_2$  show constituent quark (NCQ) scaling.

➤ Nuclei are expected to form at a later stage due to their low binding energy

→ Can we expect mass number scaling of nuclei  $v_2$  ?

→ How does nuclei and anti-nuclei  $v_2$  compare?

→ Is there any centrality dependence of nuclei  $v_2$  ?

*J. I. Kapusta, Phys. Rev. C 21, 1301 (1980)*


*R. Scheibl, U. Heinz, Phys. Rev. C 59, 1585 (1999)*

*D. Molnár, S. A. Voloshin, Phys. Rev. Lett. 91, 92301 (2003)*

Figure ref: *Phys. Rev. C 75, 054906 (2007)*, *Phys. Rev. Lett. 99, 112301 (2007)*, *Phys. Rev. Lett. 98, 162301 (2007)*

# The STAR experiment

## The Solenoidal Tracker At RHIC (STAR)



### 1. Time Projection Chamber (TPC)

pseudo-rapidity window:  $-1.0 < \eta < 1.0$   
full azimuthal coverage.

### 2. Time of Flight (ToF)

pseudo-rapidity window:  $-0.9 < \eta < 0.9$   
full azimuthal coverage


Using TPC and ToF  $\pi$ ,  $K$ ,  $p$  can be identified up to  $p_T \sim 3.0$  GeV/c,

### Light nuclei identification using TPC


d, dbar, triton:  $p_T \sim 1.0$  GeV/c, and  ${}^3\text{He}$  up to 4.5 GeV/c

### Light nuclei identification using ToF

d, dbar, triton:  $p_T \sim 4.0$  GeV/c,




# Measurement of nuclei $v_2$



- ✓ Elliptic flow of  $d$ ,  $\bar{d}$ ,  $t$ ,  ${}^3\text{He}$ ,  ${}^3\bar{\text{He}}$  measured at mid-rapidity.
- ✓  $\eta$  sub-eventplane method was used with  $\eta$ -gap = 0.1

proton  $v_2$  from *Phys. Rev. C* 88, 014902 (2013)

# Mass ordering of $v_2$




→ Nuclei  $v_2$  shows mass ordering at low  $p_T$  similar to hadrons

hadron  $v_2$  from Phys. Rev. C 88, 014902 (2013)





# $v_2$ of triton ( $t$ ) and ${}^3\text{He}$



$\rightarrow v_2$  of  $t$  and  ${}^3\text{He}$  are of similar magnitude (within statistical uncertainty)

# $v_2$ of particles and anti-particles




- Nuclei and anti-nuclei shows similar magnitude of  $v_2$
- Statistics for anti-nuclei at beam energies below 19.6 GeV is too low to check this consistency.
- $\Delta v_2$  for 10-40% centrality is similar to minimum bias result
- Centrality dependence not observed in  $\Delta v_2$
- $\Delta v_2$  relative to proton  $v_2$  shows a centrality dependence

Note: The difference of  $d, \bar{d}$   $v_2$  for 200 GeV in this slide are different than that shown in Quark Matter 2014 conference.

$v_2^{norm} = v_2$  of proton





# Centrality dependence of nuclei $v_2$



→ Nuclei  $v_2$  shows centrality dependence for all energies


# NCQ scaling of hadron $v_2$



- ✓ NCQ scaling observed for particle and anti-particle groups separately for beam energy  $\geq 19.6$  GeV
- ✓ Scaling holds for  $1.5 < p_T < 5.0$  GeV/c
- ✓ More statistics is needed for 7.7 and 11.5 GeV/c

Figures: *Phys. Rev. C* 88, 014902 (2013)

# Precision measurement of $v_2$ of $\phi$ and $\Omega$




- ✓ Mass ordering observed for  $p_T < 2.0$  GeV/c
- ✓ Baryon – meson splitting for  $2.0 < p_T < 5.0$  GeV/c

→ *High precision measurement of  $\phi$  and  $\Omega$   $v_2$  agree with the previous physics conclusion of partonic collectivity at 200 GeV*




# Mass number scaling of $v_2$




Nuclei  $v_2$  show mass number scaling for  $p_T / A \sim 1.5$  GeV/c for all beam energies  
 $\rightarrow$  Support the general idea that nuclei are formed by coalescence of nucleons

(anti-) proton  $v_2$  from Phys. Rev. C 88, 014902 (2013)



# Coalescence model results




*Coalescence model agrees with data*

*→ Another indication of coalescence of nucleons to form nuclei*

- ✓ Probability for producing a nucleus is given by the overlap of nucleon phase-space distribution with the Wigner phase-space function of nucleons inside the nuclei.
- ✓ Nucleon phase space information used from a transport (AMPT) model.

*R. Mattiello et al. Phys. Rev. Lett. 74, 2180 (1995), L. W. Chen et al. Phys. Rev. C 68, 017601 (2003)  
AMPT model: Zi-Wei Lin et al. Phys. Rev. C 72, 064901 (2005)*



# Summary

## (A) New Measurement presented:

- ✓ Energy ( $\sqrt{s_{NN}} = 7.7, 11.5, 19.6, 27, 39, 62.4$  and 200 GeV) and centrality dependence of nuclei  $v_2$  presented.
- ✓ Centrality dependence of difference in  $v_2$  of proton and anti-proton at  $\sqrt{s_{NN}} = 7.7, 11.5, 19.6, 27, 39, 62.4$  presented.

## (B) Observation and Physics conclusion:

1. Nuclei  $v_2$  versus  $p_T$  shows a clear centrality dependence and mass ordering when compared to identified hadrons at all beam energies studied  
→ *Mass ordering of  $v_2$  occurs naturally in a hydrodynamic model.*
2. Nuclei  $v_2$  versus  $p_T$  shows mass number scaling upto  $p_T/A = 1.5$  GeV/c and the magnitude of nuclei  $v_2$  versus  $p_T$  are reproduced by a Coalescence model.  
→ *Both these support the physics picture of coalescence of nucleons as the dominant mechanism of nuclei production.*
3. The difference in  $v_2$  of proton and anti-proton is observed to be similar at all collision centralities studied for the BES energies. A centrality dependence appears when this difference is normalized to proton  $v_2$  at the respective beam energies  
→ *The results implies hadronic interactions play an important role at lower beam energies.*





## Other interesting results on flow from STAR (Posters):

Triangular Flow of Identified Hadrons in Au+Au Collisions at  $\sqrt{s_{NN}} = 39$  and 200 GeV

- Xu Sun (*Poster Id: H-37*)

Measurement of higher harmonic flow of  $\phi$  meson in STAR at RHIC


- Mukesh Sharma (*Poster Id: H-03*)

# Thanks..

### Acknowledgements:

STAR Collaboration, NERSC Grid (LBNL), RCAS Grid (BNL),  
VECC TIER2 Grid (VECC), KONARK Grid (NISER).



Rihan Haque is supported by DAE-BRNS project grant No. 2010/21/15-BRNS/2026.



Back up is here...


Anti-particles:

particles:





# Baryon meson ratio



$v_2$  vs.  $m_T - m_0$  data fitted

$v_2$  baryon to  $v_2$  meson ratio taken at  $m_T - m_0 = 2.0$  GeV/c for baryons and (2/3) of that value for mesons

- Splitting larger for particles than for anti-particles
- Centrality dependence only for anti-particles
- No energy dependence