


D⁺ production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV measured by the STAR experiment



Yuanjing Ji, for the STAR Collaboration
University of Science and Technology of China
State Key Laboratory of Particle Detection and Electronics



Abstract


One of the goals of heavy-ion collisions is to search for the Quark-Gluon Plasma (QGP) and study its properties. Due to their large masses, heavy quarks are mainly produced in the initial hard scatterings during the early stage of heavy-ion collisions and experience the entire space-time evolution of the system. At the STAR experiment, utilizing high-precision secondary vertex reconstruction provided by the Heavy Flavor Tracker (HFT), D⁰ mesons have been comprehensively studied to investigate the charm quark transport in the QGP. Measurement of D⁺ production is complementary to the D⁰ measurement in studying the medium modification to the open charm meson production. It also provides useful information on feed-down contributions to the D⁰ yields. In this poster, measurement of D⁺ production at mid-rapidity ($|y| < 1$) in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV is reported. D⁺ are reconstructed via the hadronic decay channel (D⁺ → D⁰π⁺, D⁰ → K⁻π⁺, and its charge conjugate channel) utilizing the STAR HFT detector. The invariant yields of D⁺ and the ratios of D⁺/D⁰ yields are shown as a function of transverse momentum in different centralities.

Motivation

- Charm mesons as a sensitive probe of QGP via energy loss measurements;
- Study the D⁺/D⁰ ratio;
 - c → D⁰ (61.41% ± 0.73%), c → D^{+/−} (23.86% ± 0.46%) [1];
 - D⁺ feed-down contribution to D⁰ yields;


$$D^{*+} \rightarrow D^0 \pi^+$$
 - Hot medium effects:
 - D⁺ life time could become shorter in hot medium;
 - Regeneration and re-scattering [2].

Resonance	K*(892)	D ⁺ (2010)
Decay channel	Kπ	D ⁰ π
Branching Ratio %	~100	67.7
Width	50.7 MeV	83.3 KeV
Life time	4 [fm/c]	~2[pm/c]




Reconstruction Efficiency

- D⁺ efficiency
 - D⁰ efficiency ⊗ π_{soft} efficiency;
 - Vertex resolution correction;
- D⁰ efficiency
 - D⁰ reconstruction efficiency ← data-driven simulation;
 - Mass cut efficiency ← Real data D⁰ signal;
- π_{soft} efficiency
 - TPC tracking efficiency ← TPC embedding;
 - TOF matching efficiency ← Real data;
 - PID efficiency ← Extracted using the pure pion sample from K_s⁰ decay.




STAR Detector




Heavy Flavor Tracker
Inner tracking system (2014-2016):
 ◇ Silicon Strip Detector: r ~22 cm
 ◇ Intermediate Silicon Tracker: r ~14 cm
 ◇ PIXEL detector: r ~2.8 & 8 cm, MAPS, 20.7x20.7 μm², 0.5%X₀ thick, air-cooled

Systematic Uncertainties



Results



- ✓ D⁺/D⁰ ratio in Au+Au collisions at 200 GeV is consistent with that in Pb+Pb collisions at 5.02 TeV [4].
- ✓ The p_T dependence of D⁺/D⁰ ratio shows similar trend in different centrality bins.
- ✓ D⁺/D⁰ ratio as a function of p_T is consistent with PYTHIA8 (STAR-HF Tune) prediction.
- ✓ Ratio of integrated yields (2 < p_T < 10 GeV/c) of D⁺ to D⁰ shows no strong dependence on centrality. No significant effect of hot medium on the D⁺ life time has been observed.



D⁺ reconstruction

- Dataset: Au+Au @ 200 GeV recorded in 2014; ~900 Million minimum-bias events.
- Reconstruction method
 - D⁺ → D⁰π_{soft}⁺ (B. R. = 67.7%),
 - D⁰ → K⁻π⁺ (B. R. = 3.89%),
 - and its charge conjugate channel.

- D⁰ reconstruction cuts:
 - $|y|_{D^0} < 1$;
 - K/π : p_T > 0.3 GeV/c;
 - K/π : $|\eta| < 1$;
 - K/π : at least one hit in each layer of PXL and IST;
 - K/π PID : if TOF available, TOF & TPC; otherwise TPC only.
- D⁰ topological cuts:

D ⁰ p _T (GeV/c)	0-1	1-2	2-3	3-5	5-15
decay length (μm) >	145	181	212	247	259
DCA between 2 daughters (μm) <	84	66	57	50	60
DCA between D ⁰ and PV (μm) >	61	49	38	38	40
DCA between π and PV (μm) >	110	111	86	81	62
DCA between K and PV (μm) >	103	91	95	79	58

- Kπ invariant mass range for D⁰ candidates: 1.83 GeV/c² < M(Kπ) < 1.90 GeV/c²
- π_{soft} cuts:
 - DCA_{PV} ≤ 3 cm, not refitted with the PV;
 - At least 20 space points in the TPC, (no requirement to leave hits in HFT);
 - p_T > 0.15 GeV/c;
 - $|\eta| < 1$;
 - PID: TOF and TPC if TOF is available, otherwise TPC only.



- ✓ Background is estimated by a mixed-event method.

Summary

- ✓ D⁺ p_T spectra and D⁺/D⁰ ratio have been measured for different centralities of Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV;
- ✓ The dependence of D⁺/D⁰ ratio on p_T is similar in different centrality bins, and is compatible to that in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV.
- ✓ Ratio of integrated yields (2 < p_T < 10 GeV/c) of D⁺ to D⁰ shows no strong dependence on centrality. No significant effect of hot medium on the D⁺ life time has been observed.

Reference

- [1] M. Lisovsky, et al., Eur. Phys. J. C (2016) 76: 397.
- [2] M. M. Aggarwal et al. Phys. Rev. C (2011) 84.3: 034909.
- [3] L. Adamczyk et al. Phys. Rev. Lett. (2017) 118.21: 212301.
- [4] ALICE Collaboration. arXiv:1804.09083

