- ¹ Deuteron number fluctuations and proton-deuteron correlations in
- ² high energy heavy-ion collisions in STAR experiment at RHIC
- 3

Debasish Mallick

(for the STAR Collaboration)

National Institute of Science Education and Research, HBNI, Jatni-752050, INDIA

XXIXth International Conference on Ultra-relativistic Nucleus-Nucleus Collisions

The production mechanism of deuterons, which have a binding energy of 2.2 MeV, is a topic of current interest in high energy heavy-ion collisions, where the system undergoes kinetic freeze-out at temperatures around 100 MeV. Two possible scenarios include (a) statistical thermal process and (b) coalescence of nucleons. Cumulants of deuteron number distributions and proton-deuteron correlations are sensitive to these physics scenarios. In addition, they are also sensitive to the choice of canonical versus grand canonical ensemble in statistical thermal models.

We report the first systematic measurements of collision energy and centrality depen-11 dence of cumulants (up to fourth order) of deuteron number distributions in Au+Au colli-12 sions at $\sqrt{s_{NN}} = 7.7, 11.5, 14.5, 19.6, 27, 39, 54.4, 62.4, and 200 GeV. We will also discuss$ 13 new measurements on proton-deuteron correlations. The measurements are performed in 14 the STAR experiment at mid-rapidity (|y| < 0.5) and within transverse momentum range 15 $0.8 < p_T(\text{GeV}/c) < 4.0$, using Time Projection Chamber and Time-of-Flight detectors. 16 The experimental results are compared to the statistical thermal model calculations with 17 a grand canonical, canonical ensemble, and the UrQMD model that incorporates the coa-18 lescence of nucleons close by in space and momentum to form deuterons. These theoretical 19 comparisons with the experimental measurements provide key insights into the mechanism 20 of deuteron production in high-energy heavy-ion collisions. 21