




# Beam energy and collision species dependences of photon-induced lepton pair production at STAR

---



Xiaofeng Wang (王晓凤)

For the STAR Collaboration  
Shandong University (山东大学)

---



# $\gamma\gamma \rightarrow l^+ l^-$ Process




- Highly Lorentz-contracted charged nuclei produce electromagnetic fields (EM)
- Equivalent Photon Approximation (EPA): EM fields  $\rightarrow$  a flux of **quasi-real photons**  
*Weizsäcker, C. F. v. Zeitschrift für Physik 88 (1934): 612*
- 1934 Breit & Wheeler : “Collision of two Light Quanta”  
*G. Breit and J. A. Wheeler. Physical Review 46 (1934): 1087*
- High photon density with highly charged nuclei ( $\propto Z^2$ )

# $\gamma\gamma \rightarrow l^+l^-$ in Peripheral Collisions



STAR: Phys.Rev.Lett. 121, 132301 (2018)



Observation of  $\gamma\gamma \rightarrow e^+e^-$  in hadronic heavy ion collisions at STAR


Energy dependence?  
Centrality dependence?

Di-muon?

Collision species dependence?





# The Solenoid Tracker At RHIC (STAR) and PID




Time Projection Chamber (TPC): momentum and energy loss

Time Of Flight (TOF): velocity



# Transverse Momentum Distribution



$e^+e^-$  pairs

Excesses above hadronic production are observed at low- $p_T$

Lowest order EPA-QED predictions are consistent with observed excesses

Energy dependence  
**54.4 GeV**, 200 GeV

Centrality dependence  
 40-60%, 60-80%, **80-100%**



EPA-QED: W. Zha et al, Phys.Lett.B 800 (2020) 135089

# Transverse Momentum Distribution




$\mu^+ \mu^-$  pairs

Poster by Jian Zhou (04/06/22 6:30-7:30)



Similar excesses at low- $p_T$  observed in the  $\mu^+ \mu^-$  channel

# Invariant Mass Distribution at Low- $p_T$



$$\gamma\gamma \rightarrow e^+e^-$$

Excesses (Data - Cocktail) are extracted

No vector meson observed  
( $\gamma\gamma \not\rightarrow$  vector meson)


Excesses are well described by lowest order EPA-QED predictions

# Invariant Mass Distribution at Low- $p_T$



$$\gamma\gamma \rightarrow \mu^+\mu^-$$

Poster by Jian Zhou (04/06/22 6:30-7:30)



EPA-QED predicts different cross sections due to electron and muon mass difference

Excesses (Data - Cocktail) are extracted


Excesses are well described by lowest order EPA-QED predictions



# Energy Dependence of Excess Yield



$$\gamma\gamma \rightarrow e^+e^-$$




**Excess yield increase with beam energy**

**EPA-QED predicts similar energy dependence**

# Energy and Centrality Dependence of $\sqrt{\langle p_T^2 \rangle}$



$$\gamma\gamma \rightarrow e^+e^-$$




$\sqrt{\langle p_T^2 \rangle}$  is sensitive to  $p_T$  broadening

$\sqrt{\langle p_T^2 \rangle}$  decreases from semi-peripheral to peripheral collisions

Initial state effect: Impact parameter dependence

Energy dependence ( $3.7\sigma$  compared to 200 GeV QED) and/or final state effect ( $1.8\sigma$ )

# Application: Constrain Charge Distribution



$\gamma\gamma \rightarrow l^+l^-$  can be used to constrain nucleus charge distribution at RHIC energy

STAR data compared to EPA-QED

Low energy scattering:  $R=6.38$  fm,  $d=0.535$  fm

R. C. Barrett and D. F. Jackson, Nuclear Sizes and Structure (Oxford University Press, 1977)

200 GeV vs 54 GeV: maybe due to energy dependence of charge distribution


Low-energy vs RHIC ( $3\sigma$  difference): maybe due to energy dependence of charge distribution and/or final state effect

EPA-QED: J. D. Brandenburg et al, Eur. Phys. J. A 57 (2021) 299.

# Collision Species Dependence ( ${}^{96}_{44}\text{Ru} + {}^{96}_{44}\text{Ru}$ , ${}^{96}_{40}\text{Zr} + {}^{96}_{40}\text{Zr}$ )



$$\gamma\gamma \rightarrow e^+e^-$$



At very low  $p_T$  ( $< 0.15$  GeV/c),  $e^+e^-$  pairs dominated by  $\gamma\gamma \rightarrow e^+e^-$



Ratio is consistent with  $(\frac{44}{40})^4$  at very low  $p_T$

**Initial EM field is different in Ru + Ru and Zr + Zr ( $\sim 3\sigma$ )**

At  $p_T > 0.15$  GeV/c, hadronic production contributions to  $e^+e^-$  pairs are similar in Ru + Ru and Zr + Zr

Poster by Kaifeng Shen (04/06/22 6:30-7:30)

# Collision Species Dependence ( ${}^{96}_{44}\text{Ru} + {}^{96}_{44}\text{Ru}$ , ${}^{96}_{40}\text{Zr} + {}^{96}_{40}\text{Zr}$ )




Poster by Kaifeng Shen (04/06/22 6:30-7:30)

- At very low  $p_T$ ,  $J/\psi$  dominated by  $\gamma A \rightarrow J/\psi$
- Ratio is consistent with  $\left(\frac{44}{40}\right)^2$  at very low  $p_T$
- Initial EM field is different in Ru + Ru and Zr + Zr ( $\sim 1.7\sigma$ )
- At  $p_T > 0.2$  GeV/c, hadronic production contributions to  $J/\psi$  are similar in Ru + Ru and Zr + Zr



# Collision Species Dependence ( $^{96}_{44}\text{Ru} + ^{96}_{44}\text{Ru}$ , $^{96}_{40}\text{Zr} + ^{96}_{40}\text{Zr}$ )



$$\gamma A \rightarrow J/\psi$$

$$J/\psi \text{ excess yield} \propto Z^2$$

Photoproduced  $J/\psi$  yields seem to be independent of form factor and impact parameter

Poster by Kaifeng Shen (04/06/22 6:30-7:30)

Published Results: J. Adam et al. (STAR Collaboration), Phys.Rev.Lett. 123 (2019) 132302

EPA: W. Zha et al, Phys. Lett. B 789 (2019) 238–242

- Beam energy and centrality dependences of  $\gamma\gamma \rightarrow l^+l^-$  have been measured at STAR
  - ✓ Excess yield: **Increases with beam energy**
  - ✓  $\sqrt{\langle p_T^2 \rangle}$ : Decreases with increasing impact parameter
  - ✓  $\sqrt{\langle p_T^2 \rangle}$ : **Energy dependence** ( $3.7\sigma$  compared to 200 GeV QED)
  - ✓ **Application:**  $\gamma\gamma \rightarrow l^+l^-$  can be used to **constrain nuclei charge distribution at RHIC energy**
- Collision species dependence of  $\gamma\gamma \rightarrow l^+l^-$  and  $\gamma A \rightarrow J/\psi$  have been measured at STAR
  - ✓ **Initial EM field is different** in Ru + Ru and Zr + Zr ( $\sim 3\sigma$  in  $e^+e^- \oplus \sim 1.7\sigma$  in  $J/\psi$ )
  - ✓ Photon-induced  $J/\psi$  yield  $\propto Z^2$