

${}^4_{\Lambda}\text{He}$ Production in $\sqrt{s_{NN}} = 3$ GeV Au+Au Collisions

Fengyi Zhao

(for the STAR Collaboration)

October 9, 2024

Abstract

Hypernuclei, bound states of nuclei with one or more hyperons, serve as a natural laboratory to investigate the hyperon-nucleon (YN) interaction. High-baryon density environment can be created by heavy-ion collisions at a collision energy of a few GeV/u, which provides a unique opportunity to study the YN interaction and production mechanism of hypernuclei. Comparison of ${}^4_{\Lambda}\text{He}$ yield with its isobar ${}^4_{\Lambda}\text{H}$ may shed light on the isospin related Λ -t and Λ - ${}^3\text{He}$ interactions.

In this poster, we will present the new results on the ${}^4_{\Lambda}\text{He}$ differential yield as a function of rapidity and transverse momentum. The yield is measured in $\sqrt{s_{NN}} = 3$ GeV Au+Au 0-50% central collisions, and the data is collected by the STAR experiment with fixed-target mode. ${}^4_{\Lambda}\text{He}$ is identified via its three-body decay channel ${}^4_{\Lambda}\text{He} \rightarrow {}^3\text{He} + p + \pi^-$. We found that the rapidity distributions of ${}^4_{\Lambda}\text{He}$ are similar to those of ${}^4_{\Lambda}\text{H}$, and the differential yield ratios of ${}^4_{\Lambda}\text{He}/{}^4_{\Lambda}\text{H}$ are consistent with those of ${}^3\text{He}/t$ within uncertainties. The observed yield ratios can be described by both the JAM plus a coalescence afterburner and the canonical thermal model.