



Investigating system size dependence of strange hadron production at 200 GeV at STAR

Xiongxiong Xu (ailondx@rcf.rhic.bnl.gov)
Tsinghua University, for the STAR Collaboration



Abstract

There are significant discussions in the field about the initial conditions, including the size of the system, needed to generate a quark-gluon plasma (QGP). Strangeness production serves as a sensitive probe into the properties of the QGP. It is expected that the Ω/ϕ ratios in different colliding systems may reveal the minimum colliding system size required to produce QGP.

In this poster, we will present the transverse-momentum (p_T) spectra of strange hadron ($\phi, \Omega, \bar{\Omega}$) in isobar (Ru+Ru and Zr+Zr) and O+O collisions at $\sqrt{s_{NN}} = 200$ GeV at mid-rapidity ($|y| < |0.5|$) and the Ω/ϕ ratios in those colliding systems. The O+O system has the extended kinematic coverage benefit from the iTPC upgrade, which extended the rapidity coverage and enhanced the particle identification capability compared with previous results.


Motivation

- Update Ω/ϕ ratio as a function of p_T in different colliding systems to explore the minimum colliding system size required to produce QGP.
- Precise measurement of $\Omega(\bar{\Omega})$ yields to investigate the system size dependence of strangeness production.

STAR Detector


iTPC (STAR Inner Sector TPC Upgrade) :

- Larger rapidity coverage — $|y|$ from 1.0 to 1.5
- Better PID — improved dE/dx resolution
- Lower p_T limit — from 125 to 60 MeV/c
- Efficiency of strangeness reconstruction improved significantly





Analysis

- Dataset: Run18 isobar 200 GeV and Run21 O+O 200 GeV
- $\sim 730M$ minimum bias events for isobar
- $\sim 370M$ minimum bias events for O+O
- Particle identification with TPC(dE/dx) and TOF($1/\beta$)
- KFParticle method used in Ω reconstruction — efficiency improved at high p_T
- p_T region of Ω : $0.8 \sim 4.6$ GeV/c
- p_T region of ϕ : $0.4 \sim 5.0$ GeV/c
- Signal extraction: rotational (for Ω) and mix-event (for ϕ) for combinational background, polynomial fit for residual background & double gaussian fit for signal



ϕ Meson p_T Spectra

- Precise measurement for 7 centrality (parameter indicating how central the event is) bins
- Maximum $p_T \sim 4.5$ GeV/c



Supported in part by the

U.S. DEPARTMENT OF
ENERGY Office of
Science




清华大学
Tsinghua University

清华大学
Tsinghua University

Ω Baryon p_T Spectra


- Precise measurement for 7 centrality bins
- Maximum $p_T \sim 4$ GeV/c
- Fraction of the integral yield covered by the measured data points: $62 \sim 70\%$



dN/dy Yields

- N_{part} (number of participant nucleons) scaled Ω yields increase faster with the increasing N_{part} from large (Au+Au) towards small (O+O) systems.
- May result from higher N_{coll} (number of binary collisions) corresponding to the same N_{part} in smaller systems.


Run11 data points N_{part} shifted for clarity
 $p+p$: STAR, Phys. Rev. C 75 (2007) 064901
 $Cu+Cu$: STAR, Phys. Rev. Lett. 108 (2012) 072301



Ω/ϕ Ratio

- Significant Ω over ϕ enhancement observed in central isobar collisions — compatible with the **existence of QGP in central isobar collisions**.
- Moderate enhancement in central O+O, consistent with 40-60% isobar (similar $N_{part} \sim 30$).

$p+p$ 200 GeV $\Omega + \bar{\Omega}$: X. Zhu, QM2014;
 $p+p$ 200 GeV ϕ : STAR, Phys. Rev. C 79(2009) 064903
Theory: Phys. Rev. C, 2007, 75: 054904.



Summary and Outlook

- Precise measurement of $\phi, \Omega, \bar{\Omega}$ p_T spectra and $\Omega, \bar{\Omega}$ yields in isobar and O+O collisions at 200 GeV are achieved.
- N_{part} scaled Ω yield increases faster with the increasing N_{part} from large (Au+Au) towards small (O+O) system.
- Significant Ω over ϕ enhancement at intermediate p_T is observed in central isobar collisions — compatible with the **existence of QGP in central isobar collisions**.
- Moderate Ω over ϕ enhancement in central O+O is observed, consistent with 40-60% isobar, where N_{part} is similar ~ 30 ; analysis with high-multiplicity triggered O+O events is ongoing.



The STAR
Collaboration