



Measurement of K^0 and K^{*+} in Ru+Ru and Zr+Zr collisions at RHIC


Subhash Singha (subhash@impcas.ac.cn), for the STAR Collaboration
Institute of Modern Physics Chinese Academy of Sciences, Lanzhou



Abstract

The neutral K^0 and charged K^{*+} vector mesons have similar masses and total isospin (I), but their third component of isospin (I_3) has opposite sign, and their quark magnetic moment differ by a factor of five. These properties making them ideal probes for studying Landau level splitting in presence of magnetic (B) field in heavy-ion collisions. We present the reconstruction of K^0 and K^{*+} mesons in RHIC isobar (Ru+Ru and Zr+Zr) collisions. We report on the yield and $\langle p_T \rangle$ of K^0 mesons as well as K^0/K ratio in these collisions as a function of collision centrality. The analysis of $K^{*0,+}$ involving isospin-asymmetric nuclei ($Au+Au$, $Ru+Ru$ and $Zr+Zr$) and isospin-symmetric nuclei ($O+O$), as well as in $p+p$ collisions at 200 GeV is underway. This study can offer insights into Landau levels, isospin violation, and late-stage B-fields in QCD medium.

Introduction:



K. Xu et al, Phys Lett B 809, 135706 (2020)

The energy level for a point-like charged particle under static uniform magnetic field (B)

$$\epsilon_{n,s_z}^2(p_z) = p_z^2 + (2n - 2 \text{sign}(q)s_z + 1)|qB| + m^2$$


q = electric charge
 n = Landau level
 s_z = projection of spin along magnetic field
 p_z = momentum along magnetic field

- Under B -field, $N_{K^{*+}} > N_{K^0}$ is expected from Landau level splitting (isospin violation)
- Neutral and charged vector meson's yield ratio can be used to constraint B -field at freeze-out


Signal reconstruction:

Invariant mass method is used to reconstruct $K^{*0,+}$ signal; combinatorial background is constructed via track-rotation technique. A clear signal is observed on top of a combinatorial background

K^0



K^{*+}




Signal is fitted with a non-relativistic Breit-Wigner function and a second order polynomial for residual background

$$\frac{Y}{2\pi(m-m_0)^2 + (\Gamma_0/2)^2} + AM^2 + BM + C$$

Experimental details:

STAR offers uniform acceptance, full azimuthal coverage and excellent particle identification using Time Projection Chamber (TPC) and Time Of Flight (TOF) detectors




Event selection:

- Vertex: $-35 < V_z < 25$ cm, $V_r < 2$ cm; Centrality from TPC; Rejection of pile-up events


Track selection:

- No. of TPC hits > 15 ,
- $K^0 \rightarrow \pi^+ \pi^-$, $K^{*0} \rightarrow K_S^0 \pi^-$
- K^0 : $p_T > 0.2$ GeV/c, $DCA < 2$ cm, $|\eta| < 1.0$ for K and π
- K^{*+} : STAR helix topology method for K_S^0 selection, $p_T > 0.2$ GeV/c, $DCA < 2$ cm, $|\eta| < 1.0$ for π
- K and π 's are identified using TPC and TOF detectors


Yield, $\langle p_T \rangle$ and particle ratios:



- K^0 dN/dy increase with $\langle N_{part} \rangle$



- K^0 $\langle p_T \rangle$ similar to protons (mass dependence)



- $(K^0/K)_{\text{central}} < (K^0/K)_{\text{peripheral}}$ ($\sim 3.3 \sigma$)

- Evidence of late-stage hadronic re-scattering

Summary and outlook:

- K^0 production \rightarrow studied in isobar collisions ($Ru+Ru$, $Zr+Zr$)
- $K^0/K \rightarrow$ evidence of late-stage hadronic re-scattering

- $K^{*+} \rightarrow$ analysis in $Au+Au$, $Ru+Ru$, $Zr+Zr$, $O+O$ and $p+p$ collisions is underway
- Yield difference between neutral K^0 and charged K^{*+} in heavy-ion collisions can help constrain the B-field at freeze-out

Supported in part by the



This work is supported by grant from the Chinese Academy of Sciences

The STAR Collaboration

