

Beam Energy Dependence of Directed Flow of pions and Kaons in Au+Au Collisions from STAR

Ze Qiu
(for the STAR Collaboration)

1 The first harmonic coefficient of the Fourier expansion of the final-state momentum-
2 space azimuthal distribution of produced particles relative to the event plane is called
3 directed flow (v_1), and describe the collective sideward motion of emitted particles. The
4 v_1 is sensitive to the equation of state of nuclear matter, making it a useful probe for
5 studying the phase transition. Results from RHIC Beam Energy Scan-I (BES-I) program
6 on directed flow show that the proton and net-proton slope parameter $\frac{dv_1}{dy}|_{y=0}$ exhibits a
7 minimum between 11.5 and 19.6 GeV. Additionally, the net-proton $\frac{dv_1}{dy}|_{y=0}$ changes sign
8 twice between 7.7 and 39 GeV.

9 In this poster, we will present measurements of the directed flow of π^+, π^-, K^+, K^- and
10 K_s^0 in Au+Au collisions at $\sqrt{s_{NN}} = 7.7, 9.2, 11.5, 14.6, 17.3$ and 19.6 GeV from the RHIC
11 Beam Energy Scan-II (BES-II) program. The rapidity and centrality dependence of v_1 ,
12 as well as the energy dependence of the v_1 slope, will be discussed. These results will also
13 be compared with predictions from the ultra-relativistic quantum molecular dynamics
14 (UrQMD) model.