



Supported in part by:

U.S. DEPARTMENT OF
ENERGY

Office of
Science



Evidence of $\Lambda\bar{\Lambda}$ spin correlation in p+p collisions at STAR

Jan Vanek, for the STAR Collaboration

Brookhaven National Laboratory

QM 2025, Frankfurt, Germany

7. 4. 2025



STAR Collaboration

NON-PERTURBATIVE QCD: QUARK CONFINEMENT




Low temperature/density

- Confinement of partons in hadrons
- Chiral symmetry broken
- Hadronic matter



High temperature/density

- Deconfined matter of quarks and gluons
- Chiral symmetry (partially) restored
- Early universe, Quark-Gluon Plasma



NON-PERTURBATIVE QCD: QUARK CONFINEMENT


Low temperature/density

- Confinement of partons in hadrons
- Chiral symmetry broken
- Hadronic matter



High temperature/density

- Deconfined matter of quarks and gluons
- Chiral symmetry (partially) restored
- Early universe, Quark-Gluon Plasma



- Hadron-parton transition
 - Temperature, nature of phase transition?
- Chiral symmetry breaking/restoration
 - Same temperature as hadron-parton transition?
- How do hadrons acquire their properties?
 - Parton vs. hadron properties (mass, spin...)

Uncovering how confinement shapes hadron properties is one of the key questions in nuclear and particle physics.

EMERGENT HADRON PROPERTIES



- **Mass**

- Hadron masses far exceed those of their constituent partons
 - Chiral symmetry breaking
 - Dynamical mass generation from **QCD vacuum**



EMERGENT HADRON PROPERTIES


■ Mass

- Hadron masses far exceed those of their constituent partons
 - Chiral symmetry breaking
 - Dynamical mass generation from **QCD vacuum**

■ Spin

- Proton spin crisis
 - Valence quarks account for only $\sim 30\%$ of proton's spin
 - Gluons carry next $\sim 30\%$ and sea quarks the rest
- Λ^0 hyperon polarization puzzle
 - Polarized Λ^0 observed in collisions of unpolarized particles

- Our goal is to experimentally reveal the origin of these emergent hadron properties




VACUUM: MAXIMALLY ENTANGLED QUARK PAIRS



- Quark condensate has up ($u, \bar{u}$), down ($d, \bar{d}$), and strange ($s, \bar{s}$) quarks
- **Quark pairs from the vacuum** are expected to emerge as **maximally entangled spin-triplet states** with aligned spins
 - The only way to preserve the vacuum's quantum numbers $J^{PC} = 0^{++}$

J. R. Ellis, D. Kharzeev, and A. Kotzinian, Z. Phys. C 69, 467 (1996)

- We aim to experimentally track the evolution of these entangled $s\bar{s}$ quark pairs into the hadronic level




NEW EXPERIMENTAL APPROACH

- Λ^0 hyperon pair spin-spin correlations in $p+p$ collisions:


(a)

High energy proton-proton collisions liberate quarks and gluons




(b)

Collisions that excite the vacuum can produce quark-antiquark pairs




(c)

The quarks undergo the hadronization process which turns them into hadrons due to confinement



(d)

Λ and $\bar{\Lambda}$ hadron can be formed in the hadronization process, and each has its hadron spin polarization




- We use the spin-spin correlation of $\Lambda^0 \bar{\Lambda}^0$ hyperon pairs measured in $p+p$ collisions to study the hadronization of the entangled $s\bar{s}$ quark pairs from the QCD vacuum

EXPERIMENTAL METHOD

- Find a Λ^0 hyperon pair (any combination) in one event
 - Decay channel $\Lambda^0 \rightarrow p\pi^+$ and charge conjugate
- Boost proton (anti-proton) from decay of the corresponding Λ^0 ($\bar{\Lambda}^0$) to **rest frame of its mother**
- Measure angle θ^* between the two **boosted protons**
 - The distribution of pair angle is given by:


$$\frac{1}{N} \frac{dN}{d\cos(\theta^*)} = \frac{1}{2} [1 + \alpha_1 \alpha_2 P_{\Lambda_1 \Lambda_2} \cos(\theta^*)]$$
 - α_1 and α_2 are weak decay parameters of Λ^0 or $\bar{\Lambda}^0$ (α_- or α_+)
- A non-zero $P_{\Lambda_1 \Lambda_2}$ would indicate spin correlation between the two Λ^0 ($\bar{\Lambda}^0$) hyperons**
 - No global single Λ^0 hyperon polarization expected at STAR at mid-rapidity



SOLENOIDAL TRACKER AT RHIC (STAR)




- Key subsystems for this analysis:
 - Solenoidal magnet
 - 0.5 T magnetic field **allowing low p_T coverage**
 - Time Projection Chamber (TPC)
 - Measurement of charged particle transverse momentum (p_T)
 - Particle identification (PID) based on energy loss in TPC gas
 - **Full azimuthal coverage for $|\eta| < 1$**



EVENT AND TRACK SELECTION, PID




- Data-set:
 - $p+p$ collisions at $\sqrt{s} = 200$ GeV (2012)
 - Ca. 600M minimum bias events
- Track selection to ensure good track quality within geometrical acceptance
- Particle identification (PID) to obtain pure proton and pion sample
- Decay topology to suppress combinatorial background from tracks originating from close to primary vertex



Λ^0 SIGNAL EXTRACTION



- Signal extraction determined from 2D M_{inv} distributions of unlike-sign (US) πp pairs
- Three components:
 - a) Peak: πp from Λ^0 ($\bar{\Lambda}^0$) decay paired with another πp from Λ^0 ($\bar{\Lambda}^0$) decay
 - b) Ridges: πp from Λ^0 ($\bar{\Lambda}^0$) decay paired with combinatorial background
 - c) Continuum: combinatorial background paired with combinatorial background
- Contributions (b) and (c) are subtracted from (a) and fitted with 2D Gaussian function
 - Signal region is defined as mean $\pm 2\sigma$



ACCEPTANCE EFFECT CORRECTION




- Measured $dN/d \cos(\theta^*)$ distributions have significant slope originating from acceptance effect
 - Predominantly from lower p_T cut on daughter π
- Correction done using mixed-event (ME) hyperon pairs
 - ME definition example:

Same-event $\Lambda \bar{\Lambda}$



Single $\bar{\Lambda}$ from different event




Mixed-event $\Lambda \bar{\Lambda}$


$$\begin{aligned} |p_{T,SE} - p_{T,single}| &< 0.1 \text{ GeV}/c \\ |y_{SE} - y_{single}| &< 0.1 \\ |\phi_{SE} - \phi_{single}| &< 0.1 \end{aligned}$$

- Verified using standalone PYTHIA simulation (shown on the right), as well as full simulation with detector effects

Before correction



After correction



SPIN-SPIN CORRELATION EXTRACTION




- Spin-spin correlation is extracted by fitting $dN/d \cos(\theta^*)$ distributions after ME correction for signal+background (P_{S+B}) and background (P_B) using:

$$dN/d \cos(\theta^*) = A[1 + B \cos(\theta^*)]$$

- A and B are parameters of the fit
- A is normalization, $B = \alpha_1 \alpha_2 P_{\Lambda_1 \Lambda_2}$
- Signal (P_S) is calculated according to:


$$P_{S+B} = f_S P_S + (1 - f_S) P_B$$
 - f_S is signal fraction
- Signal extracted for two relative pair kinematics:
 - Short-range:** $|\Delta y| < 0.5$ and $|\Delta \phi| < \pi/3$
 - Long-range:** $0.5 < |\Delta y| < 2.0$ or $\pi/3 < |\Delta \phi| < \pi$



Λ SPIN-SPIN CORRELATIONS



- Spin-spin correlation for short-range (a) and long-range (b) hyperon pairs
 - Compared to null measurement of K_S^0 and PYTHIA calculation
- Short-range $\Lambda\bar{\Lambda}$ pairs show non-zero spin-spin correlation
 - $P_{\Lambda_1\Lambda_2} = 0.181 \pm 0.035_{stat} \pm 0.022_{sys}$
 - Significance 4.4 standard deviation
- All other pairs are consistent with zero



Λ SPIN-SPIN CORRELATIONS




- Spin-spin correlation for short-range (a) and long-range (b) hyperon pairs

- Compared to null measurement of K_S^0 and PYTHIA calculation

- Short-range $\Lambda\bar{\Lambda}$ pairs show non-zero spin-spin correlation

- $P_{\Lambda_1\Lambda_2} = 0.181 \pm 0.035_{stat} \pm 0.022_{sys}$
- Significance 4.4 standard deviation

- All other pairs are consistent with zero



MAXIMUM EXPECTED SPIN-SPIN CORRELATION



- **Pure spin triplet $s\bar{s}$ pairs** from QCD vacuum: $P_{s\bar{s}} = 1/3$ N. A. Tornqvist, Phys. Lett. A542117, 1 (1986)
 - What happens when such pair hadronizes into $\Lambda^0\bar{\Lambda}^0$ pair?

- **Expected maximum for $\Lambda^0\bar{\Lambda}^0$ pairs** in our dataset based on models and feed-down from decay of heavier hyperons:

Single Λ^0 ($\bar{\Lambda}^0$) polarizations depending on its mother particle from SU(6) and BJ models

Λ 's parent	SU(6)	BJ model
Primary	1	0.63
Σ^0	1/9	0.15
Ξ^0	0.6	-0.37
Ξ^-	0.6	-0.37
Σ^*	5/9	N/A

- $P_{\Lambda_1\Lambda_2,\text{SU}(6)} = 0.096 \pm 0.004$
- $P_{\Lambda_1\Lambda_2,\text{BJ}} = 0.015 \pm 0.002$


- Model prediction has two components:
 - Single Λ^0 ($\bar{\Lambda}^0$) polarization depending on its mother particle from two models:
 - Non-relativistic SU(6) quark model and Burkardt-Jaffe (BJ) model
 - Feed-down mixture for $\Lambda^0\bar{\Lambda}^0$ pairs from PYYHIA 8 + Geant simulation

M. Burkardt and R. L. Jaffe, Phys. Rev. Lett. 70, 2537 (1993)
J. Ellis, *et al.*, Eur. Phys. J. C 52, 283–294 (2007)

$\Lambda\bar{\Lambda}$ SPIN-SPIN CORRELATIONS




- Spin-spin correlation of $\Lambda\bar{\Lambda}$ pairs as a function of $\Delta R = \sqrt{\Delta y^2 + \Delta\phi^2}$
 - Compared to SU(6) and BJ models
- Short range $\Lambda\bar{\Lambda}$ pairs appear to be in triplet state**
 - First direct probe of QCD vacuum?
- Possible observation of quantum decoherence**
 - Spin-spin correlation seem to “turn off” with larger pair separation



PHYSICS IMPLICATIONS




- We observed the **first evidence of a positive spin-spin correlation** of short-range $\Lambda\bar{\Lambda}$ pairs from p+p collisions at 200 GeV.
- **This may represent a new experimental approach to probing the quark condensate**
 - Our results align with short-range $\Lambda\bar{\Lambda}$ pairs being in a spin-triplet state, as expected for $\Lambda\bar{\Lambda}$ pairs from the QCD vacuum
- **The $s\bar{s}$ pair's spin-spin correlation** appears to largely **survive hadronization**
 - First time, we seem to track the spin degree of freedom through QCD quark-to-hadron transition
 - A key insight into hadronization process of (s) quarks



PHYSICS IMPLICATIONS CONT.



- **Result support a scenario where s quarks carry most of the hyperon's spin**
 - Results offer valuable insight into how partons contribute to the Λ hyperon's spin
- **Possible experimental approach to observe quantum decoherence**
 - Spin-spin correlation weakens with increasing pair separation
 - Can be extended to p +Au or Au+Au to study influence of cold nuclear matter effects and the QGP
- **Provides new insight into the Λ hyperon polarization puzzle and spin transfer**
 - New experimental constraints on both initial- and final-state models



FUTURE PROSPECTS




- Extension of kinematic coverage beyond $|y| < 1$
 - Access to region with non-zero single Λ^0 hyperon polarization
 - **STAR forward upgrade**
- High p_T Λ^0 in jets – study of $g \rightarrow s\bar{s}$
 - Spin configuration of $s\bar{s}$ given by gluons
- Collision energy dependence
 - From RHIC to LHC
- Other collision systems
 - $p+A$ – cold nuclear matter effects
 - $A+A$ – Quark Gluon Plasma, chiral symmetry restoration

STAR forward upgrade

$2.5 < |\eta| < 4$

Forward Tracking and Calorimeter Systems

Taking data since 2022



THANK YOU FOR ATTENTION



BACKUP

EVENT AND TRACK SELECTION, PID



- Data-set:
 - p+p collisions at $\sqrt{s} = 200$ GeV (2012)
 - Ca. 600M minimum bias events
- Events with primary vertex close to center of STAR detector selected
- Track selection to ensure good track quality within geometrical acceptance
- Particle identification to obtain pure proton and pion sample
- Decay topology to suppress combinatorial background from tracks originating from close to primary vertex



Event selection	$ V_z < 60$ cm
	$p_T > 150$ MeV/c
Track selection	$ \eta < 1$
	nHitsFit > 20
	nHitsFit/nHitsMax > 0.52
Particle identification	$ n\sigma_\pi < 3$
	$ n\sigma_p < 2$
Decay topology Λ^0	DCA _{π-PV} > 0.3 cm
	DCA _{p-PV} > 0.1 cm
	DCA _{Λ-PV} < 1.0 cm
	DCA _{pair} < 1.0 cm
	$2 \text{ cm} < L_{\text{dec}} < 25 \text{ cm}$
	$\cos(\theta) > 0.996$

EXPERIMENTAL METHOD





- Find a Λ^0 hyperon pair (any combination) in one event
 - Decay channel $\Lambda^0 \rightarrow p\pi^+$ and charge conjugate
 - $p_{\Lambda^0,lab} = p_{p,lab} + p_{\pi^-,lab}$
- Boost (anti-)proton from decay of the corresponding Λ^0 ($\bar{\Lambda}$) to **rest frame of its mother**
 - Proton momenta in mother rest frame: $p_{p,CMS_\Lambda}, p_{\bar{p},CMS_{\bar{\Lambda}}}$
- Measure angle θ^* between the two **boosted protons**
 - The distribution of pair angle is given by:

$$\frac{1}{N} \frac{dN}{d\cos(\theta^*)} = \frac{1}{2} [1 + \alpha_1 \alpha_2 P_{\Lambda_1 \Lambda_2} \cos(\theta^*)]$$
 - α_1 and α_2 are α_+ or α_- , depending on Λ^0 hyperon pair
- A non-zero $P_{\Lambda_1 \Lambda_2}$ would indicate spin correlation between the two Λ^0 ($\bar{\Lambda}^0$) hyperons**



EXPERIMENTAL METHOD

- Find a Λ^0 hyperon pair (any combination) in one event
 - Decay channel $\Lambda^0 \rightarrow p\pi^+$ and charge conjugate
 - $p_{\Lambda^0,lab} = p_{p,lab} + p_{\pi^-,lab}$
- Boost to the center of mass of the Λ^0 pair (the mother's CMS)
 - This experimental method is sensitive to selection criteria and detector acceptance
- Major source is lower cut on p_T of π
 - Preferential selection of Λ decays, where π is emitted along Λ momentum
 - This is called the **acceptance effect** and is corrected in this analysis
- A non-zero $P_{\Lambda_1\Lambda_2}$ would indicate spin correlation between the two Λ^0 hyperons



FEED-DOWN ESTIMATION



- Feed-down estimation for $\Lambda^0 \bar{\Lambda}^0$ in $p+p$ collisions at $\sqrt{s} = 200$ GeV
 - PYTHIA 8 + Geant simulation

