

Precision measurements of light hypernuclei lifetime and branching ratio fraction R_3 by the STAR experiment

Yuanjing Ji (LBNL), Xiujun Li (USTC)
for the STAR Collaboration

2021-11-25

1 Hypernuclei are bound states of nucleons and hyperons, and thus naturally correlated hyperon-
2 baryon systems. Hypernuclei are regarded a unique laboratory to study the hyperon-nucleon (Y - N)
3 interaction. The Y - N interaction is an important ingredient, not only in the equation-of-state (EoS) of
4 astrophysical objects such as neutron stars, but also in the description of the hadronic phase of a heavy-
5 ion collision. The strength of the Y - N interaction can be investigated by measuring the properties of
6 hypernuclei. Precise determination of hypernuclei structure parameters, such as Λ separation energy
7 B_Λ , lifetime, and branching ratios, may also shed light on the role that two-body Y - N and three-body
8 Y - N - N interactions play in the density regime of neutron stars.

9 In this talk, we report precision measurements of the lifetime of ${}^3_\Lambda\text{H}$, ${}^4_\Lambda\text{H}$ and ${}^4_\Lambda\text{He}$ obtained from
10 Au+Au collisions collected by STAR during the Beam Energy Scan Phase-II program. Hypernuclei
11 are reconstructed via charged pion decay channels including both two-body and three-body decay
12 modes. We also present the relative branching ratio R_3 of ${}^3_\Lambda\text{H}$ and ${}^4_\Lambda\text{H}$, where R_3 is the fraction of
13 the two-body decay rate out of the sum of two-body and three-body decay rates. The results will be
14 compared with model calculations and physics implications will be discussed.