

Elliptic flow of light nuclei in Au+Au collisions at $\sqrt{s_{NN}} = 14.6, 19.6, 27, \text{ and } 54.4 \text{ GeV}$

Rishabh Sharma (for the STAR Collaboration)

Indian Institute of Science Education and Research (IISER) Tirupati

Abstract

1 High energy heavy-ion collisions provide an opportunity to study the production mechanism
2 of light (anti)nuclei. There are two main possible models to explain the production mechanism
3 - the thermal model and the coalescence model. The thermal model has been quite successful
4 in explaining the yields of light (anti)nuclei produced in heavy-ion collisions. Thermal model
5 suggests that the light (anti)nuclei are produced from a thermal source, where they are in equi-
6 librium with other species present in the fireball. However, due to the small binding energies, the
7 produced (anti)nuclei are not likely to survive in the high temperature conditions of the fireball.
8 The coalescence model, on the other hand, predicts that light (anti)nuclei can be formed by the
9 final state coalescence of nucleons. Studying the azimuthal anisotropy of light (anti)nuclei and
10 comparing them with that of nucleons, therefore, will give insights into the production mechanism
11 of light (anti)nuclei.

12 In this poster, we will present the energy and centrality dependence of elliptic flow (v_2) of d ,
13 t , and ${}^3\text{He}$ and their antiparticles in Au+Au collisions at $\sqrt{s_{NN}} = 14.6, 19.6, 27, \text{ and } 54.4 \text{ GeV}$.
14 Mass number and number of constituent quarks (NCQ) scalings will also be shown. The $v_2(p_T)$
15 of light (anti)nuclei will be compared with the coalescence and hydrodynamic models.