

Investigating Entanglement Enabled Spin Interference in photonuclear $\rho^0 \rightarrow \pi^+\pi^-$ and $\gamma\gamma \rightarrow \pi^+\pi^-$ in Au+Au collisions at STAR

Sam Corey (for the STAR Collaboration)

Abstract

1 In ultraperipheral collisions, the invariant mass spectrum of $\pi^+\pi^-$ pairs is very complex
2 due to the numerous production channels and intermediate states. The quantum ambiguity
3 between production channels, referred to as the Entanglement Enabled Spin Interference
4 (EESI) effect, leads to angular anisotropy in the final state. The most dominant contribution
5 to the invariant mass spectrum of $\pi^+\pi^-$ is $\gamma A \rightarrow \rho^0(770) \rightarrow \pi^+\pi^-$, but other photonuclear
6 (γA) and light-by-light ($\gamma\gamma$) channels also must be considered. EESI between the $\gamma\gamma$ and
7 γA channels is expected to produce $A_{1\Delta\phi}$ and $A_{3\Delta\phi}$ signals. This new window into hadronic
8 light-by-light production may provide new theoretical constraints on the anomalous magnetic
9 moment of the muon, where the hadronic light-by-light contribution is one of the largest
10 uncertainties.

11 In this talk, the first measurement of EESI between photonuclear and light-by-light pro-
12 duction of $\pi^+\pi^-$ pairs, including the strong EESI signal associated with the $f_2(1270)$ res-
13 onance, will be presented. The EESI observables are then used to isolate $\gamma\gamma \rightarrow \pi^+\pi^-$ in
14 ultraperipheral $Au + Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV.