



XI International Conference on New Frontiers in Physics



Search for the Chiral Magnetic Effect by the STAR Experiment

Gang Wang (UCLA)


for the STAR Collaboration

Supported in part by




U.S. DEPARTMENT OF
ENERGY

Office of
Science




Chiral Magnetic Effect: $J \propto \mu_5 B$


1 Chiral symmetry restoration (massless quarks)



2 Chirality imbalance (finite μ_5 , Local Parity Violation)




3 Strong magnetic field ($B \sim 10^{18}$ Gauss)



Experimental manifestation: charge separation across the reaction plane

4 Chiral Magnetic Effect ($J \parallel B$)



Observables in search of CME

$$\frac{dN_\alpha}{d\phi^*} \approx \frac{N_\alpha}{2\pi} [1 + 2v_{1,\alpha} \cos(\phi^*) + 2v_{2,\alpha} \cos(2\phi^*) + 2v_{3,\alpha} \cos(3\phi^*) + \dots + 2a_{1,\alpha} \sin(\phi^*) + \dots] \quad \phi^* = \phi - \Psi_{RP}$$

CME-sensitive observables on the market:

- γ correlator

S.A. Voloshin, Phys. Rev. C,70, 057901 (2004)

- R correlator

N. N. Ajitanand *et al.*, Phys. Rev. C83, 011901(R) (2011)

- Signed balance functions


A. H. Tang, Chin. Phys. C,44, No.5 054101 (2020)

Here we focus on

$$\gamma_{112} \equiv \langle \cos(\phi_\alpha + \phi_\beta - 2\Psi_2) \rangle$$




And the CME signal should cause

$$\Delta\gamma_{112} \equiv \gamma_{112}^{OS} - \gamma_{112}^{SS} > 0$$



S. Choudhury *et al.* (STAR), Chinese Phys. C 46 (2022) 014101. AVFD simulations show that these methods have **similar sensitivities** to the CME signal and to the background.

γ_{112} measurements at RHIC/LHC



In various collision systems and at different beam energies, positively **finite** $\Delta\gamma_{112}$ meets the CME expectation, **but** could contain contributions from:



- Flow-related background $\propto v_2$ (elliptic flow)
- Nonflow-related background (di-jets)

κ_{112} measurements at RHIC/LHC

$$\kappa_{112} \equiv \Delta\gamma_{112}/(v_2\Delta\delta)$$

$$\delta \equiv \langle \cos(\phi_\alpha - \phi_\beta) \rangle$$


Normalized quantity facilitates comparison between data and model calculations (AMPT).



Compared with a pure-background model, the **CME signal** seems to **disappear at 7.7 GeV and 2.76 TeV**.

- very low beam energies: **no chiral symmetry restoration?**
- very high energies: **no duration of the magnetic field?**

Isobar collisions: prospect




Compare the two isobaric systems:

- CME: $B\text{-field}^2$ is $\sim 15\%$ larger in Ru+Ru
- Flow-related BKG: utilize $\Delta\gamma_{112}/v_2$
- Nonflow-related BKG: almost same



Isobar collisions provide best possible control of signal and background.

2.5 B events per species:

- uncertainty of **0.4%** in the $\Delta\gamma/v_2$ ratio.
- if $f_{\text{CME}} > 14\%$, $\Delta\gamma_{112}/v_2$ difference $> 2\%$, yielding a **5σ** significance.
- f_{CME} is the unknown CME fraction in $\Delta\gamma_{112}$.



Isobar program: data collection in 2018



Successful data taking of isobar collisions at RHIC/STAR

PHYSICAL REVIEW C
covering nuclear physics



Highlights Recent Accepted Collections Authors Referees Search Press About

Search for the chiral magnetic effect with isobar collisions at $\sqrt{s_{NN}} = 200$ GeV by the STAR Collaboration at the BNL Relativistic Heavy Ion Collider

M. S. Abdallah *et al.* (STAR Collaboration)
Phys. Rev. C **105**, 014901 – Published 3 January 2022

First publication after 3 years and many people's efforts...


Isobar blind analysis



Blinding committee decides the procedure.

STAR, Nucl. Sci. Tech. 32 (2021) 48

Five independent groups run each other's frozen code.



No access to species-specific information until last step.
Everything documented (**not written** → **not allowed**)

Case for CME & interpretation must be pre-defined.

Centrality definition

Blind analysis: compare observables at **matching centrality** between two isobar systems.

STAR, Phys. Rev. C 105 (2022) 14901



Case-1 [83]			
Nucleus	R (fm)	a (fm)	β_2
$^{96}_{44}\text{Ru}$	5.085	0.46	0.158
$^{96}_{40}\text{Zr}$	5.02	0.46	0.08

Deng *et al.*, PRC 94, 041901 (2016)

Case-2 [83]			
Nucleus	R (fm)	a (fm)	β_2
$^{96}_{44}\text{Ru}$	5.085	0.46	0.053
$^{96}_{40}\text{Zr}$	5.02	0.46	0.217


Xu *et al.*, PRL. 121, 022301 (2018)

Case-3 [113]			
Nucleus	R (fm)	a (fm)	β_2
$^{96}_{44}\text{Ru}$	5.067	0.500	0
$^{96}_{40}\text{Zr}$	4.965	0.556	0



MC-Glauber model fits the uncorrected multiplicity distribution. Woods-Saxon parameters with thicker neutron skin in Zr (no deformation) gives the best fit of the multiplicity distributions.

Multiplicity mismatch



Case-3 (thicker neutron skin in Zr and zero β_2) gives the **best fit** of the multiplicity distributions.


However, multiplicity (efficiency uncorrected) is larger in Ru+Ru than in Zr+Zr in such a matching centrality.

This can affect background (and signal) difference between the two isobaric systems.

Case-1 and **Case-2** give (almost) the **same multiplicity** in Ru+Ru and Zr+Zr, but they don't describe the multiplicity distribution so well.

In the end, the blind analysis sticks to **Case-3**.


v_2 and $\Delta\delta$




STAR has multiple sets of results with different kinematic cuts. I will use the set with smallest statistical errors as a demonstration.

Both v_2 and $\Delta\delta$ contribute to the background, and their **ratios** of Ru+Ru to Zr+Zr are **not exactly unity**.

At matching centrality, the below-unity $\Delta\delta$ ratio could even fake a CME signal.




$\Delta\gamma_{112}$ and κ_{112}




Both $\Delta\gamma_{112}$ and κ_{112} ratios of Ru+Ru to Zr+Zr go **significantly below unity!**

How can we understand this?




Matching centrality or matching multiplicity?



Qualitative change at matching multiplicity: κ_{112} ratios are more consistent with unity.

κ_{112} ratio $\approx 1 + 15\% f_{\text{CME}}$



$$\kappa_{112} \equiv \frac{\Delta\gamma_{112}}{v_2 \cdot \Delta\delta}$$


Pre-defined CME signature,

$$\frac{\kappa_{112}^{\text{Ru+Ru}}}{\kappa_{112}^{\text{Zr+Zr}}} > 1$$

is NOT seen.

Upper limit (95% CL):



1.0066 for κ_{112} ratio;
 ~4% for f_{CME} .



Small interpolation before taking ratios

Post-blinding

$\Delta\gamma_{112}$ results are **consistent with preliminary background estimate** within current uncertainty.




Why is f_{CME} so small?

AVFD simulation:


f_{CME} is smaller in isobar than Au+Au, especially when using the participant plane.
smaller system \rightarrow larger fluctuation \rightarrow larger BKG
& smaller CME signal \rightarrow lower f_{CME}

The bright side





H-J. Xu, et al, CPC 42 (2018) 084103;

S. A. Voloshin, Phys. Rev. C 98 (2018) 054911



The difference between different event plane types indicates a **finite f_{CME} in Au+Au at 200 GeV**. More data to come!




Event-shape engineering

- suppresses v_2 -related background
- enhances f_{CME} (**finite at 27 GeV?**)

Backup slides

Matching centrality or matching multiplicity?



$$\rho(r, \theta, \phi) \propto \frac{1}{1 + e^{[r - R_0(1 + \beta_2 Y_2^0(\theta, \phi) + \beta_3 Y_3^0(\theta, \phi))]/a_0}}$$

J. Jia, G. Wang, C. Zhang, arXiv:2203.12654

The difference between matching centrality and matching multiplicity comes from a_0 , surface diffuseness.

Isobar: charge separation measured with R_{ψ_2}

$$R_{\psi_2}(\Delta S) = C_{\psi_2}(\Delta S) / C_{\psi_2}^{\perp}(\Delta S)$$

$$C_{\psi_2} = \frac{N_{\text{real}}(\Delta S)}{N_{\text{shuffled}}(\Delta S)}$$


$$\Delta S = \left\{ \frac{\sum_{i=1}^{n^+} w_i^+ \sin(\phi_i - \psi_2)}{\sum_{i=1}^{n^+} w_i^+} - \frac{\sum_{i=1}^{n^-} w_i^- \sin(\phi_i - \psi_2)}{\sum_{i=1}^{n^-} w_i^-} \right\}$$

σ_{ψ_2} is the Gaussian width of the respective $R(\Delta S)$

Measurement of the in-plane and out-of-plane distributions of the dipole separation event by event.

STAR Isobar blind analysis, $\sqrt{s_{\text{NN}}} = 200$ GeV

STAR, Phys. Rev. C 105 (2022) 14901



Predefined CME signature:

$$1/\sigma_{\psi_2}^{\text{Ru+Ru}} > 1/\sigma_{\psi_2}^{\text{Zr+Zr}}$$

No significant difference is observed between the two isobaric systems