



Initial Stages 2023


Copenhagen, June 19-23

Measurements of jet substructure in $p+p$
and jet-event activity correlations in $p+Au$
collisions at $\sqrt{s_{NN}} = 200 \text{ GeV}$ at STAR

David Stewart for the STAR Collaboration




WAYNE STATE
UNIVERSITY





supported in part by
U.S. DEPARTMENT OF
ENERGY

Office of
Science



Jets: probe initial stages





This presentation

$p + p$


Collisions:

- Measure hadronization scale
- Measure gluon and quark jet fractions
- Measure parton showers, correlate hard and soft processes
- Compare/benchmark measurements to MC


$p + Au$

Collisions:

- Report unexpected anticorrelation in hard and soft particle production across wide rapidity gap
- Probe if anticorrelation results from jet quenching



Experimental data: particles in STAR detector



Subsystems (full azimuth coverage):

Barrel ElectroMagnetic Calorimeter (BEMC):

- EM interacting particles
- $|\eta| \leq 1$

Time Projection Chamber (TPC):


- Charged particles
- $|\eta| \leq 1$

Beam Beam Counter (BBC):

- Charged particles
- $|\eta| \in [3.4,5]$

Particles → Jets


- Charged jets: TPC particles
- Full jets: TPC + BEMC particles
- Jet finding/clustering: anti- k_T algorithm



pp



Measure QCD hadronization scale?

- “Energy-Energy Correlators” (EEC, [Lee, Meçaj, and Moult, arXiv 2205.03414](#)) probe hadronization scale ([Andres, et al., arXiv 2209.11236](#))
- Normalized EEC $\equiv \frac{1}{\sum_{\text{Jets}} \sum_{i \neq j} \frac{E_i E_j}{p_{T,\text{jet}}^2}} \frac{d\left(\sum_{\text{Jets}} \sum_{i \neq j} \frac{E_i E_j}{p_{T,\text{jet}}^2}\right)}{d(\Delta R)}$
- EEC (ΔR) corresponds to
 - at small ΔR : free hadrons
 - at large ΔR : perturbative quark & gluon interactions
 - at ΔR between: $\Delta R_{\text{Turnover}} \propto \frac{\Lambda_{\text{QCD}}}{p_{T,\text{Jet}}}$ ([Komiske et al, 2023, PRL 130, 051901](#))





→ $\Lambda_{\text{QCD}} \propto \Delta R_{\text{Turnover}} p_{T,\text{Jet}}$ at all scales

First EEC measurement at RHIC



- $p_{T,\text{Low}}^{\text{Jet}} \Delta R_{\text{Turnover}} = 10^{-0.75} \left(15 \frac{\text{GeV}}{c} \right) = 2.7 \frac{\text{GeV}}{c}$
 \rightarrow scale appears universal between RHIC and LHC (CMS $\approx 2.5 \frac{\text{GeV}}{c}$)
- $\Delta R_{\text{Turnover}}$ depends on quark/gluon fraction







pp

Measure initiating quark/gluon fractions?

Probe with jet charge: $Q_{\kappa}^{\text{jet}} \equiv \sum_{j \in \text{jet}} \left(\frac{p_T^j}{p_{T,\text{jet}}} \right)^{\kappa} Q_j$




- Goal: template fit PYTHIA (u, d, g, other) Q^{jet} to data
→ measure proper (u, d, g, other) fraction of initiators
- Q^{jet} in PYTHIA8 increases and is more quark dominated with increasing p_T^{jet}




pp

Measure initiating quark/gluon fractions?

Probe with jet charge: $Q_\kappa^{\text{jet}} \equiv \sum_{j \in \text{jet}} \left(\frac{p_T^j}{p_{T,\text{jet}}} \right)^\kappa Q_j$





- Both PYTHIAs perform well relative to data
 - Hint of preference for PYTHIA-6 Perugia tune
 - Data mean shifts to the right -> more up quark dominated



p
p

Jet radial constituent p_T profile



$$\rho(\Delta R) \equiv \lim_{\delta r \rightarrow 0} \left\langle \frac{1}{\delta r} \frac{\sum_{|\Delta R| < \delta r} p_{T,i}}{p_{T,\text{jet}}} \right\rangle$$

PYTHIA-8 overestimates broader (gluon-like?) jets relative to data

- Provides vacuum-like baseline for Au+Au collisions to measure hot-dense QCD effects

*Plot in backup shows same result for jet girth



Probe parton shower more differentially

Represent jet as a parton shower

- Contains both soft- and hard-scale physics
- SoftDrop criterion selects “first hard branching”:

$$\frac{\min(p_{T,1}, p_{T,2})}{p_{T,1} + p_{T,2}} > 0.1 \quad (0.1 \text{ from generalized case } z_{\text{cut}} \left(\frac{\Delta R_{12}}{R_{\text{jet}}} \right)^\beta \text{ with } z_{\text{cut}} = 0.1 \text{ & } \beta = 0)$$

- ML software (**MULTIFOLD**) corrects unbinned observables for detector effects (**MULTIFOLD**, applied for first time at RHIC)
→ Probe correlations between observables and hard and soft scales






Correlation: soft radiation to hard splitting angle

$p\bar{p}$


Ratio of jet mass removed (groomed): $\frac{\Delta M}{M} = \frac{M - M_g}{M}$

Hard splitting angle: $R_g = \sqrt{(\phi_1 - \phi_2)^2 + (\eta_1 - \eta_2)^2}$



$\Delta M / M$ distribution is anti-correlated with groomed jet radius R_g
→ Consistent with angular ordering the parton shower

- Large groomed jet radius → small $\Delta M / M$, little/no soft wide-angle radiation in shower
- MC models fit trends in data



Jets in p+A collisions

Benchmark cold nuclear matter effects (Cronin, nuclear
calculate R_{AA} in Au+Au collisions





→ Quantify jet quenching in Au+Au via R_{AA} : $\frac{\text{geom. scaling} \times \text{jet yield from pp}}{\text{jet yield from } p\text{-A}}$

Classify events by “Event Activity” (EA) – soft particle production

In p+A, no jet quenching (i.e. $R_{AA} \approx 1$) anticipated

Results: (p/d)+A

Inclusive
measurements @
all EA: $R \approx 1 \rightarrow$
don't see quenching





John Laijola for PHENIX, Moriond 2021 (link).
Supersedes previously published result (PRL 116, 122301, erratum in preparation).




Surprises: R_{AA} in EA selected events

- Suppressed for high-EA (“central”) central events
- Enhanced for low-EA (“peripheral”) events




John Lajoia for PHENIX, Moriond 2021 ([link](#)).
Supersedes previously published result
(PRL 116, 122301, erratum in preparation)

- Why is there a jet spectra to EA correlation:
Jet quenching? Centrality selection? Phase space configuration? ...





- EA \equiv sum of signal in BBC inner tiles
- At STAR kinematics, there **is not** enough phase space for an event to have
 - (1) trigger in BEMC
 - (2) jet in TPC, and
 - (3) a dijet partner in BBC
- No “auto-correlation” from directly measuring high EA from dijet partners hitting BBC
- Per-trigger normalization doesn’t use geometric scaling used in R_{AA}

EA distribution anticorrelated with $p_{T,\text{lead}}^{\text{reco}}$




- **Anti-correlation:** EA distribution is
 - softest in events with a **high- p_T jet**
 - harder in events with a **mid- p_T jet**
 - hardest in events with a **low- p_T jet**
- i.e.: significant drop in mean EA with increasing $p_{T,\text{lead}}^{\text{reco}}$


Jet spectra per triggered event suppressed at high-EA




- **Anti-correlation**
- Both trigger- and recoil-side jets suppressed in high EA relative to low EA
- Qualitatively different from quenching in QGP in A+A collisions:
In A+A collisions, away-side jets are preferentially more quenched due to trigger surface bias




Test for $p + Au$ jet quenching via jet mass



Jet mass distribution not EA dependent
→ null result for jet quenching




Test for $p + \text{Au}$ jet quenching via jet mass





M distribution in (high-EA) $p + \text{Au}$ consistent with $p + p$

Test for p + Au jet quenching via dijet p_T balance



Require:


- p_T cut on leading (highest p_T) & subleading (2nd highest p_T jet)
- Require azimuthally recoiling:
 $|\phi_{\text{lead}} - \phi_{\text{sub}}| > (\pi - R)$



Dijet p_T balance distribution not EA dependent
→ null result for jet quenching

Jets are uncorrected for detector effects


Events corrected for luminosity and underlying event density → detector and event effects cancel in the ratio




Measurement Summary

Anti-correlation, over large rapidity gap,
between hard and soft particle production


- No jet quenching within kinematics measured
- Rapidity gap shows correlation results from earliest collision stages and/or pre-collision dynamics





p_p Measurement Summary

- Energy-Energy Correlator
 - measured QCD hadronization scale
 - shows universal scale at LHC and RHIC energies
- Jet charge and radial density profiles
 - sensitive to jet initiating partons (gluon vs quark and quark flavor)
 - PYTHIA simulations generally validated
 - future looking: measure distribution of gluon, quark, and quark-flavor of jets
- Jet substructure groomed by SoftDrop
 - groomed mass ratio vs hard splitting angle: consistent with angular ordering
 - mass shared fraction in hard splitting: parton virtuality decreases faster in more collinear splittings



Backup Slides

MultFold Variables


- Jet measurements need to be corrected for detector effects for comparison with theory/model
- Unfolding methods:
 - Iterative Bayesian unfolding ([D'Agostini. arXiv:1010.0632\(2010\)](#))
 - **MultiFold** ([Andreassen et al. PRL 124, 182001 \(2020\)](#))
 - Machine learning driven
 - Unbinned
 - **Simultaneously** unfolds multiple observables → **Correlation** information is retained!
- First application of MultiFold on RHIC data!

- Jet observables
 - p_T : transverse momentum
 - $Q^\kappa = \frac{1}{(p_{T\text{jet}})^\kappa} \sum_{i \in \text{jet}} q_i \cdot (p_{Ti})^\kappa \rightarrow$ Choose K=2
 - $M = |\sum_{i \in \text{jet}} p_i| = \sqrt{E^2 - |\vec{p}|^2}$
 - 4-momentum of the constituent i
 - R_g : groomed jet radius
 - z_g : shared momentum fraction
 - M_g : groomed jet mass


All 6 observables are simultaneously unfolded in an unbinned way!

- Uncertainties due to prior choice accounted for through 6D reweighting based on PYTHIA8 or HERWIG

Multifold vs RooUnfold



Collinear drop groomed jet mass




PYTHIA6 Perugia 2012 STAR tune: Skands. PRD 82, 074018 (2010)
 PYTHIA8 Detroit tune: Aguilar et al. PRD 105, 016011(2022)
 HERWIG7: Bellm, et al. PRC 76, 1-8 (2016)

Measurement excludes jets with $\Delta M = 0$ (45.5% of jets in this jet p_T range)


- First collinear drop groomed jet measurement, sensitive to soft radiation within jets
- MC predictions qualitatively consistent with data; some tension from HERWIG in small ΔM region
- MultiFold allows us to correlate (combinations of) unfolded quantities

Soft radiation vs hard splitting momentum imbalance





R_g and z_g are correlated, $\Delta M/M$ affects R_g
 \rightarrow correlation between $\Delta M/M$ and z_g

- The more mass that is groomed away relative to the original mass, the flatter the z_g distribution is
 - Demonstrates that **early** soft wide angle radiation constrains the momentum imbalance of & the amount of npQCD contributions to **later** splittings
- MC models describe the trend of data



Correlate: groomed mass fraction “ μ ” vs $p_{\text{T},\text{jet}}$ vs R_g

$$\mu = \frac{\max(m_1, m_2)}{m_2 + m_1}$$





μ smaller at smaller R_g

→ indicates faster reduction of virtuality in the jet shower

Correlate splitting scale to hard splitting angle

Splitting scale: $k_T = z_g p_{T,\text{jet}} \sin(R_g)$






pp


Measure initiating quark/gluon fractions?

Measure jet girth




PYTHIA-8 overestimates broader (gluon-like?) jets relative to data


- Provides vacuum-like baseline for Au+Au collisions to measure hot nuclear effects



Small system jet modification score card



PHENIX measurement: $\left\langle R_{dAu, EXP}^{\pi^0} \right\rangle$




PHENIX 2023, arXiv: <https://arxiv.org/pdf/2303.12899.pdf>


STAR spectra modification *not* due to dijets hitting BBC

 Outer BBC $\eta \in (-2, -3.4)$

 Inner BBC $\eta \in (-3.4, -5)$



Shared momentum fraction vs mass pruned



STAR EEC for three different jet p_T selections

