

Status Report on the Analyses of Proton-Number Cumulants in the STAR Fixed-Target Program

*Meeting of the APS Division of Nuclear Physics
and the Physical Society of Japan*

Zachary Sweger
University of California, Davis
For the STAR Collaboration

11/30/23

Waikoloa Village, Hawaii

UC DAVIS
UNIVERSITY OF CALIFORNIA



Supported in part by




QCD Phase Diagram

- Quarks and gluons experience confinement at low temperatures and densities.
- At high temperatures and densities, there is a deconfined phase, a quark-gluon plasma.

Beam Energy Scan (BES)

- BES program at the Relativistic Heavy-Ion Collider scans phase space of QCD matter by colliding gold ions at varying energies
- Seeking to map onset of deconfinement, and the predicted QCD critical point




QCD Phase Diagram

- Quarks and gluons experience confinement at low temperatures and densities.
- At high temperatures and densities, there is a deconfined phase, a quark-gluon plasma.

Beam Energy Scan (BES)

- BES program at the Relativistic Heavy-Ion Collider scans phase space of QCD matter by colliding gold ions at varying energies
- Seeking to map onset of deconfinement, and the predicted QCD critical point



Cumulants of a distribution are defined as

$$C_1 = \langle N \rangle \equiv \mu \text{ [mean]}$$

$$C_2 = \langle (N - \mu)^2 \rangle \equiv \sigma^2 \text{ [variance]}$$


$$C_3 = \langle (N - \mu)^3 \rangle$$


$$C_4 = \langle (N - \mu)^4 \rangle - 3\langle (N - \mu)^2 \rangle^2$$

The standardized moments of a distribution are

$$S\sigma = C_3/C_2 \text{ [skewness]} \quad \text{measure of distribution's asymmetry}$$


$$\kappa\sigma^2 = C_4/C_2 \text{ [excess kurtosis]} \quad \text{measure of distribution's tails}$$






STAR, Phys. Rev. Lett. 128, 202303 (2022); arXiv : 2209.11940.
 Phys. Rev. Lett. 126, 092301 (2021); Phys. Rev. C 104, 024902 (2021)

Predicted Fluctuation in C_4/C_2 Near Critical Point




M. Stephanov. J. Physics G.: Nucl. Part. Phys. **38** (2011) 124147

- Non-monotonic energy dependence was observed in BES-I
- Recent measurement at 3 GeV demonstrates a return to the UrQMD baseline.
- High-statistics data (BES-II collider mode) with detector improvements have been taken from 7.7 GeV to 27 GeV.
- Data have been collected to fill the large gap between 3.0 and 7.7 GeV in STAR's Fixed-Target Program



STAR, Phys. Rev. Lett. 128, 202303 (2022); arXiv: 2209.11910
 Phys. Rev. Lett. 126, 092301 (2021); Phys. Rev. C 104, 024902 (2021)

Predicted Fluctuation in C_4/C_2 Near Critical Point



M. Stephanov. J. Physics G.: Nucl. Part. Phys. **38** (2011) 124147




- Non-monotonic energy dependence was observed in BES-I
- Recent measurement at 3 GeV demonstrates a return to the UrQMD baseline.
- High-statistics data (BES-II collider mode) with detector improvements have been taken from 7.7 GeV to 27 GeV.
- Data have been collected to fill the large gap between 3.0 and 7.7 GeV in STAR's Fixed-Target Program

Fixed-Target (FXT) Program at STAR

- Test run with gold target in 2015
- First physics runs at $\sqrt{s_{NN}} = 3.0$ GeV and 7.2 GeV in 2018
- Now have data at $\sqrt{s_{NN}}$ of 3.0, 3.2, 3.5, 3.9, 4.5, 5.2, 6.2, 7.2, and 7.7 GeV


Challenges for FXT

- Shifting asymmetric acceptance with respect to midrapidity
- At 7.7 GeV, the midrapidity moves to edge of Time Projection Chamber (TPC) acceptance
- Boost at higher energies shifts particle identification (PID) to rely more on TOF than TPC



ETOF Details

- CBM-TOF group provided ETOF system
- Provides particle identification over $1.55 < \eta < 2.2$
- Collected data for the Fixed-Target Program
- Calibrations completed at $\sqrt{s_{NN}} = 3.5, 3.9, 4.5$ GeV




Analysis Strategy



- Midrapidity analyses will be performed at $\sqrt{s_{NN}} = 3.2, 3.5, 3.9, 4.5$ GeV since we have near-full coverage over $-0.5 < y - y_{CM} < 0$ and $0.4 < p_T < 2$ GeV/c
- At $\sqrt{s_{NN}} = 5.2, 6.2, 7.2,$ and 7.7 GeV proton cumulants will be analyzed away from midrapidity
 - We can map proton cumulants as a function of μ_B

Top priority for publication →

Secondary for publication →



Expectations for Critical Point Location



From this year's 30th International Conference on Ultra-relativistic Nucleus-Nucleus Collisions (Quark Matter 2023)

Nominal v_s (GeV)	Chemical Potential μ_B (MeV)
3.2	697
3.5	666
3.9	632
4.5	589
5.2	541
6.2	487
7.2	443
7.7	420


Expectations for Critical Point Location



Nominal vs (GeV)	Chemical Potential μ_B (MeV)
3.2	697
3.5	666
3.9	632
4.5	589
5.2	541
6.2	487
7.2	443
7.7	420

From this year's 30th International Conference on Ultra-relativistic Nucleus-Nucleus Collisions (Quark Matter 2023)

- Wei-jie Fu: "Recent studies of QCD phase structure from both fRG and DSE have shown convergent estimate for the location of CEP: $600 \lesssim \mu_B \lesssim 650$ MeV"





Expectations for Critical Point Location



Nominal \sqrt{s} (GeV)	Chemical Potential μ_B (MeV)
3.2	697
3.5	666
3.9	632
4.5	589
5.2	541
6.2	487
7.2	443
7.7	420

From this year's 30th International Conference on Ultra-relativistic Nucleus-Nucleus Collisions (Quark Matter 2023)

- Wei-jie Fu: "Recent studies of QCD phase structure from both fRG and DSE have shown convergent estimate for the location of CEP: $600 \lesssim \mu_B \lesssim 650$ MeV"
- Maurício Hippert: holographic Bayesian analysis gives $560 \lesssim \mu_B \lesssim 625$ MeV





Expectations for Critical Point Location




Nominal \sqrt{s} (GeV)	Chemical Potential μ_B (MeV)
3.2	697
3.5	666
3.9	632
4.5	589
5.2	541
6.2	487
7.2	443
7.7	420

From this year's 30th International Conference on Ultra-relativistic Nucleus-Nucleus Collisions (Quark Matter 2023)

- Wei-jie Fu: “Recent studies of QCD phase structure from both fRG and DSE have shown convergent estimate for the location of CEP: $600 \lesssim \mu_B \lesssim 650$ MeV”
- Maurício Hippert: holographic Bayesian analysis gives $560 \lesssim \mu_B \lesssim 625$ MeV
- Jishnu Goswami: extrapolation using machine-learning model from hot QCD: $\mu_B \cong 600 \pm 80$ MeV




Jishnu Goswami, “Exploring the Critical Points in QCD with Multi-Point Padé and Machine Learning Techniques in (2+1)-flavor QCD” Quark Matter (2023)



Detector Acceptances




- We have near full acceptance in our analysis window ($-0.5 < y - y_{CM} < 0$, $0.4 < p_T < 2$ GeV/c) up to 4.5 GeV
- We rely on much more time-of-flight for particle identification for $\sqrt{s_{NN}} = 3.5, 3.9, 4.5$ GeV



Lessons: FXT and Timing Fluctuations




- STAR TPC has 40 μs drift time



Lessons: FXT and Timing Fluctuations




- STAR TPC has $40\ \mu\text{s}$ drift time \rightarrow occasionally a second collision will occur within that time



Lessons: FXT and Timing Fluctuations





- STAR TPC has 40 μs drift time \rightarrow occasionally a second collision will occur within that time
- Tracks from a second collision that are included in the event are known as pileup
- Time-of-flight (TOF) detectors use nanosecond-scale timing resolution to identify particles
- Pileup tracks which are out-of-time will often not be well-identified by TOF



Lessons: FXT and Timing Fluctuations





- When we use the time-of-flight for particle ID, protons from out-of-time pileup are not counted by the fast TOF
- The TPC still identifies all the pileup tracks
- Centrality is determined by the TPC multiplicity
- A pileup event may be classified as very central, but have few protons
- For each centrality bin, this leads to a low-proton-number tail



Lessons: FXT and Timing Fluctuations




- When we use the time-of-flight for particle ID, protons from out-of-time pileup are not counted by the fast TOF
- The TPC still identifies all the pileup tracks
- Centrality is determined by the TPC multiplicity
- A pileup event may be classified as very central, but have few protons
- For each centrality bin, this leads to a low-proton-number tail
- Long tails mean large kurtosis



Lessons: FXT and Timing Fluctuations





- We can simulate this in UrQMD by sampling two events at a rate of 0.0001
- In the experiment we use TOF PID for tracks above a certain momentum
- We can simulate this by only including pileup tracks with momenta below the threshold for using TOF
- The more TOF PID we use, the more the pileup causes a tail



Lessons: FXT and Timing Fluctuations



- We can simulate this in UrQMD by sampling two events at a rate of 0.0001
- In the experiment we use TOF PID for tracks above a certain momentum
- We can simulate this by only including pileup tracks with momenta below the threshold for using TOF
- The more TOF PID we use, the more the pileup causes a tail
- This causes instability in C_4/C_2 and other cumulants
- **Conclusion:** remove pileup when using TOF for proton ID. *Do not* correct for it



- Recent data from the Fixed–Target Program will extend our knowledge of the proton cumulant ratios at low energies (3.2-7.7 GeV)
- Non-monotonic variation in proton higher moments would suggest proximity to a critical point in the QCD phase diagram
- Many theoretical approaches now suggest critical point is accessible in the STAR Fixed-Target regime
- Midrapidity measurements will be performed at $\sqrt{s_{NN}} = 3.2, 3.5, 3.9, \text{ and } 4.5$ GeV
- Rapidity-dependent study will be done at $\sqrt{s_{NN}} = 5.2, 6.2, 7.2, \text{ and } 7.7$ GeV.
- The Fixed-Target analysis comes with unique challenges which we are working to understand

