

Measuring Away-Side Jet Modifications in Au+Au Collisions at RHIC

Kun Jiang

Purdue University
University of Science and Technology of China (USTC)

Outline




- Motivations
- Methodology
- Correct for detector efficiency/acceptance
- Systematic error study
- Results and discussion
- Summary


Motivation



- Energetic partons are predicted to lose energy due to interactions in the dense medium





PRC 82 (2010) 024912



- Measurements of medium modifications of jets have so far been obscured by the large anisotropic flow background. Flow shape and amplitude are not precisely known.
- We devise a method to subtract flow background using data itself



Methodology



close-region 2p corr.
= flow + near-side jet + away-side jet * fraction_close



far-region 2p corr.
= flow + near-side jet + away-side jet * fraction_far

- Select events with a large recoil P_x from a high- p_T trigger particle within a given eta window (0.5 - 1 or -1 - -0.5) to enhance away-side jet population
- Analyze di-hadron correlations in close-region and far-region respectively
- Flow contributions to close-region and far-region are equal!!

"Px" calculation

- Recoil Px from a high-pT trigger particle:


$$P_x |_{\eta_1 \leq \eta \leq \eta_2} = \sum_{assoc} pT_{assoc} \cdot \cos(\phi_{assoc} - \phi_{trig})$$



- For each centrality, cut on the left tail of the distribution (fraction of events) to enhance away-side jet population





Correct for ϕ -dependent efficiency * acceptance



- Normalize the single particle phi distribution to average unity. The inverse of that will be the phi-dependent efficiency
- Done run-by-run (and runs with same efficiency grouped together)
- Corrections are done as a function of centralities
- Apply phi-dependent efficiency correction for Px calculation

Correct for η -dependent efficiency * acceptance





- Use the $-2 < z_{vtx} < 2$ cm distribution and make it symmetric by taking the average of the two sides. Treat the symmetrized $dN/d\eta$ as "truth"
- Take the ratio of the $dN/d\eta$ in each z_{vtx} bin to this "truth". Use the inverse of the ratio as the η - and z_{vtx} -dependent correction
- Apply η -dependent efficiency correction for P_x calculation and di-hadron correlations

STAR Run11 AuAu 200 GeV



trigger particle: $p_T > 3$ GeV, associated particle: $1 < p_T < 2$ GeV, minbias
Near-side almost equal as expected




close-region 2p corr.
= flow + near-side jet + away-side jet * fraction_close

far-region 2p corr.
= flow + near-side jet + away-side jet * fraction_far

STAR TPC





Fit away-side jet shape with a Gaussian




Systematic study: Varying "Px" cut

varying Px cut percentage: 2%, 5%, 10% (default), 15%, 20%, 30%, 50%
 In principle: σ should not change with Px cut, only jet fraction (or amplitude) changes




- Take the ratio of the σ under other "Px" cut to σ under default cut
- The Sigma of the distribution gives a systematic error of 3.1% from "Px" cut





Systematic study: Varying track quality cuts



Default: $\text{dca} \leq 2$ $\text{nHitsFit} \geq 20$

Loose cut: $\text{dca} \leq 3$ $\text{nHitsFit} \geq 15$


Tight cut: $\text{dca} \leq 1$ $\text{nHitsFit} \geq 25$

dca: distance of closest approach to the collision vertex

nHitsFit: number of hits in the TPC

- Take the ratio of the σ under other track quality cuts to σ under default track quality cuts
- The Sigma of the distribution gives a systematic error of 3.8% from track quality cuts

Result: the width of the away-side jet




- Away-side jets are modified:
- \bullet Moderate to high p_T associated particles: broaden with increasing centrality
 - \bullet Low p_T associated particles: no change
 - \bullet In central collisions, particles of all p_T tending towards same distribution

The horizontal caps indicate the systematic error



Result: the width of the away-side jet



The leftmost 3 sets of data are for
PHENIX p+p
PHENIX d+Au
STAR d+Au
minbias

PRD 74 (2006) 072002
PRC 73 (2006) 054903

- Peripheral data are consistent with pp/dAu

Summary




- A novel method was devised to measure away-side jet modifications with clean, robust flow subtraction
- Away-side jets are modified:
 - Moderate to high p_T associated particles: broaden with increasing centrality
 - Low p_T associated particles: no change
 - In central collisions, particles of all p_T tending towards same distribution
- Potentially powerful method to study jet modification in medium

Backup slides



STAR detector






Data sets and cuts


- Data sets
 - Au+Au@200 GeV run11
- Event and track cuts
 - $|Vz| < 30$ cm
 - $Vr < 3$ cm
 - track quality cut > 0.52
 - nHitsFit > 20
 - dca < 2 cm

Run by run phi-dependent efficiency correction




- Histogram single particle phi distribution in one run
- Take a ratio between adjacent runs and fit by a constant
- Combine runs with same detector efficiency into one run block

#1




a jump in efficiency, grouped into 2 groups

#2







single bad run

two groups can be catenated



- Total 1297 runs in Run11. Grouped into 144 groups
- Exclude 134 single bad buns

Correct for detector inefficiency









Varying "Px" cut

study sigma vs Px cut percentage: 2%, 5%, 10%, 15%, 20%, 30%, 50%

default "Px" cut
10%



associate pT
0.5 - 1 GeV









Varying "Px" cut

study sigma vs Px cut percentage: 2%, 5%, 10%, 15%, 20%, 30%, 50%

default "Px" cut
10%



associate pT
1 - 2 GeV









Varying "Px" cut

study sigma vs Px cut percentage: 2%, 5%, 10%, 15%, 20%, 30%, 50%

default "Px" cut
10%




associate pT
2 - 3 GeV






Px distributions




Px distribution
estimated flow distribution

peripheral 60-70%



midcentral 30-40%




central 0-5%


- recoil Px is a convolution of away-side jet and flow
- assume that flow has a symmetric distribution, small px are almost from flow contribution
- flip the histogram on the right of the maximum bin to the left to estimate the flow contribution



Jet fractions



shape comparison: jet and far-region corr.



- For far-region corr, low p_T is dominated by flow. High p_T is dominated by jet, so it has same shape as jet.