


Supported in part by



# Global polarization of $\Xi$ hyperons in Au+Au collisions in the STAR experiment

## Outline:


- Introduction
- Global hyperon polarization
- Motivation
- The STAR experiment
- Hyperon polarization measurements
- Results
- Conclusions

Egor Alpatov (for the STAR Collaboration)


National Research Nuclear University MEPhI

# Introduction

- The Quark-Gluon Plasma (QGP) formed in non-central nucleus-nucleus collisions is associated with large angular momentum, that leads to vorticity in the medium
- Spin-orbit coupling aligns spin directions of produced particles with the direction of vorticity
  - Z.-T. Liang and X.-N. Wang, PRL94, 102301 (2005)
  - S. A. Voloshin, arXiv:nucl-th/0410089
- Another possible source of particle polarization is magnetic field, created in non-central collisions in the initial stage
  - D. Kharzeev, L. McLerran, and H. Warringa, Nucl.Phys.A803, 227 (2008)
  - McLerran and Skokov, Nucl. Phys. A929, 184 (2014)



# Vorticity



- In non-central HIC the initial collective longitudinal flow velocity depends on x:



$$\omega_y = \frac{1}{2} (\nabla \times v)_y \approx -\frac{1}{2} \frac{dv_z}{dx}$$

- For small polarization:

Becattini, Karpenko, Lisa, Upsal, Voloshin PRC95.054902 (2017)

$$P_\Lambda \approx \frac{1}{2} \frac{\omega}{T} + \frac{\mu_\Lambda B}{T}$$

$$P_{\bar{\Lambda}} \approx \frac{1}{2} \frac{\omega}{T} - \frac{\mu_\Lambda B}{T}$$



# How to measure global polarization?


- Hyperons are “self-analyzing” due to weak decay properties:
  - Daughter baryons are preferentially emitted along parent spin direction
- Daughter baryons of hyperons with polarization ( $\vec{P}$ ) follows the distribution:

$$\frac{dN}{d\Omega^*} = \frac{1}{4\pi} (1 + \alpha_H |\vec{P}| \cdot \widehat{p}_b^*) = \frac{1}{4\pi} (1 + \alpha_H P \cos \theta^*)$$


- $\alpha_H$  - decay parameter, unique for each hyperon species
- $\widehat{p}_b^*$  is the daughter baryon momentum in the parent frame
- Projection to the transverse plane can be measured:


$$P_H = \frac{8}{\pi \alpha_H} \frac{\langle \sin(\psi_1 - \varphi_p^*) \rangle}{Res(\psi_1)}$$

- $\psi_1$  is the reaction plane angle
- $\psi_1$  and its resolution  $Res(\psi_1)$  can be calculated with spectator’s signal.
- $\Xi$  global polarization could also be measured via its daughter  $\Lambda$  polarization with transfer factor  $C_{\Xi\Lambda} = 0.932$



STAR *Nature* 548, 62–65 (2017)







# Motivation

- Global polarization of  $\Lambda$  hyperons was measured for  $\sqrt{s_{NN}} = 7.7-200$  GeV at STAR
- $P_H$  decreases with increasing collision energy
- Difference between  $P_\Lambda$  and  $P_{\bar{\Lambda}}$  maybe due to B-field effect
- Theoretical calculations can quantitatively explain the energy dependence of the  $\Lambda$  polarization, but many of them fail to explain differential measurements
- Nowadays there is a growing interest to measure the global polarization of other hyperons such as  $\Xi$ .
- $\Xi$  polarization may provide new input for global polarization and vorticity studies

STAR PRC 98, 014910 (2018)



# The STAR experiment



## Hyperon reconstruction:

- Time Projection Chamber  
 $|\eta| \in [-1, 1]$
- Time-Of-Flight  
 $|\eta| \in [-0.9, 0.9]$

## Event plane angle measurement:

- Beam-Beam Counter  
 $|\eta| \in [3.3, 5.0]$
- Event-Plane Detector  
 $|\eta| \in [2.1, 5.1]$
- Zero Degree Calorimeter  
 $|\eta| > 6.3$

# Experimental technique




- **Event plane**  $\Psi_1$  be measured by detectors at forward rapidity where directed flow is large

$$\Psi_1 = \tan^{-1} \left( \frac{\sum w_i \sin(\phi_i)}{\sum w_i \cos(\phi_i)} \right), \text{ where } w_i \text{ is detector's tile ADC}$$



- $Res(\Psi_{1,Forward} \eta) = Res(\Psi_{1,Backward} \eta) = \sqrt{\langle \cos(\Psi_{1,Forward} \eta - \Psi_{1,Backward} \eta) \rangle}$
- BBC and ZDC for  $\sqrt{s_{NN}} = 54.4$  GeV, EPD and BBC for  $\sqrt{s_{NN}} = 27$  GeV

A. M. Poskanzer, S. A. Voloshin, PRC58.1671(1998)

- **Hyperon reconstruction** performed via decay topology
- $\Lambda$  daughters identified via TPC and TOF
- $\Xi$  were reconstructed via  $\Xi \rightarrow \Lambda + \pi$



# Global polarization: event-plane method




- Build invariant mass distribution for each  $(\Psi_1 - \varphi^*)$  bin
- Subtract background and integrate subtracted distribution
- Fit  $dN/d(\Psi_1 - \varphi^*)$  with Fourier function up to 2<sup>nd</sup> order
- Coefficient for  $\sin(\Delta\varphi^*)$  term is what we're looking for

$$\Delta\varphi^* = \Psi_1 - \varphi^*$$




# Results



- Global polarization of  $\Xi$  hyperons consistent with model predictions
- Experimental results for  $\Xi$  global polarization congruence with each other at different collision energies
- $\Xi$  global polarization, measured via its daughter  $\Lambda$  decays is consistent with direct method
- $\Xi$  and inclusive  $\Lambda$  global polarization are consistent within statistical uncertainties

## Global hyperon polarization $\sqrt{s_{NN}}$ dependence



# Conclusions

- We presented first results of  $\Xi + \bar{\Xi}$  global polarization measurements in Au+Au collisions at 54.4 GeV
- $\Xi + \bar{\Xi}$  global polarization is comparable to  $\Lambda + \bar{\Lambda}$  global polarization within uncertainties
- We are looking forward to continuing this measurements at other energies

Thank you for your attention!