

## Abstract

The observation of hyperon global polarization along the system's angular momentum has revealed the existence of large vorticities in the medium created by heavy-ion collisions. Using the high-statistics data collected by the STAR experiment during the RHIC Beam Energy Scan II (BES-II) program with upgraded detector systems, we present the global polarization measurements of  $\Xi$  and  $\Omega$  hyperons in Au+Au collisions at BES-II energies ( $\sqrt{s_{NN}} = 7.7, 9.2, 11.5, 14.6, 17.3, 19.6$ , and 27 GeV). Specifically, we focus on the polarization behaviors observed in different hyperons ( $\Lambda, \Xi, \Omega$ ). These results provide new insights into the polarization mechanism and vorticity fields in heavy-ion collisions as well as additional constraints on the properties and dynamics of the hot and dense matter created in these collisions.

## Introduction

The polarization along the initial angular momentum direction, can be described as [1,2]


$$P_H = \frac{8}{\pi a_H A_0} \frac{(\sin(\Psi_1 - \phi_B^*))}{\text{Res}(\Psi_1)}$$

$\Psi_1$ : first-order event plane  
 $a_H$ : hyperon decay parameter  
 $\phi_B^*$ : azimuthal angle of the daughter baryon in hyperon rest frame  
 $A_0$ : acceptance correction factor

## Motivation


Multi-strange hyperon global polarization properties

- Collision energy, centrality,  $p_T, \eta$  dependence?
- Possible  $\Lambda, \Xi, \Omega$  global polarization difference?




## Global Polarization Signal Extraction

- Direct measurement : via daughter  $\Lambda$  angle distribution in  $\Xi, \Omega$  rest frame
- Indirect measurement : via daughter  $\Lambda$  polarization with spin transfer factor ( $C_{\Xi \rightarrow \Lambda} \sim 0.944, C_{\Omega \rightarrow \Lambda} \sim 1.0$  is assumed)
- Generalized Invariant Mass method(IM) for signal extraction  
 $(\sin(\Psi_1 - \phi_B^*)) = (1 - f^{Bg}(M_{inv}))(\sin(\Psi_1 - \phi_B^*))^{Sig} + f^{Bg}(M_{inv})(\sin(\Psi_1 - \phi_B^*))^{Bg}$   
 $(\sin(\Psi_1 - \phi_B^*))^{Sig} = (\sin(\Psi_1 - \phi_B^*))^{true} + c \sin(\phi_H - \phi_B^*)$   
 $f^{Bg}(M_{inv})$  is background fraction as a function of invariant mass,  $c$  is  $v_1$  factor



## The STAR Detector



### Time Projection Chamber

- Upgrade with inner TPC
- Better track quality
- Larger acceptance
- $|\eta| < 1.0 \rightarrow |\eta| < 1.5$


### Time Of Flight

- PID via particle velocity
- $|\eta| < 0.9$


### Event Plane Detector

- Event plane reconstruction
- $2.1 < |\eta| < 5.1$


## Results



## Hyperon and Event Plane Reconstruction



- $\Lambda, \Xi, \Omega$  hyperons reconstructed using KF Particle package



- First-order event plane reconstructed by EPD
- Event plane correction:  $\eta$ -weight,  $\phi$ -weight, recenter, flattening
- Two-sub event plane method to estimate resolution with Bessel function

- Significant  $\Xi^- + \Xi^+$  global polarization observed ( $\sim 5\sigma$ )
- Global polarization of  $\Xi^- + \Xi^+$  and  $\Omega^- + \Omega^+$  seems to decrease with increase in collision energy
- $\Xi^- + \Xi^+$  global polarization are consistent between direct and indirect measurement methods
- No significant difference between  $\Lambda + \bar{\Lambda}$  and  $\Xi^- + \Xi^+$  global polarization within uncertainties
- A hint of larger  $\Omega^- + \Omega^+$  polarization than  $\Lambda + \bar{\Lambda}$  and  $\Xi^- + \Xi^+$  in lower energies

## Summary

- The first measurement of  $\Xi^- + \Xi^+$  and  $\Omega^- + \Omega^+$  global polarization vs. collision energy at  $\sqrt{s_{NN}} = 7.7, 9.2, 11.5, 14.6, 17.3, 19.6$ , and 27 GeV
- Global polarization of  $\Xi^- + \Xi^+$  and  $\Omega^- + \Omega^+$  seems to decrease with collision energy, with a hint of larger  $\Omega^- + \Omega^+$  polarization

## References

- [1] Z.-T. Liang and X.-N. Wang, PRL 94, 102301 (2005)
- [2] STAR Collaboration, Nature 548, 62 (2017).
- [3] STAR Collaboration, PRL 126, 162301(2021)
- [4] Hui Li et al., PLB 827, 136971(2022)

