

Status of the Analyses for the Proton Higher-Order Fluctuations in the STAR Fixed-Target Program from $\sqrt{s_{NN}} = 3.2$ to 7.7 GeV

1. Introduction

Critical Fluctuations

- Fluctuations in conserved charges (such as baryon number) expected near critical point
- Proton number is a proxy for baryon number
- Measure cumulants of net-proton number distributions as a function of collision energy
- QCD calculations predict non-monotonic cumulants vs $\sqrt{s_{NN}}$ if close to critical point

Cumulants and Moments

Cumulants of a distribution are defined as:



- $C_1 = \langle N \rangle \equiv \mu$
- $C_2 = \langle (N - \mu)^2 \rangle \equiv \sigma^2$
- $C_3 = \langle (N - \mu)^3 \rangle$
- $C_4 = \langle (N - \mu)^4 \rangle - 3\langle (N - \mu)^2 \rangle^2$

Some standardized moments are:

- $S\sigma = C_3/C_2$ [measures asymmetry]
- $\kappa\sigma^2 = C_4/C_2$ [measures "tailedness"]

Previous Results


- Non-monotonicity in Beam Energy Scan I net-proton C_4/C_2 with 3.1σ significance
- Recent $\sqrt{s_{NN}} = 3$ GeV measurement returns to non-critical baseline
- High-statistics collider data from Beam Energy Scan II will improve measurements from $\sqrt{s_{NN}} = 7.7-27$ GeV
- Fixed-target data will fill the gap between $\sqrt{s_{NN}} = 3$ and 7.7 GeV



2. The STAR Fixed-Target Program

- Relativistic Heavy-Ion Collider (RHIC) can only collide Au+Au down to $\sqrt{s_{NN}} = 7.7$ GeV
- Gold target installed on west end of STAR TPC
- New data at $\sqrt{s_{NN}} = 3.0, 3.2, 3.5, 3.9, 4.5, 5.2, 6.2, 7.2, 7.7$ GeV

$\sqrt{s_{NN}}$ (GeV)	3.2	3.5	3.9	4.5	5.2	6.2	7.2	7.7
μ_B (GeV)	0.70	0.67	0.63	0.59	0.54	0.49	0.44	0.42
y_{CM}	1.139	1.254	1.375	1.522	1.683	1.867	2.021	2.102
# good evts produced	201M	116M	117M	108M	103M	118M	317M	113M
	yes	yes	yes	yes	yes	yes	no	yes




Challenges

- Acceptance is asymmetric w.r.t. midrapidity
- Limited midrapidity acceptance at highest energies
- Time-of-flight (TOF) required for particle ID


4. Timing Fluctuations

- Fixed-target analysis uses TOF and TPC for proton ID
- Centrality determined by TPC multiplicity
- TOF relies on precision timing and often misses protons from pileup with timing offset
- TPC multiplicity includes pileup tracks
- Pileup events classified as central collisions but have few TOF-identified protons
- This causes tail on proton-number distribution, which creates false signal for high-order cumulants



Solution: Reject pileup, do not correct for it. Usual cumulant correction for pileup doesn't fix timing fluctuations


3. Acceptance



- Standard analysis window shown in red box
- We have near-full acceptance up to $\sqrt{s_{NN}} = 4.5$ GeV
- Top energy analyses must be away from midrapidity

Overlap at $\sqrt{s_{NN}} = 7.7$ GeV

- Data was taken at $\sqrt{s_{NN}} = 7.7$ GeV in both the collider and fixed-target configurations
- There is significant overlap in acceptance for a comparison
- Overlap analysis will validate fixed-target and collider methodologies



References

- [1] Xiaofeng Luo and Nu Xu. In: *Nuclear Science and Techniques* 28 (Aug. 2017).
- [2] M. S. Abdallah et al. In: *Phys. Rev. Lett.* 128 (20 2022), p. 202303.
- [3] M A Stephanov. In: *Journ. of Phys. G: Nucl. and Part. Phys.* 38.12 (2011), p. 124147.

5. Conclusions

- Fixed-Target Program extends energy range of RHIC down to $\sqrt{s_{NN}} = 3$ GeV
- Fixed-target proton fluctuations analyses are underway from $\sqrt{s_{NN}} = 3.2-7.7$ GeV
- Overlap acceptance for fixed-target and collider data at $\sqrt{s_{NN}} = 7.7$ GeV allows methodology validation
- Reliance on TOF for particle ID makes measurements sensitive to timing fluctuations, which can be managed
- Prior measurements show non-monotonic variation in C_4/C_2 as a function of $\sqrt{s_{NN}}$ at 3.1σ significance
- Fixed-target data will add mid-rapidity C_4/C_2 measurements at $\sqrt{s_{NN}} = 3.2-4.5$ GeV to energy scan